

On the conversion of module representations for higher dimensional supersingular isogenies

AUREL PAGE¹, DAMIEN ROBERT¹, AND JULIEN SOUMIER²

ABSTRACT. We expand the well developed toolbox between quaternionic ideals and supersingular elliptic curves into its higher dimensional version, namely (Hermitian) modules and maximal supersingular principally polarized abelian varieties. One of our main result is an efficient algorithm to compute an unpolarized isomorphism $A \simeq E_0^g$ given the abstract module representation of A . This algorithm relies on a subroutine that solves the Principal Ideal Problem (PIP) in matrix rings over quaternion orders, combined with a higher dimensional generalisation of the Clapotis algorithm. To illustrate the flexibility of our framework, we also use it to reduce the degree of the output of the KLPT² algorithm, from $O(p^{25})$ to $O(p^{15.5})$.

1. INTRODUCTION

As a serious candidate for post-quantum cryptography standardization due to its compactness [AAA+25], isogeny-based cryptography has seen a lot of algorithmic improvements over the last decade.

A major tool in this context is the Deuring correspondence [Ler25; DKLPW20]. It allows to represent supersingular elliptic curves as quaternion orders, as soon as their endomorphism rings are known. Efficient algorithms to convert between ideals and isogenies are crucial backbones both for instantiating cryptographic protocols, such as SQISign [DKLPW20; DLRW24; BDF+24], but also for security reductions [EHLMP18; Wes21; HW25]. The first algorithm in this context was the KLPT algorithm [KLPT14], which allows to (heuristically) smoothen an ideal. The fall of the SIDH protocol in [CD23; MMPPW23; Rob23] led to the consideration of higher dimensional isogenies, and the discovery of the Clapotis algorithm [PR23] that uses higher dimensional isogenies as “shortcuts” for the ideal to isogeny conversion problem. Clapotis has proven polynomial time complexity, without Generalized Riemann Hypothesis (GRH), but in practice faster heuristic variants are used [PPS25; DEF+25; BCE+25; DEIV26].

A recent trend in isogeny based cryptography has been to consider the higher dimensional maximal supersingular graphs for their own sake, rather than just as providing convenient speed ups for computing dimension 1 isogenies. See for instance [CDS20; KMM+25] for a generalisation of the CGL Hash function [CLG09] in higher dimension. More recently [Rob24] proposed MIKE, a new non-interactive key exchange that makes crucial use of the dimension 2 and 4 isogeny graphs.

There are two slightly different versions of the Deuring correspondence. The first one, as mentioned represents a supersingular curve $E/\mathbb{F}_{p^2}$ by its endomorphism ring $\mathcal{O} = \text{End}(E)$, as a maximal order in $\mathcal{B}_{p,\infty}$, the quaternion algebra ramified at

¹CENTRE INRIA DE L’UNIVERSITÉ DE BORDEAUX

²UNIVERSITÉ DE LORRAINE, CNRS, INRIA

p and ∞ . The second one is to fix a base curve $E_0/\mathbb{F}_{p^2}$ with endomorphism ring $\mathcal{O}_0$, and represent E by the left $\mathcal{O}_0$ -module $\text{Hom}_{\mathbb{F}_{p^2}}(E, E_0)$. In fact, if $\varphi: E_0 \rightarrow E$ denotes any isogeny between E_0 and E , the latter module induces a left $\mathcal{O}_0$ -ideal $I = \text{Hom}_{\mathbb{F}_{p^2}}(E, E_0)\varphi$. Given I , $\mathcal{O} = \text{End}(E)$ is the right order of I . Conversely given $\mathcal{O}$, we can recover an ideal equivalent to I , through any connecting ideal between $\mathcal{O}_0$ and $\mathcal{O}$. (A slight subtlety is that there are two classes of such connecting ideals, I and $\mathfrak{p}I$ where $\mathfrak{p}$ is the two-sided ideal of $\mathcal{O}_0$ of norm p . This is because the endomorphism ring representation does not distinguish between E and its $\mathbb{F}_{p^2}/\mathbb{F}_p$ -Galois conjugate E^σ).

These two versions of the Deuring correspondance extend to higher dimension. We can represent a g -dimensional maximal, *i.e.*, $\#A(\mathbb{F}_{p^2}) = (p+1)^{2g}$, supersingular principally polarized abelian variety (ppav) (A, λ_A) by its polarized endomorphism ring $(\text{End}(A), r_A)$ where r_A is the Rosati involution. We can also use the module representation $M = \text{Hom}(A, E_0)$, endowed with the unimodular Hermitian form H induced by the principal polarizations: $H(\varphi_1, \varphi_2) = \varphi_1 \circ \lambda_A^{-1} \circ \varphi_2^\vee \circ \lambda_0 \in \mathcal{O}_0$, where λ_0 is the canonical principal polarization on E_0 , and $\varphi_2^\vee$ the dual of φ_2 . This latter representation due to Serre, was introduced in the context of higher dimensional isogeny based cryptography in [Rob24].

We note that there is a phase shift compared to the dimension 1 case: in dimension 1 there are many maximal orders $\mathcal{O}$ in $\mathcal{B}_{p,\infty}$, which all admit a unique positive Rosati involution (given by the quaternionic conjugation). From the ideal point of view, there are many classes of ideals (up to isomorphism as a $\mathcal{O}_0$ -module), which all admit a unique principal polarization (*i.e.* a unimodular Hermitian form) $H_I(x, y) = x\bar{y}/N(I)$. On the geometric side, this corresponds to the fact that the Néron-Severi group satisfies $\text{NS}(E) = \mathbb{Z}$, and all elliptic curves have a canonical principal polarization, *i.e.* the one induced by (0_E) . So the polarization lies in the background in dimension 1. In higher dimension $g > 1$, Deligne¹ proved that a maximal superspecial abelian variety $A/\mathbb{F}_{p^2}$ is always isomorphic to E_0^g . As a consequence, the endomorphism ring $\text{End}(A)$ of A is isomorphic to the ring of matrices $\mathcal{M}_g(\mathcal{O}_0)$, and the module $M = \text{Hom}(A, E_0)$ is isomorphic to $\mathcal{O}_0^g$. So this time, it is crucial to keep track of the principal polarization λ_A of A , either as the Rosati involution for the endomorphism representation, or as an Hermitian form in the module representation.

In particular, we can represent (A, λ_A) by a Hermitian unimodular definite matrix $S \in \mathcal{M}_g(\mathcal{O}_0)$, whose associated Hermitian form on $\mathcal{O}_0^g$ is $H_S(x, y) = xSy^*$, and whose associated Rosati involution on $\mathcal{M}_g(\mathcal{O}_0)$ is SM^*S^{-1} , where $M \mapsto M^*$ is the conjugate transpose. This is the “IKO representation”, used in [CDK+25] in their (heuristic) generalisation of the KLPT algorithm to dimension 2.

In the following, we frequently use the KLPT algorithm as a black box. Although the original analysis of its efficiency relied on heuristic assumptions [KLPT14], a provably polynomial-time version was later established under GRH in [Wes21]. Throughout this work, “KLPT” will implicitly refer to this latter version, even though it outputs larger ideals, so that the original KLPT is often used in practice.

1.1. Main contributions. We give a brief summary of our main contributions, and refer to the section below for more details. Our main algorithmic tools are two fold:

¹Deligne’s theorem is over $\overline{\mathbb{F}_p}$, but [JKP+18] gives a version over $\mathbb{F}_{p^2}$

- A polynomial time algorithm (under GRH) to solve the Principal Ideal Problem in a matrix ring over a maximal quaternion order $\mathcal{O}_0$ of $\mathcal{B}_{p,\infty}$,
- A polynomial time algorithm that, given a unimodular Hermitian module (M, H) of rank g , constructs an efficient representation of a double path between E_0^g and the ppav (A, λ_A) associated with (M, H) .

As applications of these algorithmic results (and of further tools developed in this paper), we derive:

- Polynomial time algorithms to convert between different abstract representations of a maximal supersingular abelian variety (A, λ_A) : the Hermitian module representation, the endomorphism ring representation (along with the associated Rosatti involution). In this text we prefer to work with rank g modules over $\mathcal{O}_0$ rather than ideals of $\mathcal{M}_g(\mathcal{O}_0)$, but see Proposition 3.5 for how to go from one to the other.
- Algorithms to convert from these abstract representations into concrete representations, and conversely. Notably, given any module morphism $\psi : (M_2, H_2) \rightarrow (M_1, H_1)$ between two unimodular torsion free Hermitian modules, we can build an explicit representation of the associated principally polarised abelian varieties and morphism (which need not be an isogeny): $\phi : (A_1, \lambda_1) \rightarrow (A_2, \lambda_2)$.
- As an illustration of our tools we derive improved bound for the KLPT² algorithm from [CDK+25].

We note that although our results extend some of the security reductions in $g = 1$ (see e.g. [EHLMP18; Wes21; HW25]), the security picture is not yet as complete as in the $g = 1$ case (for instance we do not provide an equivalent of the OneEnd to EndRing reduction of [PW24]).

Although Clapotis was the first algorithm allowing the computation of the ideal class group action of an arbitrary ideal in polynomial time, it required a lot of improvements to be made practical (see [PPS25; DEF+25; DEIV26]). Likewise, in this paper, while we do provide an HD generalisation of Clapotis, which in theory allows us to evaluate the module action of [Rob24] for arbitrary modules, more work is needed for the algorithm to really be practical.

1.2. Organization of the paper. Section 2 is dedicated to the introduction of mathematical background.

Section 3 presents what we call the abstract and standard representations for polarized abelian varieties of dimension $g \geq 2$. On the one hand, an *abstract* representation informally refers to a description in which only the underlying algebraic structure of the mathematical objects is specified. For example, an abstract Hermitian module representation (M, H_M) consists of a finitely presented torsion-free module M , given by a finite set of generators, together with a Hermitian form H_M represented by a Gram matrix. On the other hand, we define the notion of a *standard* representation. Such representations are abstract representations of a particular canonical form; for instance, in the module setting, one may require $M = \mathcal{O}_0^g$. A standard representation may therefore be viewed as an analog of a normal form within a given equivalence class.

We then explain how to convert between the different abstract and standard representations. In particular, we show that computing isomorphisms between $\text{End}(A)$ (resp. M) and $\mathcal{M}_g(\mathcal{O}_0)$ (resp. $\mathcal{O}_0^g$) reduces to solving the Principal Ideal

Problem in $\mathcal{M}_g(\mathcal{O}_0)$. To that end, we provide a new algorithm (see Corollary 3.8) to build an explicit isomorphism between two isomorphic central simple algebras A, A' over $\mathbb{Q}$, given maximal orders $\Lambda \subset A$ and $\Lambda' \subset A'$. This algorithm builds on [IRS12], but uses the extra information given by Λ, Λ' to bypass the need for a factorization oracle.

In Section 4 we present a polynomial time solution to solve the principal ideal problem in matrix rings over maximal quaternion orders. This allows us to conclude in Section 4.4 that we provide efficient algorithms to convert between the abstract and standard representations.

Then in Section 5, we develop very general frameworks that will allow us to go back and forth between explicit and abstract representations of abelian varieties and their morphisms. Informally, the *explicit* representations also grasps the geometrical properties of the mathematical objects involved. For instance, by an explicit representation of an isogeny $\varphi: A \rightarrow B$, we mean that A and B are given in a convenient way (*e.g.* by level 4 theta null points), and that we can efficiently evaluate φ .

We explain first, given an explicit representation of morphisms $\varphi_1, \dots, \varphi_m \in \text{Hom}(A_1, A_2)$ that generates it (as an abelian group), how to recover its representation as an abstract abelian group. Given also explicit representations of generators of $\text{End}(A_1), \text{End}(A_2)$, we also explain how to recover their ring structure and the $(\text{End}(A_2), \text{End}(A_1))$ -bimodule structure of $\text{Hom}(A_1, A_2)$. For this, we use the same method as already used in dimension 1 for endomorphism rings [EHLMP18; Wes21], namely we embed $\text{Hom}(A_1, A_2)$ into $\mathbb{Z}^m$ thanks to the trace pairing $(\varphi_1 | \varphi_2) = \text{Tr}(\varphi_1 \tilde{\varphi}_2)$. The trace pairing method was already used in higher dimension to compute, from an explicit representation, an abstract representation of endomorphisms rings, see [ABIGW25, § 3], and [Mon26] when $g = 2$. We show that the method works just as well for Hom modules than for endomorphism rings. As an aside, both references above assume that they are given an explicit representation not only of the $\varphi_i \in \text{End}(A)$, but also of their polarized duals $\tilde{\varphi}_i$, so as to be able to compute the action of the $\varphi_i \tilde{\varphi}_j$ in order to compute their traces. In this paper we sidestep this by using the Weil pairing to recover from the action of the φ_i on the ℓ -torsion, the action of its polarized dual $\tilde{\varphi}_i$. The Weil pairing trick is folklore in isogeny based cryptography (see for instance [Wes22, § 3.1]), and works just as well for general morphisms.

For the other direction, in Proposition 5.16 we show how double paths allow to compute an explicit representation of an homomorphism set in polynomial time, given its abstract representation. A summary of the translations between the different representations may be found in Appendix A.

Motivated by the previous paragraph, in Section 6 we develop the technical tools needed to compute the double paths using Hermitian module theory, leading to an higher dimensional version of the Clapotis algorithm. That such an extension was possible was conjectured in [Rob24]. Given that the module action constitutes a higher-dimensional analog of the class group action, our algorithm computing this action may be regarded as a corresponding generalization of the classical Clapotis family of algorithms.

Our generalised Clapotis (HD-Clapotis for Higher Dimensional Clapotis ²) allows to make our abstract representations effective. For instance given the module representation (M, H_M) of (A, λ_A) , we can recover A (e.g. as given by theta constants) along with the explicit action of $M = \text{Hom}(A, E_0)$ and $\text{End}(A) \simeq \text{End}_{\mathcal{O}_0}(M)$ on A . We mean by *explicit action*, that we can evaluate the corresponding morphisms on points of A . The subscript $\mathcal{O}_0$ emphasizes that M is viewed as a $\mathcal{O}_0$ -module, and we may drop it when it will be clear from the context. Other applications of the HD-Clapotis algorithm, such as the computation of (non polarized) isomorphism $A \simeq E_0^g$, are summarized in Section 6.3.

To illustrate the flexibility of our algorithms, in Section 7 we revisit the KLPT² algorithm in dimension 2 from [CDK+25] and show how the module point of view combined with the tools developed above allows to improve the length of the resulting isogeny from $O(p^{25})$ to $O(p^{15.5})$.

All our algorithms are in classical polynomial time in $\log(p)$. They are mostly deterministic, except that every time we use the HD representation, such as in the HD-Clapotis algorithm, we need to express an integer as a sum of four squares, which is only known to be randomized polynomial time [PT18]. Our PIP algorithm relies on the KLPT algorithm and therefore can only be proven to run in polynomial time under GRH. It also depends on Algorithm 1, which is randomized. Finally our improvements to KLPT² are heuristic (it uses the same heuristics as in [CDK+25]).

We provide a proof of concept implementation of our algorithm for solving the Principal Ideal Problem in $\mathcal{O}_0^g$, available at:

<https://gitlab.inria.fr/module-representation/PIP-quaternion-mat> .

An explicit example may be found in Appendix B.

1.3. Concurrent work. In an independent unpublished work by Castryck, Eriksen, Invernizzi, and Vercauteren [CEIV26], the authors also solve the Principal Ideal Problem in the matrix ring $\mathcal{M}_g(\mathcal{O}_0)$, using a different approach. A noteworthy difference is that they do not rely on the KLPT algorithm as a subroutine: this makes their output smaller. On the other hand, their algorithm is only heuristically in polynomial time, while ours is proven to be polynomial time under GRH.

In another independent work, made recently available in [Mon26], Montessinos also explains how to switch, in dimension 2, between the different endomorphism ring representations above (abstract, effective, IKO), and how to compute an explicit isomorphism $A \simeq E_0^2$. His approach is complementary to ours: to build this isomorphism out of an IKO representation of A , he uses the (heuristic) KLPT² algorithm to build a 2^e -isogeny $E_0^2 \rightarrow A$, and then he carefully iteratively builds isomorphisms $E_0^2 \simeq A_i$ at each step of the 2^e -isogeny chain. In this paper, we instead rely on the HD-Clapotis algorithm to construct a double path $A \rightarrow E_0^2$, which also allows us to explicitly compute the isomorphism.

A difference with [Mon26] is that in this paper we mainly adopt the module point of view, which is not used in [Mon26]. Thanks to the HD-Clapotis algorithm, available in any dimension, this allows us to treat any dimension g , and also to handle the missing reduction between Problem 3.3 and Problem 3.4 in [Mon26] (which is only treated for polarized products of two elliptic curves there), see Remark 5.10.

²Following feedback from the isogeny community regarding the “tradition” of naming Clapotis-style IdealToIsogeny algorithms with variants of /klapoti/, and following a suggestion of Houben, we also propose the name *Chlapotis*

Both papers have benefited from each other. In [Mon26], Montessinos uses our solution to the PIP to find the IKO matrix. On the other hand, Montessinos pointed out to us a crucial ambiguity that we had missed in a preliminary draft version of this paper. Indeed, although all automorphisms of $\mathcal{M}_g(\mathcal{B}_{p,\infty})$ are inner automorphisms, this is no longer the case for $\mathcal{M}_g(\mathcal{O}_0)$, where there is one extra outer automorphism (see [Mon26, § 6.1.1] for more details). Thus the endomorphism ring representation $(\text{End}(A), r_A)$, where r_A is the Rosati involution, has an ambiguity. As explained in the introduction, this ambiguity is already present in dimension 1. Montessinos solves this ambiguity by keeping track of the action by $\text{End}(A)$ on global differentials of A . In this paper we follow his lead and use the same solution. (Although it may seem that our notion of refined endomorphism ring representation in Remark 3.2 does not involve the global differentials, by Proposition 5.11 hidden in the module representation.)

1.4. History. This work started as a follow up of the paper [GSS25] by Gaudry, Soumier, and Spaenlehauer. In that paper, the authors explain how to compute an explicit isomorphism $E_1 \times E_2 \times \cdots \times E_g \simeq E'_1 \times E'_2 \times \cdots \times E'_g$ between products of maximal supersingular curves over $\mathbb{F}_{p^2}$, given the endomorphism rings of the E_i, E'_i .

In a private discussion with these authors, the second author of this paper had sketched out an *heuristic algorithm* on how to compute an isomorphism $A \simeq E_0^g$ for a general maximal supersingular abelian variety $A/\mathbb{F}_{p^2}$ (which exists by Deligne's theorem), given the *module representation* M of A , as introduced in [Rob24].

This algorithm relied on two crucial ingredients. First we needed to solve the purely number theoretic problem of computing an isomorphism $\psi: M \simeq \mathcal{O}_0^g$. For this we turned to an algorithm due to the first author, who in 2014 wrote a note on how to extend the KLPT principalization algorithm to any central simple algebra. Unfortunately this note was never published. Also, and unlike the original KLPT algorithm, that algorithm is at best subexponential in general. However, in our context, the first author argued that it could be made polynomial time. In this paper we present a drastically simplified version of this general algorithm, adapted to our special case and using the module point of view. This simplified version was fully worked out, along with an implementation, by the third author. A fun fact that we realised when writing this paper is that the PIP for $\mathcal{M}_g(\mathcal{O}_0)$ had already been implicitly solved in [GSS25], see Remark 3.6!

Once we have the abstract isomorphism $\psi: M \simeq \mathcal{O}_0^g$, it remains to make it effective (working coordinate by coordinate, this is the same as making the module action effective, i.e. given $v \in \mathcal{O}_0^g$, being able to evaluate $\varphi_i = \psi^{-1}(v_i) \in M = \text{Hom}(A, E_0)$ on any point of A for $1 \leq i \leq g$). The second crucial ingredient relied on the *heuristic* approach in [Rob24] for extending Clapotis from ideals to modules. Here we give a rigorous version of HD-Clapotis, using as a crucial tool the Poincaré decomposition for modules.

1.5. Technical overview. In Section 2 we recall some facts about abelian varieties, and especially about maximal abelian varieties, which are isomorphic to a product of supersingular elliptic curves E^g . Then we highlight how these varieties connect to the theory of free Hermitian module through an anti-equivalence of categories. Namely, the module version of Deligne's theorem on superspecial abelian varieties is that a torsion free rank g $\mathcal{O}_0$ -module M is isomorphic to $\mathcal{O}_0^g$ ([JKP+18, Theorem 3.3]).

We use this in Section 3, to present the different ways we have to represent principally polarized abelian varieties. More precisely we emphasize that we can represent g -dimensional principally polarized maximal abelian varieties (A, λ_A) as unimodular positive definite Hermitian modules (M, H_M) of rank g over a quaternionic order $\mathcal{O}_0$, where $E_0/\mathbb{F}_{p^2}$ is a maximal supersingular elliptic curve and $\mathcal{O}_0 = \text{End}(E_0)$. Moreover, we show that specificities of such modules allows us to represent the polarized abelian varieties (A, λ_A) as Hermitian modules $(\mathcal{O}_0^g, S)$ where S is the Gram matrix of the Hermitian form induced by H_M over $\mathcal{O}_0^g$. Indeed, in our cases, there exists an isomorphism of modules $\varphi : \mathcal{O}_0^g \rightarrow M$, so that (M, H_M) is isomorphic, as an Hermitian module, to $(\mathcal{O}_0, \varphi^* H_M)$. We also explore the fact that, as in dimension 1 thanks to the Deuring correspondence, we can represent a module (M, H_M) by its endomorphism ring $\text{End}(M)$ enriched by the data of an involution (up to an ambiguity). Knowing that $M \simeq \mathcal{O}_0^g$, we have $\text{End}(M) \simeq \mathcal{M}_g(\mathcal{O}_0)$ and the involution can be described by an Hermitian positive definite matrix over $\mathcal{O}_0$, as in the *IKO* representation of [CDK+25]. We also explain how to navigate between the standard representations $(\mathcal{O}_0^g, S)$ and $(\mathcal{M}_g(\mathcal{O}_0), r_S)$ (resp. abstract representations (M, H_M) and $(\text{End}(M), r_M)$).

In order to compute the standard Hermitian module representation $(\mathcal{O}_0^g, S)$ from an abstract one (M, H_M) we thus need to compute an isomorphism $\mathcal{O}_0^g \rightarrow M$. In Section 3.2 we also show that computing such an isomorphism is in fact equivalent to solving a Principal Ideal Problem over $\mathcal{M}_g(\mathcal{O}_0)$. In Sections 4.1, 4.2 and 4.3 we present an algorithm to solve this problem in polynomial time in $O(\log(p))$. The idea is to work locally and proceed in two steps, assuming that we know a small prime $\ell \in \mathcal{O}_0$ and an element $u \in \mathcal{O}_0$ such that $\text{Nrd}(u) = \ell$. Let $I \subset \mathcal{M}_g(\mathcal{O}_0)$ be an ideal. First, thanks to the KLPT algorithm in dimension 1, we can principalize I locally almost everywhere, except at ℓ . This means that we can compute an $\alpha \in I[\frac{1}{\ell}]$ such that $I[\frac{1}{\ell}] = \alpha \mathcal{M}_g(\mathcal{O}_0)[\frac{1}{\ell}]$. Then we propose an algorithm that allows us to compute a $\delta \in \mathcal{O}[\frac{1}{\ell}]^\times$ so that $\delta \alpha^{-1} I \otimes \mathbb{Z}_\ell = \mathcal{M}_{2g}(\mathbb{Z}_\ell)$, thanks to the element $u \in \mathcal{O}_0$. Then we conclude by noticing that I and $\alpha \delta^{-1} \mathcal{M}_g(\mathcal{O}_0)$ are equal locally everywhere, so they are equal as $\mathbb{Z}$ -lattices. This Section 4 finally leads to our Algorithm 4 that allows us to compute generators of ideals of $\mathcal{M}_g(\mathcal{O}_0)$ in time polynomial in $\log(p)$, in the case where $\mathcal{O}_0$ contains a known quadratic order of small discriminant (see Theorem 4.10). Remark that this set up includes the cryptographic use case where $p \equiv 3 \pmod{4}$ and $\mathcal{O}_0 = \mathbb{Z} + i\mathbb{Z} + \frac{i+j}{2}\mathbb{Z} + \frac{1+k}{2}\mathbb{Z} \subset \mathcal{B}_{p,\infty}$, with the special order $\mathbb{Z}[i] \subset \mathcal{O}_0$. As explained below in Section 2.1, the generic case of a maximal order $\Lambda \subset \mathcal{M}_g(\mathcal{B}_{p,\infty})$ reduces (under GRH) to the one handled above.

Then in Section 5 we show how we can switch between explicit representation (given by a theta null point of level 4 for instance) and abstract representations of polarized abelian varieties as described above. Especially we show in Section 5.1 how the bilinear form $(\varphi_1, \varphi_2) \mapsto \text{Tr}(\lambda_A^{-1} \circ \varphi_1^\vee \circ \lambda_B \circ \varphi_2)$ over $\text{Hom}_k(A, B)$ allows us to efficiently compute the abstract $\mathbb{Z}$ -module structure of $\text{Hom}_k(A, B)$ from explicit representations of its generators, for (A, λ_A) and (B, λ_B) principally polarized abelian varieties. In our general Theorem 5.8, we also prove that we can recover the composition maps, and therefore the ring structure of $\text{End}_k(A)$ in the case where $A = B$. In Section 5.2 we show that the abstract Hermitian module representation already contains the refined information for acting over global differentials. Then in Section 5.3 we show how to recover an explicit representation from an abstract one, using coprime double path.

Section 6 is then dedicated to the construction of double paths between A and E_0^g using the module correspondence. We begin in Section 6.1 by expliciting a technical tool over Hermitian modules, Proposition 6.1. It can be seen as an analog of the Poincaré decomposition, translated in the world of Hermitian modules. In particular, it allows us to compute polarized module isogenies between (M, H_M) and a splitted Hermitian module $M_1 \oplus M_2$, while controlling its polarized degree in Corollary 6.2. This corollary will be the cornerstone of our improvement of KLPT² using module correspondence. We then use the Poincaré decomposition in Section 6.2, to prove an analog of the Clapotis method for higher dimensional modules. Our method relies on the fact that we can compute specific submodules $M_1 \hookrightarrow M$ (resp. $M_2 \hookrightarrow M$) inducing a $2^{n_1} D'_1$ -similitude (resp. $2^{n_2} D'_2$ -similitude) $\Psi_1, \Psi_2: M^4 \rightarrow \mathcal{O}_0^{4g}$, with D'_1 and D'_2 coprime integers, and such that the $2^{\max(n_1, n_2)}$ -torsion on E_0 is accessible (see Proposition 6.6). Then applying the splitting lemma to $\Psi_2 \circ \Psi_1^\vee \in \text{End}(\mathcal{O}_0^{4g})$, and computing the few steps involving 2-isogenies, leads to an efficient representation of the sought double paths. We conclude in Section 6.3 by applying these results to the context of maximal principally polarized abelian varieties (A, λ_A) . Using the previous sections and our construction of double paths, we provide Theorem 6.9 inducing that from the abstract module representation (M, H_M) we can: 1) recover the corresponding principally polarized abelian variety (A, λ_A) along with an effective description of the module action on A , getting rid of the heuristic [Rob24, Heuristic 2.13]; 2) compute an unpolarized isomorphism $A \simeq E_0^g$ (using our solution to the PIP), generalizing [GSS25] in some sense.

Finally, as a last illustration of the flexibility of the module representation, we end this paper by presenting in Section 7 a way to use this theory in order to lower the degree of outputs of the KLPT² algorithm. In Section 7.1, we use the tools of [CDK+25] in order to apply Corollary 6.3, and then compute a polarized module isogeny with a smooth polarized degree reaching a split module $M_1 \oplus M_2$. We finally conclude in Section 7.2 by showing how our method allows for the computation of a polarized isogeny of polarized degree in $O(p^{15.5})$ between polarized abelian surfaces, compared to $O(p^{25})$ given by the KLPT² algorithm. We emphasize that our algorithm inherits their heuristics. The key flexibility that the module representation gives is that we can easily work with submodules $M' \subset M$ given by an orthogonal decomposition $M' = I_1 \oplus^\perp I_2$ of rank 1 submodules. By contrast, the IKO representation $(M_2(\mathcal{O}_0), r)$ of [CDK+25] needs to write down an isomorphism $I_1 \oplus I_2 \simeq \mathcal{O}_0^2$ and can only work at the level of rank 1 submodules implicitly.

1.6. On the module versus the endomorphism ring representation. In [Rob24] it was argued that the module representation is more convenient to use for isogeny based cryptography than the endomorphism ring representation. We expand a bit on this here.

First, we have an actual anti-equivalence of category, see Theorem 2.5, a morphism $\varphi: A_1 \rightarrow A_2$ is represented by an explicit morphism $\varphi^*: M_2 = \text{Hom}(A_2, E_0) \rightarrow \text{Hom}(A_1, E_0)$. By contrast, in the endomorphism ring representation, a morphism $\varphi: A_1 \rightarrow A_2$ is implicitly represented via the image in $\text{End}(A_1)$ of the embedding of the $(\text{End}(A_1), \text{End}(A_2))$ -bimodule $\text{Hom}(A_2, A_1)$ via the inclusion $\varphi^*: \text{Hom}(A_2, A_1) \rightarrow \text{End}(A_1)$. This makes translating the standard categorical constructions (pullback, pushforward, ...) more awkward.

In dimension 1, this awkwardness was less apparent because translating between ideals and orders is easier, so often the arguments and algorithms would implicitly

switch to the module representation on the fly. (Although even in dimension 1, considering invertible ideals only rather than rank 1 projective modules make some constructions more awkward. An invertible ideal I is the same thing as a rank 1 projective module M along with a choice of a specific embedding $i : M \rightarrow R$; which on the geometric side corresponds to a choice of isogeny $i : E_0 \rightarrow E$. In other words this imposes to work in the pointed overcategory, and for the categorical constructions to be compatible with the choice of pointed morphisms. But in practice we would like to work with E without imposing a specific choice of pointed isogeny $i : E_0 \rightarrow E$.)

Secondly, the endomorphism ring representation is not available in the commutative case: already in dimension 1 all (primitively) oriented elliptic curves $E/\mathbb{F}_q$ by a quadratic imaginary order R have the same oriented endomorphism ring $\text{End}_R(E) = R$. By contrast, the R -module representation works just as well in the commutative case, and often allows to treat the maximal case and the oriented case in a uniform way. For instance in Section E, we sketch how to extend HD-Clapotis to the oriented case. Also, even in the maximal supersingular case, the endomorphism ring representation alone introduces an ambiguity, since is not sufficient to distinguish between A and A^σ .

Finally, we remark that the IKO representation, although very compact since it represents (A, λ_A) by a unimodular Hermitian matrix $S \in \mathcal{M}_g(\mathcal{O}_0)$, is quite difficult to work with algorithmically. For HD-Clapotis, we need to work with submodules M' of our initial module M ; which is not easy to do in the IKO representation. (For instance M' may not admit a principal polarization. And even if it did, the IKO representation needs an explicit $\mathcal{O}_0$ -basis of M' , which may be hard to find, and worse does not exist in general for rank 1 submodules). The IKO representation only works for $g > 1$, while the module representation works just as well for $g = 1$ as with $g > 1$. For instance, in our improvement of KLPT², we make crucial use of being able to sample nice rank 1 (but not necessarily free) submodules M' of our initial rank 2 module M .

1.6.1. Acknowledgements. We are grateful to the authors of [CEIV26; Mon26] for sharing with us preliminary versions of their work. We are especially thankful to Montessinos for pointing to us the ambiguity in the endomorphism ring representation due to the outer automorphism of $\mathcal{M}_g(\mathcal{O}_0)$. As mentioned in Section 1.3 we use his solution to clear this ambiguity. We thank Sam Frengley for pointing us misprints. We thank the anonymous reviewers for their insightful comments, which helped clarify and strengthen the manuscript. This work received funding from the France 2030 program managed by the French National Research Agency under grant agreement No. ANR-22-PETQ-0008 PQ-TLS.

2. BACKGROUND

2.1. Notations. In this paper we fix a prime $p > 3$. We denote by $\mathcal{B}_{p,\infty}$ the unique (up to isomorphisms) quaternion algebras over $\mathbb{Q}$, ramified at p and ∞ . We will denote E_0 a fixed supersingular elliptic curve over $\mathbb{F}_{p^2}$, and let $\mathcal{O}_0 = \text{End}(E_0)$. We let $\mathfrak{p}$ be the unique two-sided ideal of $\mathcal{O}_0$, so that $(p) = \mathfrak{p}^2$ and $\mathcal{O}_0/\mathfrak{p} \simeq \mathbb{F}_{p^2}$. If E_0 is defined over $\mathbb{F}_p$, then $\mathfrak{p} = (\pi_{E,p})$ is principal and generated by the Frobenius endomorphism.

In Section 4, we will require the endomorphism ring $\mathcal{O}_0$ associated with an elliptic curve to be extremal in the sense of [KLPT14], i.e. whose endomorphism ring

contains a quadratic order of small discriminant Δ . Such a curve always exist under GRH. In Section 5, to convert between the abstract module and endomorphism ring representations to explicit representations, we will assume that the action of $\mathcal{O}_0$ on E_0 is explicit. This is the case for the special curve E_0 as above. See [HW25] for a detailed discussion about what can be proven without assuming GRH. In our case we will need to assume the GRH anyway for our solution to the PIP.

Typically for isogeny based cryptography, when $p \equiv 3 \pmod{4}$, the base curve is $E_0 : y^2 = x^3 + x$ whose endomorphism ring is isomorphic to the maximal order $\mathcal{O}_0 := \mathbb{Z} + i\mathbb{Z} + \frac{i+j}{2}\mathbb{Z} + \frac{1+ij}{2}\mathbb{Z} \subset \mathcal{B}_{p,\infty}$ where $i^2 = -1$ and $j^2 = -p$, see e.g. [FMP23; Ler25; BDF+24], and in particular contains $\mathbb{Z}[i]$ so is extremal. We emphasize here that solving the PIP for $\mathcal{M}_g(\mathcal{O}_0)$ suffices to solve the general case of any maximal order $\Lambda \subset \mathcal{M}_g(\mathcal{B}_{p,\infty})$. Indeed, as in the proof of Proposition 3.9, at the cost of finding a connecting ideal between $\mathcal{M}_g(\mathcal{O}_0)$ and Λ (which only involves basic linear algebra) we reduce to the extremal case. (This point was also remarked in [CEIV26, Corollary 1.6], to which we refer for more details.) Likewise, if we have the $\mathcal{O}_0$ -module representation $M = \text{Hom}_{\mathbb{F}_{p^2}}(A, E_0)$ of A with respect to E_0 , and we know $I = \text{Hom}_{\mathbb{F}_{p^2}}(E_0, E'_0)$, then $M' = I \otimes_{\mathcal{O}_0} M$ gives the $\mathcal{O}'_0$ -module representation $\text{Hom}_{\mathbb{F}_{p^2}}(A, E'_0)$ of A with respect to E'_0 (see [Rob24, Lemma 4.13]).

For an abelian variety A over a finite field $\mathbb{F}_q$, we denote by $\text{End}(A)$ its endomorphism ring over an algebraic closure $\overline{\mathbb{F}_q}$, and by $\text{End}_{\mathbb{F}_q}(A) \subset \text{End}(A)$ the $\mathbb{F}_q$ -rational endomorphisms.

In this paper, by polynomial time or efficient, we mean polynomial in $\log p$. Some of our arguments work for R -oriented abelian varieties over $\mathbb{F}_q$, in which case we mean polynomial in $\log q$ and Δ_R . (In the maximal supersingular case, $q = p^2$ and $\mathcal{O}_0$ is of discriminant p , so this reduces to $\log p$). We assume that g is fixed throughout. In particular we allow exponential complexity with respect to g . For instance, representing an abelian variety in dimension g via level 2 theta constants requires 2^g coordinates already. When we work with an abstract module or endomorphism ring representations, we assume that our coefficients are also polynomial in $\log p$. Given an arbitrary representation we can always use a lattice reduction algorithm to reduce to this case. When we say that an integer N is (power)smooth, we mean that the largest prime factor ℓ (resp. prime power factor) is polynomial in $\log p$. The N -torsion on E_0 is said to be accessible when one can compute a basis of it in polynomial time in $\log p$; this is the case if N is powersmooth.

2.2. Maximal abelian varieties and Hermitian modules. In this section, we will introduce an equivalence of categories between principally polarized maximal supersingular abelian varieties over $\mathbb{F}_{p^2}$ and suitable unimodular Hermitian modules. Recall that a g -dimensional abelian variety A defined over $\mathbb{F}_{p^2}$ is *maximal*, when $\#A(\mathbb{F}_{p^2}) = (p+1)^{2g}$. Maximal abelian varieties are exactly the varieties reaching the maximal amount of points, prescribed by the Hasse-Weil bound. We will see that a maximal abelian variety over $\mathbb{F}_{p^2}$ is automatically superspecial, hence supersingular, so we will often speak about a “maximal supersingular abelian variety” to emphasize this fact, even though “supersingular” is redundant. The main features of maximal abelian varieties are summarised in the following Theorem 2.2 for which we refer to [EPSV24, Lem. 4] and [JKP+18, Prop. 5.1, Theorem 5.3]:

Example 2.1. If $E/\mathbb{F}_p$ is supersingular, then E is maximal over $\mathbb{F}_{p^2}$. If $E/\mathbb{F}_{p^2}$ is a maximal elliptic curve, then it is supersingular and its endomorphism ring $\text{End}(E)$ is isomorphic to a maximal order in $\mathcal{B}_{p,\infty}$ [Voi21, Thm. 42.1.9].

Theorem 2.2. *Let $A/\mathbb{F}_{p^2}$ be a maximal abelian variety of dimension g . Then A is superspecial, and $\text{End}_{\mathbb{F}_{p^2}}(A) = \text{End}(A)$. Conversely, any polarized superspecial abelian variety $(A_{\overline{\mathbb{F}}_p}, \lambda_{\overline{\mathbb{F}}_p})$ over $\overline{\mathbb{F}}_p$ admit a unique (up to automorphisms) polarized model $(A_{\mathbb{F}_{p^2}}, \lambda_{\mathbb{F}_{p^2}})$ over $\mathbb{F}_{p^2}$ where $A_{\mathbb{F}_{p^2}}$ is maximal.*

If $g > 1$, and $E/\mathbb{F}_{p^2}$ is any maximal supersingular elliptic curve, then A is isomorphic over $\mathbb{F}_{p^2}$ to E^g as an unpolarized abelian variety, so in particular if $\mathcal{O} = \text{End}(E)$, $\text{End}(A) \simeq \mathcal{M}_g(\mathcal{O})$.

Essentially the same result holds for minimal abelian varieties; in fact A is maximal if and only if its quadratic twist is minimal. The choice to work with maximal supersingular abelian varieties over $\mathbb{F}_{p^2}$ rather than superspecial ones is made in order to avoid additional complications arising from extra twists. (In dimension 1, only the curves with $j = 0, 1728$ can have supersingular twists that are neither minimal nor maximal over $\mathbb{F}_{p^2}$, but there are many more abelian varieties with extra twists in higher dimension). In particular, the maximal supersingular isogeny graph in $\mathbb{F}_{p^2}$ is isomorphic to the superspecial isogeny graph in $\overline{\mathbb{F}}_p$. Recall that $\mathcal{O}_0 = \text{End}(E_0)$ is a maximal quaternionic order in $\mathcal{B}_{p,\infty}$.

Definition 2.3. A unimodular positive definite Hermitian $\mathcal{O}_0$ -module (M, H_M) is a torsion-free left $\mathcal{O}_0$ -module finitely generated together with a Hermitian bilinear form $H_M : M \times M \rightarrow \mathcal{O}_0$ which is positive definite and unimodular (which means that H_M induces an isomorphism of M with its dual $M^\vee$).

The rank of M is the dimension of $M \otimes_{\mathbb{Z}} \mathbb{Q}$ over $\mathbb{Q}$. We denote by $q_M : M \rightarrow \mathcal{O}_0$ the quadratic form associated with H_M : if $x \in M$, $q_M(x) = H_M(x, x)$.

We will often simply say that (M, H_M) is unimodular as an abbreviation of unimodular positive definite torsion free and finitely presented. We will also abbreviate *finitely presented torsion free* by f.p.t.f. . We refer to [Rob24, § 2] for some basis theory of unimodular Hermitian modules. In the quaternionic case, forgetting about the Hermitian form, the module structure is very simple:

Theorem 2.4. [JKP+18, Theorem 3.3 (c)] *Let M be a torsion-free left $\mathcal{O}_0$ -module of rank g . Then M is projective, and if $g > 1$ M is free: $M \simeq \mathcal{O}_0^g$.*

2.3. The equivalence of categories. The Deuring correspondence has been a major tool used in the recent advances regarding isogeny-based cryptography. The following can be seen as an extension of this correspondance to higher dimension:

Theorem 2.5. [JKP+18, Theorem 5.3 (b)] and [Rob24, Theorem 5.1].

The functor $A \mapsto M = \text{Hom}_{\mathbb{F}_{p^2}}(A, E_0)$ is an anti-equivalence of categories between maximal abelian varieties over $\mathbb{F}_{p^2}$ and f.p torsion-free left $\mathcal{O}_0$ -modules. Under this equivalence, a principal polarization on A corresponds to an unimodular positive definite Hermitian form H over M .

Example 2.6. Combined with Theorem 2.4, Theorem 2.5 shows that every maximal supersingular abelian variety $A/\mathbb{F}_{p^2}$ is (non-polarized) isomorphic to E_0^g .

The converse functor can be defined as follows: to $M/\mathcal{O}_0$ we attach the abelian variety $A/\mathbb{F}_{p^2}$ whose associated fppf sheaf is given by the internal sheaf hom

Maximal abelian variety $A/\mathbb{F}_{p^2}$ of dimension g	Left f.p.t.f. $\mathcal{O}_0$ -module $M = \text{Hom}_{\mathbb{F}_{p^2}}(A, E_0)$ of rank g
The dual abelian variety $\hat{A}$	The dual module $M^\vee$ (or $M^\sharp$ if unimodular)
Principal polarization $\lambda_A: A \rightarrow \hat{A}$	U.p.d Hermitian form $H_M: M \times M \rightarrow \mathcal{O}_0$
An isogeny $\varphi: A_1 \rightarrow A_2$ of degree n	A module monomorphism $\psi: M_2 \hookrightarrow M_1$ such that $\text{Card}(M_1/\psi(M_2)) = n$
An n -isogeny $\varphi: (A_1, \lambda_1) \rightarrow (A_2, \lambda_2)$	An n -similitude $\psi: (M_2, H_2) \rightarrow (M_1, H_1)$

TABLE 1. Summary of the module correspondence.

$\mathcal{HOM}_{\mathcal{O}_0\text{-fppf}}(M, E_0)$. (It is not hard to prove that A is a proper commutative group scheme, but to prove that it is an abelian variety requires more work.) Of course this abstract definition does not allow to compute A in practice, one of the goals of this paper is to give such an effective algorithm.

Recall that $\varphi: A \rightarrow B$ is an isogeny if it is a surjective morphism between abelian varieties of the same dimension. On the module side this corresponds to an injective morphism $\psi: M_B \rightarrow M_A$ between torsion free modules of the same rank. The degree of φ is the order of the finite scheme $\text{Ker } \varphi$, which is linked to the (finite) cardinal of $M_A/\psi(M_B)$.

In practice we deal with principally polarized abelian varieties, and isogenies that respect the polarizations: $\varphi: (A, \lambda_A) \rightarrow (B, \lambda_B)$ are said to be polarized of type $[n]$ if $\tilde{\varphi} \circ \varphi = [n]$, where $\tilde{\varphi} = \lambda_A^{-1} \circ \varphi^\vee \circ \lambda_B$ is the polarized dual of φ . By abuse of notation we will also say that φ is polarized of type (or degree) n , note that in this case $\deg \varphi = n^g$. On the module side this corresponds to a n -similitude between unimodular Hermitian modules $\psi: (M_B, H_B) \rightarrow (M_A, H_A)$ satisfying $\psi^* H_A = n H_B$.

Example 2.7.

- i) The module associated with the abelian variety (E_0^g, λ_0^g) is $(\mathcal{O}_0^g, H_{\mathcal{O}_0^g})$, where $H_{\mathcal{O}_0}(x, y) = x\bar{y}$ (which we will also denote by H_0).
- ii) In the other direction, we recover the Deuring correspondence for ideals [JKP+18, Theorem 4.4 (d)]. Indeed, the variety associated with the ideal I seen as a $\mathcal{O}_0$ -module is $E_0/E_0[I]$.
- iii) Any unimodular principal Hermitian module of the form $(\mathcal{O}_0 x, \frac{H_M}{q_M(x)})$ is isomorphic as a unimodular module to $(\mathcal{O}_0, H_0)$, hence corresponds to the polarized isomorphism class of (E_0, λ_0) .

In Table 1 we summarize the main translations provided by Theorem 2.5.

Example 2.8. Let $\varphi: A_1 \rightarrow A_2$ be the n -isogeny associated with the n -similitude $\psi: (M_2, H_2) \rightarrow (M_1, H_1)$. By Proposition 3.5, $M_2 \subset M_1$ can also be equivalently described as an ideal $I \subset \text{End}_{\mathcal{O}_0}(M_1) \simeq \text{End}(A_1)$. Then $\text{Ker } \varphi = \{P \in A_1 \mid \gamma(P) = 0 \forall \gamma \in M_2\} = \{P \in A_1 \mid \gamma(P) = 0 \forall \gamma \in I\}$, and conversely $M_2 = \{\gamma \in M_1 \mid \gamma(\text{Ker } \varphi) = 0\}$, $I = \{\gamma \in \text{End}(A_1) \mid \gamma(\text{Ker } \varphi) = 0\}$. This allows to go from φ to ψ (resp. I) whenever n is smooth, the n -torsion is accessible on A_1 , and the module action is explicit (resp. the endomorphism action is explicit). In particular, if $\psi: M \rightarrow \mathcal{O}_0^g$ is an n -similitude for a powersmooth n , then we can compute the corresponding isogeny $E_0^g \rightarrow A$ and hence recover the abelian variety A , by [Rob24, Corollary 4.27] (this reference only treats the module point of view,

but the statements on I above follow immediately from the statements on M_2 and Proposition 3.5).

3. REPRESENTING POLARIZED ABELIAN VARIETIES

Let (A, λ_A) be a (polarized isomorphism class of) dimension g principally polarized maximal abelian variety over $\mathbb{F}_{p^2}$. We will denote by r_A the Rosati involution on $\text{End}(A)$ induced by λ_A (up to isomorphism).

3.1. The abstract representations.

The abstract Hermitian module representation (M, H_M) . This is the representation given by the equivalence between principally polarized abelian varieties and unimodular Hermitian modules. We assume that M is given by generators, viewed as a submodule of $\mathcal{O}_0^g$, and H_M is given by an Hermitian Gram matrix. We highlight that H_M is well defined up to congruence, meaning that for any $P \in \text{GL}_g(\mathcal{O}_0)$, (M, H_M) and $(M, PH_M \bar{P}^t)$ represent the same Hermitian module.

Remark 3.1. Note that if M is given by an abstract presentation:

$$\mathcal{O}_0^m \xrightarrow{\gamma} \mathcal{O}_0^q \longrightarrow M \longrightarrow 0, \text{ for integers } m, q \in \mathbb{Z}_{\geq 1},$$

then we can reduce to the case of generators over $\mathcal{O}_0^g$ by linear algebra. Indeed, let us denote by $G \in \mathcal{M}_{q \times m}(\mathcal{O}_0)$ the matrix given by the linear application γ in canonical basis. The columns of G are the m relations between the q generators of M given by the presentation. Now, extending the scalar, we obtain a matrix $G_{\mathbb{Q}} = G \otimes \mathbb{Q}$ representing a linear application between *vector spaces*. Therefore it is easy to compute its cokernel, given by a matrix $C \in \mathcal{M}_{g \times q}(\mathcal{B}_{p, \infty})$ (g rows since M is of rank g). Then, rescaling by the greatest common divisor of the denominators of the entries of C , we recover a matrix $C' \in \mathcal{M}_{g \times q}(\mathcal{O}_0)$. Its columns give the q generators of M seen as an $\mathcal{O}_0^g$ submodule.

The standard Hermitian module representation $(\mathcal{O}_0^g, S)$. Let (M, H_M) be the abstract module representation of (A, λ_A) . By Theorem 2.4, if $g > 1$, the $\mathcal{O}_0$ -module M is isomorphic to $\mathcal{O}_0^g$. Fixing an isomorphism $\psi: \mathcal{O}_0^g \rightarrow M$, we obtain an Hermitian form $H = \psi^* H_M$ on $\mathcal{O}_0^g$, which is represented by a matrix S in the canonical basis: $S = H_M(e_i, e_j)$, so that $H_M(u, v) = uSv^*$.

The abstract endomorphism representation $(\text{End}(M), r_M)$. Given the abstract module representation (M, H_M) , we can also construct the endomorphism representation $(\text{End}(M), r_M)$ where $\text{End}(M)$ is the endomorphism ring of M as a left $\mathcal{O}_0$ -module (so M is a $(\mathcal{O}_0, \text{End}(M))$ -bimodule), and r_M is the “Rosati involution” $\gamma \mapsto \gamma^*$, where γ^* is the adjoint of γ with respect to H_M .

In the abstract endomorphism representation, we work in the category of rings with an involution. An isomorphism $(\mathcal{A}_1, r_1)$ of $(\mathcal{A}_2, r_2)$ of rings with an involution is an isomorphism $\varphi: \mathcal{A}_1 \rightarrow \mathcal{A}_2$ such that $r_2 = \varphi r_1 \varphi^{-1}$. By the anti-equivalence of category, $(\text{End}(M), r_M) \simeq (\text{End}(A), r_A)$ as abstract rings with an involution. This is the usual representation in dimension 1: a maximal supersingular elliptic curve $E/\mathbb{F}_{p^2}$ is represented by $\mathcal{O} = \text{End}(E)$, and where the Rosati involution is implicitly given by the quaternionic conjugation on $\mathcal{O}$.

Remark 3.2 (The refined endomorphism ring representation). A very important point, is that the abstract endomorphism ring representation does not determine the Hermitian module M uniquely. This is because $\mathcal{M}_g(\mathcal{O}_0)$ has exactly one outer

automorphism as a ring, $(\text{End}(M), r_M)$ can be induced by two non isomorphic Hermitian modules. Explicitly, (M, H_M) and $(\mathfrak{p}M, H_M/p) \simeq (M, \mathfrak{p}H_M\mathfrak{p}^{-1})$ have isomorphic associated endomorphism ring with involution (on the abelian variety sides, these correspond to A and its Galois conjugate A^σ). This phenomena is well known in dimension 1: a supersingular curve E is uniquely determined by the ideal $I = \text{Hom}(E, E_0)$, but the endomorphism ring $\text{End}(E)$ only determines E up to Galois conjugation.

To distinguish the two cases, we introduce the refined ring representation. The idea is that the left $\mathcal{O}_0$ -module structure on M induces a $\mathbb{F}_{p^2}$ -module structure on $M/\mathfrak{p}M \simeq \mathbb{F}_{p^2}^g$, hence an isomorphism $\text{End}(M)/\mathfrak{p}' \simeq \mathcal{M}_g(\mathbb{F}_{p^2})$ (where $\mathfrak{p}' \subset \text{End}(M)$ is the pullback of the Jacobson radical of $\text{End}(M)/p$). In particular, M induces an $\mathbb{F}_{p^2}$ -algebra structure on $\text{End}(M)/\mathfrak{p}'$, independent of the choice of basis of $M/\mathfrak{p}M$. This extra $\mathbb{F}_{p^2}$ -algebra structure gives us our refined representation.

Notice that the $\mathbb{F}_{p^2}$ -algebra structures on $(\text{End}(M), r_M)$ associated with the two modules above differ precisely by the Galois conjugation of $\mathbb{F}_{p^2}/\mathbb{F}_p$.

The standard endomorphism representation $(\mathcal{M}_g(\mathcal{O}_0), r_S)$. Given the standard Hermitian module representation $(\mathcal{O}_0^g, S)$, we obtain the standard endomorphism representation $(\mathcal{M}_g(\mathcal{O}_0), r_S)$ where $r_S(M) = SM^*S^{-1}$ and M^* is the canonical involution $M \mapsto \overline{M}^t$. (Here the $\mathcal{O}_0/\mathfrak{p}$ -module structure on $\mathcal{M}_g(\mathcal{O}_0)/\mathfrak{p}' \simeq \mathcal{M}_g(\mathcal{O}_0/\mathfrak{p})$ should of course be the natural one.)

Recall that an involution r on $\mathcal{M}_g(\mathcal{B}_{p,\infty}) = \mathcal{M}_g(\mathcal{O}_0) \otimes_{\mathbb{Z}} \mathbb{Q}$ is positive if $\text{Tr}_{\mathcal{B}_{p,\infty}/\mathbb{Q}}(Mr(M)) > 0$ for all $M \in \mathcal{M}_g(\mathcal{B}_{p,\infty})$. If r is an involution on $\mathcal{M}_g(\mathcal{O}_0)$, it induces an involution on $\mathcal{M}_g(\mathcal{B}_{p,\infty})$ that we denote by r too. We say that r is an $\mathbb{F}_{p^2}$ -involution if $r \bmod \mathfrak{p}'$ respects the $\mathbb{F}_{p^2}$ -algebra structure of $\mathcal{M}_g(\mathcal{O}_0)/\mathfrak{p}'$. (Otherwise, it acts by the $\mathbb{F}_{p^2}/\mathbb{F}_p$ -Galois action on scalar matrices, and conjugating r by the outer automorphism we get an $\mathbb{F}_{p^2}$ -involution.)

Lemma 3.3. *Let r be a positive $\mathbb{F}_{p^2}$ -involution on $\mathcal{M}_g(\mathcal{O}_0)$. Then $r = r_S$ for some unimodular Hermitian positive definite matrix (and conversely).*

Proof. Let $r_0(M) = M^*$ be the canonical involution on $\mathcal{M}_g(\mathcal{O}_0)$, then $a = r \circ r_0$ is an automorphism. If r is an $\mathbb{F}_{p^2}$ -involution, since r_0 is an $\mathbb{F}_{p^2}$ -involution too, then a is an inner automorphism, so $r = SM^*S^{-1}$ for some matrix $S \in \mathcal{M}_g(\mathcal{O}_0)^\times$. Extending the scalars by $\mathbb{R}$, by [Voi21, Lemma 8.4.12], we see that up to replacing S by $-S$, S is Hermitian positive definite. Since its determinant is a rational number of absolute value 1, by positivity of S it is 1. The converse is immediate. $\square$

Remark 3.4. It is immediate to convert from $(\mathcal{O}_0^g, S)$ to $(\mathcal{M}_g(\mathcal{O}_0), r_S)$. In the other direction, we can recover S from r_S by solving in $X \in \text{GL}_g(\mathcal{O}_0)$ the linear system given by $r(M_i)X = X\overline{M_i}^t$, for M_i the g^2 canonical generators of $\mathcal{M}_g(\mathcal{O}_0)$. Notice that a solution to this system will be well defined up to multiplication by a scalar since $\mathcal{M}_g(\mathcal{O}_0)$ is central. But the condition of both positivity and $\det(S) = 1$ lead to a unique S .

We remark that S , the Gram matrix of the associated Hermitian form H on $\mathcal{O}_0^g$: it is precisely what is called the IKO representation in [CDK+25] when $g = 2$. Changing basis of $\mathcal{O}_0^g$ amounts to changing S to GSG^* for G an element in $\text{GL}_g(\mathcal{O}_0)$, and to conjugating $r_S : M \rightarrow SM^*S^{-1}$ by C_G : $r_{GSG^*} = C_G \circ r_S \circ C_G^{-1}$, where $C_G(M) = GMG^{-1}$.

3.2. Converting between the abstract representations.

Proposition 3.5. *The maps $M' \subset M \mapsto I_{M'} = \{\gamma \in \text{End}(M) \mid \gamma(M) \subset M'\}$ and $I \subset \text{End}(M) \mapsto M'_I = \sum_{\gamma \in I} \gamma(M) \subset M$ give converse bijections between submodules of M and ideals of $\text{End}(M)$.*

For a general module M finitely generated over a ring R , we have obvious inclusions $M'_{I_{M'}} \subset M'$ and $I_{M'_I} \supset I$. If M is projective, it is locally free over R hence these inclusions are equalities. Proposition 3.5 is a baby case of the more general Morita equivalence between $\mathcal{O}_0$ and $\mathcal{M}_g(\mathcal{O}_0)$, see Section C for more details. As we will see in Section 4, Proposition 3.5 implies that finding an explicit isomorphism $M \simeq \mathcal{O}_0^g$ given an abstract representation of M as $M \subset \mathcal{O}_0^g$ and solving the Principal Ideal Problem over $\mathcal{M}_g(\mathcal{O}_0)$ are equivalent problems.

Remark 3.6. Alternatively, one can directly solve the submodule problem as follows (hence obtain yet another solution to the PIP). First, by standard linear algebra we can write M as a direct sum of ideals: $M \simeq I_1 \oplus \cdots \oplus I_g$. This is standard in the commutative case by using pseudo-basis, and the same algorithm can be adapted to the non-commutative case, see for instance [Kir16, Algorithm 2.2.6]. It remains to find an isomorphism $I_1 \oplus \cdots \oplus I_g \simeq \mathcal{O}_0^g$, which reduces to the case $g = 2$ and is solved in [GSS25] (under GRH).

It is immediate to convert from (M, H_M) to the refined representation $(\text{End}(M), r_M)$. The other direction is less obvious, we begin by computing an isomorphism $\text{End}(M) \simeq \mathcal{M}_g(\mathcal{O}_0)$.

The first step is to compute an isomorphism $\text{End}(M) \otimes \mathbb{Q} \rightarrow \mathcal{M}_g(\mathcal{B}_{p,\infty})$. Unfortunately, there is no polynomial time algorithm (even for fixed g) in the literature to do this: the most general algorithm is [IRS12] but it requires a factorization oracle, so it only gives a polynomial time quantum algorithm. It is enough to show how to find isomorphisms $\mathcal{A} \cong \mathcal{M}_n(\mathbb{Q})$, given a central simple Q -algebra $\mathcal{A}$ as input. The algorithm proceeds by computing a maximal order Λ (using the factorization oracle) and finding a special element in a lattice constructed from Λ among its vectors in a certain ball. If the lattice does not contain very short vectors, this can be done efficiently when n is fixed, but this condition is not always satisfied. When the lattice contains very short vectors, the algorithm from [IRS12] uses a recursive step: this requires computing more maximal orders and therefore make more calls to the factorization oracle. In our setting, we can compute a maximal order in polynomial time in the initial algebra since we know a good order to start from, but not in the recursive steps since badly controlled algebras could appear. We show that we can avoid the recursion by randomizing the maximal order, using the techniques of [BPTW25] and the fact that random lattices do not have very short vectors with high probability.

Proposition 3.7. *Given as input a maximal order Λ in an algebra $\mathcal{A}$ isomorphic to $\mathcal{M}_n(\mathbb{Q})$, Algorithm 1 returns an isomorphism $\mathcal{A} \cong \mathcal{M}_n(\mathbb{Q})$ in probabilistic polynomial time for fixed n .*

Proof. The algorithm is adapted from [IRS12], so we refer to there for details of intermediate steps. We also use the terminology of Euclidean lattices, cf. [Gra84; BPTW25].

Let $c_1(n) > 0$ be a lower bound on the volume of semistable lattices of rank n . An explicit such lower bound can be derived from [SW14] or [BPTW25, Section 5].

Algorithm 1: SPLIT

Input: A maximal order Λ in an algebra $\mathcal{A}$ isomorphic to $\mathcal{M}_n(\mathbb{Q})$

Output: An isomorphism $\mathcal{A} \cong \mathcal{M}_n(\mathbb{Q})$

// For the definition of the c_i , see the proof
of Proposition 3.7

- 1 Compute an isomorphism $\iota: \mathcal{A} \otimes \mathbb{R} \cong \mathcal{M}_n(\mathbb{R})$;
 - 2 $\Lambda' \leftarrow \Lambda$, $\iota' \leftarrow \iota$, $\lambda \leftarrow \lambda_1(\Lambda, \iota)$;
 - 3 Let ℓ be the next prime after $c_3(n, \lambda)$;
 - 4 Compute an isomorphism $\Lambda/\ell\Lambda \cong \mathcal{M}_n(\mathbb{F}_\ell)$;
 - 5 **while** $\lambda_1(\Lambda', \iota') < \exp(-c_2(n))$ **do**
 - 6 Let ι' be a small random perturbation of ι ;
 - 7 Let $H \subset \mathbb{F}_\ell^n$ be a uniformly random hyperplane ;
 - 8 Let $H' \subset (\mathbb{Z}/\ell^2\mathbb{Z})^n$ be the preimage of H ;
 - 9 Let $\Lambda' = \frac{1}{\ell}\{b \in \Lambda \mid b(H') \subset \ell H'\}$
 - 10 **end**
 - 11 Compute the set X of elements $a \in \Lambda'$ with $\|\iota'(a)\| \leq n$;
 - 12 Let $a \in X$ be such that $\dim_{\mathbb{Q}}(\mathcal{A}a) = n$;
 - 13 Return the isomorphism $\mathcal{A} \cong \mathcal{M}_n(\mathbb{Q})$ derived from the action of $\mathcal{A}$ on $\mathcal{A}a$.
-

Let $c_2(n)$ be the constant c appearing in [Gra86, Theorem 2.2] when $L = M$ is a lattice of rank n . By (Λ', ι') we denote the Euclidean lattice obtained by pulling back via ι' the standard Euclidean structure on $\mathcal{M}_n(\mathbb{R})$. Note that since n is fixed, the first minimum of any lattice of rank n^2 can be computed in polynomial time, see for instance [HS07].

The order Λ is isomorphic to $\text{End}(L)$ for some Euclidean lattice L in a way that is compatible with ι . From a lower bound on $\lambda_1(\Lambda, \iota)$, by [Gra86, Theorem 2.2], we can derive a lower bound on $\lambda_1(L)$ (first of the successive minima). From [BPTW25, Theorem 3], determine a small random perturbation over $\mathbb{R}$ and a bound $c_3(n, \lambda) = \lambda^{-O_n(1)}$ such that random sublattices of L of prime index at least $c_3(n, \lambda)$ are $c_1(n)/2$ -close to equidistributed in the space of lattices of rank n . Such a bound can be derived without knowledge of L but simply from a lower bound on its first minimum (in the terminology of [BPTW25] this gives a bound on the balancedness parameter α). Note that in our case, since the base field is $\mathbb{Q}$, the Riemann hypothesis and averaging over several primes are unnecessary.

Since the isomorphism $\mathcal{A} \otimes \mathbb{R} \cong \mathcal{M}_n(\mathbb{R})$ can be computed in polynomial time, its coefficients expressed on a basis of Λ have polynomial size, and therefore $\log \lambda_1(\Lambda, \iota)$ is polynomially bounded, and similarly for $\log \ell$ when n is fixed.

In the while loop of Algorithm 1, the order Λ' that is constructed (note $\ell\Lambda \subset \Lambda' \subset \frac{1}{\ell}\Lambda$) is of the form $\text{End}(L')$ where L' is the preimage of H in L via an isomorphism $L/\ell L \cong \mathbb{F}_\ell^n$ compatible with $\text{End}(L)/\ell \text{End}(L) = \Lambda/\ell\Lambda \cong \mathcal{M}_n(\mathbb{F}_\ell)$; by construction L' is a uniformly random sublattice of index ℓ in L . Therefore, with probability at least $c_1(n)/2$, the lattice L' is semistable, and in this case by [Gra86, Theorem 2.2], the loop terminates. In particular, the expected number of loop iterations is at most $2/c_1(n)$.

Since n is fixed and $\lambda_1(\Lambda', \iota')$ is lower bounded by a function of n , the set X can be computed in polynomial time [FP85; HS07]. By [IRS12, Theorem 2], the

element $a \in X$ always exists. The last step can be computed in polynomial time as in [IRS12]. $\square$

Corollary 3.8. *Let $\mathcal{A}_1, \mathcal{A}_2$ be two isomorphic central simple algebras over $\mathbb{Q}$ of dimension g^2 . Given orders $\Lambda_1 \subset \mathcal{A}_1$ and $\Lambda_2 \subset \mathcal{A}_2$ together with factorizations of their discriminants, one can compute an isomorphism $\mathcal{A}_1 \cong \mathcal{A}_2$ in probabilistic polynomial time for fixed g .*

Proof. Since we know the factorization of the discriminants of Λ_1 and Λ_2 , by [IR93] we can efficiently compute a maximal order $\Lambda \subset \mathcal{A}$ containing $\Lambda_1 \otimes \Lambda_2^{\text{op}}$, where $\mathcal{A} = \mathcal{A}_1 \otimes \mathcal{A}_2^{\text{op}} \simeq \mathcal{M}_{g^2}(\mathbb{Q})$. Then we apply Proposition 3.7 (to $\Lambda \subset \mathcal{A}$ for $n = g^2$) and [IRS12, Section 4]. $\square$

Applying the previous corollary for the orders and $\mathbb{Q}$ -algebras $\text{End}(M) \subset \text{End}(M) \otimes \mathbb{Q}$ and $\mathcal{M}_g(\mathcal{O}_0) \subset \mathcal{M}_g(\mathcal{B}_{p,\infty})$ we can efficiently compute an isomorphism $\text{End}(M) \otimes \mathbb{Q} \rightarrow \mathcal{M}_g(\mathcal{B}_{p,\infty})$. Indeed the discriminants involved are powers of p and are therefore easy to factor. However to conclude our proof, we rely on a solution of the Principal Ideal Problem in $\mathcal{M}_g(\mathcal{O}_0)$, not yet presented. Hence, for now, we only show how computing such an isomorphism reduces to the PIP.

Proposition 3.9. *With the above notations, for $g \geq 2$. Computing an isomorphism $\text{End}(M) \simeq \mathcal{M}_g(\mathcal{O}_0)$, reduces to solving the PIP in $\mathcal{M}_g(\mathcal{O}_0)$.*

Proof. Since we can compute an isomorphism $\text{End}(M) \otimes \mathbb{Q} \simeq \mathcal{M}_g(\mathcal{B}_{p,\infty})$, we deduce an inclusion $i: \text{End}(M) \hookrightarrow \mathcal{M}_g(\mathcal{B}_{p,\infty})$. Let us denote by $\mathcal{O}_1 = \mathcal{M}_g(\mathcal{O}_0)$ and $\mathcal{O}_2 = i(\text{End}(M))$ the involved orders inside $\mathcal{M}_g(\mathcal{B}_{p,\infty})$.

Now $\mathcal{O}_2$ must be isomorphic to $\mathcal{O}_1$. Since the center $\mathbb{Q}$ has no nontrivial automorphisms, the Skolem–Noether theorem [Voi21, Theorem 7.7.1] implies that there exists a matrix $P \in \mathcal{O}_1$ such that the map $\text{End}(M) \rightarrow \mathcal{O}_1$ given by $\alpha \mapsto Pi(\alpha)P^{-1}$ is a ring isomorphism, and in particular $\mathcal{O}_2 = P\mathcal{O}_1P^{-1}$. Finding this P reduces to a PIP instance in $\mathcal{O}_1$ for an ideal connecting $\mathcal{O}_1$ and $\mathcal{O}_2$.

Indeed, let I_c be any ideal connecting $\mathcal{O}_1$ and $\mathcal{O}_2$ (which are both maximal orders). Such an I_c can, for instance, be efficiently computed as in [CEIV26, Above Corollary 1.6], $I_c = e\mathcal{O}_1\mathcal{O}_2$, for $e \in \mathbb{Z}_{\geq 1}$ the exponent of the group quotient $\mathcal{O}_2/(\mathcal{O}_1 \cap \mathcal{O}_2)$. Since the narrow class group of $\mathbb{Q}$ is trivial, by Eichler’s theorem [Voi21, Theorem 28.5.5, Corollary 28.5.17] the ideal I_c is principal. Furthermore if $I_c = \mathcal{O}_1\beta$, then $\mathcal{O}_2 = \beta^{-1}\mathcal{O}_1\beta$. That is why solving a principal generator for a connecting ideal between $\mathcal{O}_1$ and $\mathcal{O}_2$ gives such a P . $\square$

The results of the present section, especially Proposition 3.9, will enable us to switch between different representation once a solution to the PIP has been obtained. We defer a summary of the resulting translations to Section 4.4, and now focus on the Principal Ideal Problem.

4. ON THE PRINCIPAL IDEAL PROBLEM

This section is dedicated to the resolution of the Principal Ideal Problem for matrix rings over the maximal quaternionic order $\mathcal{O}_0$. In the following, $g \geq 2$.

Problem 4.1. *Let I be an integral right-ideal in $\mathcal{M}_g(\mathcal{O}_0)$, given by generators. Compute an element $P \in I$ such that $I = P\mathcal{M}_g(\mathcal{O}_0)$.*

The Principal Ideal Problem has a long history in the context of number fields (see [Coh93, Section 6.5.5]). In quaternion algebras it has been studied before [DD08; KV10; Pag14; KLPT14] and is relevant to lattice-based cryptography [CME+25]. In more general semisimple algebras it has been studied for its applications to integral representation theory [BHJ22] and very recently in the same context as the present paper [CEIV26]. As stated earlier this problem is equivalent to the submodule problem below, by Morita equivalence.

Problem 4.2. *Given generators $m_1, \dots, m_r \in \mathcal{O}_0^g$ of a free right $\mathcal{O}_0$ -module M of rank $g \geq 2$, compute a basis of M as a submodule of $\mathcal{O}_0^g$.*

In the following we choose to fit in the framework of [Rei03], so we will work with right $\mathcal{O}_0$ -modules (and right $\mathcal{M}_g(\mathcal{O}_0)$ -ideals) instead of left $\mathcal{O}_0$ -module as above. Of course, by applying transpose-conjugate, the left setting is equivalent to the right setting.

To facilitate the readability of Proposition 4.8 and Theorem 4.10 we use the vague adjective *small* for a prime ℓ and a discriminant Δ of a quadratic order. Let us make this precise. The complexity is described in terms of the parameter p , the ramified prime of the quaternion algebra we are dealing with. For both ℓ and Δ , we say that they are small to mean that they are in $O(\log(p))$.

In the following sections we present an algorithm to solve Problem 4.1 using the KLPT algorithm as a subroutine and linear algebra over $\mathcal{O}_0$ and its localizations. We divide the solution of Problem 4.1 in two steps. First we reduce to the case of a principal ideal I of $\mathcal{M}_g(\mathcal{O}_0)[\frac{1}{\ell_1 \dots \ell_r}]$, and then we try to multiply I by elements of $\mathcal{M}_g(\mathcal{O}_0)[\frac{1}{\ell_1 \dots \ell_r}]^\times$ so that it becomes locally trivial at every prime ℓ_i . This is the same strategy as in [Pag14] and was inspired by it, but the features of our special case allow us to design a polynomial time algorithm. For instance, our local algorithm in Section 4.2 has a natural geometric interpretation on the Bruhat–Tits building of $GL_{2g}(\mathbb{Q}_\ell)$ similar to [Pag14, Subalgorithm 3.9 and Figure 2].

4.1. Solving the problem almost everywhere. In the following we will denote by $\mathcal{O}$ the algebra $\mathcal{M}_g(\mathcal{O}_0)$, with $g \geq 2$. Let I be a right $\mathcal{O}$ -ideal, and $S = \{\ell_1, \dots, \ell_r\}$ a set of small primes such that $\ell_i \neq p$. In the following we denote by π the product $\ell_1 \dots \ell_r$. The first step is to find a generator of $I[\frac{1}{\pi}]$. We can handle dimension 1 thanks to the KLPT algorithm.

Lemma 4.3. *Let I be an integral right $\mathcal{O}_0$ -ideal. There exists $\alpha \in I[\frac{1}{\pi}]$ such that $I[\frac{1}{\pi}] = \alpha \mathcal{O}_0[\frac{1}{\pi}]$.*

Proof. KLPT returns an $\alpha \in \mathcal{B}_{p,\infty}^\times$ such that $I = \alpha J$, where J is an ideal of norm $\ell_1^{n_1} \dots \ell_r^{n_r}$ for positive integers n_i . Thus we can write J as a sum $J = j_1 \mathcal{O}_0 + \ell_1^{n_1} \dots \ell_r^{n_r} \mathcal{O}_0$ which leads to $J[\frac{1}{\pi}] = j_1 \mathcal{O}_0[\frac{1}{\pi}] + \ell_1^{n_1} \dots \ell_r^{n_r} \mathcal{O}_0[\frac{1}{\pi}] = \mathcal{O}_0[\frac{1}{\pi}]$. Thus $I[\frac{1}{\pi}] = \alpha J[\frac{1}{\pi}] = \alpha \mathcal{O}_0[\frac{1}{\pi}]$. Since $1 \in \mathcal{O}_0[\frac{1}{\pi}]$, it is clear that $\alpha \in I[\frac{1}{\pi}]$. $\square$

We now generalize this lemma to the right ideal I of $\mathcal{O}$. The fact that working with I a $\mathcal{O}$ -right ideal, is equivalent to working with the $\mathcal{O}_0$ -module generated by the columns of the given generators of I , is crucial for Algorithm 2.

Proposition 4.4. *Given an integral right $\mathcal{O}$ -ideal I and a set S of small primes distinct from p , Algorithm 2 returns an element $\alpha \in I[\frac{1}{\pi}]$ such that $I[\frac{1}{\pi}] = \alpha \mathcal{O}[\frac{1}{\pi}]$. Assuming GRH, this algorithm runs in polynomial time in $\log(p)$.*

Proof. Let us prove the correctness of Algorithm 2. Let us denote by $v_1, \dots, v_{ng}$ the list of the column vectors of the $n \geq 2$ generators of I , and M_π the $\mathcal{O}_0[\frac{1}{\pi}]$ module generated by the v_i . We want to compute g vectors $w_1, \dots, w_g \in \mathcal{O}_0^g[\frac{1}{\pi}]$ such that $M_\pi = \langle w_1, \dots, w_g \rangle_{\mathcal{O}_0[\frac{1}{\pi}]}$. At the cost of repeating the process below, we may assume that there is only $g+1$ vectors v_i .

By Lemma 4.3, we know that there exists $\alpha_1 \in \mathcal{O}_0[\frac{1}{\pi}]$ such that $\alpha_1 \mathcal{O}_0[\frac{1}{\pi}] = \langle v_{1,1}, \dots, v_{g+1,1} \rangle_{\mathcal{O}_0[\frac{1}{\pi}]}$, where $v_{i,j}$ denotes the j -th component of v_i . Thus there exists $c_1, \dots, c_g, d_1, \dots, d_{g-1} \in \mathcal{O}_0[\frac{1}{\pi}]$ such that:

$$w_1 := \sum_{i=1}^{g+1} v_i \cdot c_i = (\alpha_1, d_1, \dots, d_{g-1})^t.$$

Hence, since α_1 left-divided every element in $\{v_{1,1}, \dots, v_{g+1,1}\}$, by linear combinations of the v_i and w_1 we can write that: $M_\pi = \langle v'_1, \dots, v'_{g+1}, w_1 \rangle_{\mathcal{O}_0[\frac{1}{\pi}]}$ where the v'_i are of the form: $v'_i = (0 \ v'_{i,2} \ \dots \ v'_{i,g})^t$. By the same argument there exists an $\mathcal{O}_0[\frac{1}{\pi}]$ -vector w_2 , such that $M_\pi = \langle v''_1, \dots, v''_{g+1}, w_1, w_2 \rangle_{\mathcal{O}_0[\frac{1}{\pi}]}$, where this time the v''_i are of the form: $v''_i = (0 \ 0 \ v''_{i,3} \ \dots \ v''_{i,g})^t$. Repeating this process leads to g vectors w_i such that $M_\pi = \langle 0, \dots, 0, w_1, \dots, w_g \rangle_{\mathcal{O}_0[\frac{1}{\pi}]}$, which concludes.

Our algorithm only involves linear algebra and g calls to KLPT. It implies the complexity statement. $\square$

Algorithm 2: PRINCIPALIZEALMOSTEVERYWHERE

Input: An integral right $\mathcal{O}$ -ideal I given by n generators, and a set S of small primes distinct from p

Output: An $\alpha \in I[\frac{1}{\pi}]$ such that $I[\frac{1}{\pi}] = \alpha \mathcal{O}[\frac{1}{\pi}]$

- 1 Denote by L_v the list of the column vectors of the generators of I ;
- 2 Denote by $L_{res} := []$ what will be the columns of α ;
- 3 **for** i_{row} *in* $range(g)$ **do**
- 4 Write $L_v = (v_i)_{i \in [1, ng]}$;
- 5 Let I_{row} be the right $\mathcal{O}$ -ideal generated by $(v_i[i_{row}])_i$;
- 6 Call KLPT to compute $\beta \in I_{row}[\frac{1}{\pi}]$ such that $I_{row}[\frac{1}{\pi}] = \beta \mathcal{O}_0[\frac{1}{\pi}]$;
- 7 Compute the $c_i \in \mathcal{O}_0[\frac{1}{\pi}]$ such that $\sum v_i[i_{row}]c_i = \beta$;
- 8 Append $w = \sum v_i c_i$ to the list L_{res} ;
- 9 Replace $L_v = (v_i - w \cdot (\beta^{-1} v_i[i_{row}]))_i$;
- 10 **end**
- 11 Return the matrix whose columns are the elements of L_{res} .

4.2. Solving the problem locally at every split $\ell \in S$. The only obstruction left for I to be principal lies locally at primes $\ell \in S$, since we found a generator $\alpha \in I[\frac{1}{\pi}]$ suitable at each place where they are invertible. Let us denote $J := \alpha^{-1}I$, being an $\mathcal{O}$ -ideal. We aim now to trivialize J by acting by elements of $\mathcal{O}[\frac{1}{\pi}]^\times$ to preserve the fact that $J[\frac{1}{\pi}] = \mathcal{O}[\frac{1}{\pi}]$. In the following we fix an $\ell \in S$ that splits in $\mathcal{O}_0$, so that $\mathcal{O}[\frac{1}{\ell}]^\times \subset \mathcal{O}[\frac{1}{\pi}]^\times$. We fix an isomorphism $\mathcal{O} \otimes \mathbb{Z}_\ell \rightarrow \mathcal{M}_{2g}(\mathbb{Z}_\ell)$, that naturally extend to an isomorphism $\iota: \mathcal{O} \otimes \mathbb{Q}_\ell \rightarrow \mathcal{M}_{2g}(\mathbb{Q}_\ell)$, and denote by $\mathcal{O}_m$ the image of $\mathcal{O}$ by ι . We know that $J \otimes \mathbb{Z}_\ell$ embeds in $\mathcal{M}_{2g}(\mathbb{Q}_\ell)$ as a fractional principal ideal of

$\mathcal{M}_{2g}(\mathbb{Z}_\ell)$ denoted J_ℓ . Thus there exists $\beta_\ell \in \mathcal{M}_{2g}(\mathbb{Z}_\ell)$ such that $\ell^r J_\ell = \beta_\ell \mathcal{M}_{2g}(\mathbb{Z}_\ell)$, for some $r \in \mathbb{Z}_{\geq 0}$.

Note that every permutation matrix is invertible in $\mathcal{O}$. In addition, for $x \in \mathcal{O}_0$ and distinct $i, j \in \llbracket 1, g \rrbracket$, we have $Id_g + E_{ij}x \in \mathcal{O}^\times$ where $(E_{ij})_{i,j \in \llbracket 1, g \rrbracket}$ denotes the canonical basis of $\mathcal{M}_g(\mathbb{Q})$. Therefore we can act on β_ℓ by the subgroup $G \subset \mathcal{O}_m^\times$ generated by elements of the form $P \otimes Id_2$ and $Id_{2g} + (E_{ij} \otimes M)$, where P is a permutation matrix, $M \in \mathcal{M}_2(\mathbb{Z}_\ell)$, $i \neq j$ and $\otimes$ denotes the usual tensor product for matrices.

Lemma 4.5. *The subgroup $G \subset \mathcal{O}_m^\times$ acts transitively on $\mathbb{P}^{2g-1}(\mathbb{F}_\ell)$.*

Proof. Let $x := [x_{11}, x_{12}, \dots, x_{g1}, x_{g2}]$ and $y := [y_{11}, y_{12}, \dots, y_{g1}, y_{g2}]$ be two points of $\mathbb{P}^{2g-1}(\mathbb{F}_\ell)$. First by acting on x and y by the permutations $P \otimes Id_2$, we may assume that $(x_{11}, x_{12}) \neq (0, 0)$ and that $(y_{g1}, y_{g2}) \neq (0, 0)$. Thus we can suppose that $x_{11} \neq 0$ or $x_{12} \neq 0$, without loss of generality. We reduce to the case where $x_{11} \neq 0$, the other case being symmetric. We normalize $x = [1, x'_{12}, \dots, x'_{g1}, x'_{g2}]$. We can now construct the following matrix:

$$M_1 := \left(Id_{2g} + E_{21} \otimes \begin{pmatrix} y_{21} - x'_{21} & 0 \\ y_{22} - x'_{22} & 0 \end{pmatrix} \right) \times \dots \times \left(Id_{2g} + E_{g1} \otimes \begin{pmatrix} y_{g1} - x'_{g1} & 0 \\ y_{g2} - x'_{g2} & 0 \end{pmatrix} \right) \\ = \begin{pmatrix} Id_2 & 0 & 0 & \dots & 0 \\ \begin{pmatrix} y_{21} - x'_{21} & 0 \\ y_{22} - x'_{22} & 0 \end{pmatrix} & Id_2 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ \begin{pmatrix} y_{g1} - x'_{g1} & 0 \\ y_{g2} - x'_{g2} & 0 \end{pmatrix} & 0 & 0 & \dots & Id_2 \end{pmatrix} \in G,$$

such that $M_1 \cdot x = [1, x'_{12}, y_{21}, y_{22}, \dots, y_{g1}, y_{g2}]$. By a similar argument on y_{g1} and y_{g2} , we find $M_2 \in G$ such that $M_2 \cdot [1, x'_{12}, y_{21}, y_{22}, \dots, y_{g1}, y_{g2}] = [y_{11}, y_{12}, \dots, y_{g1}, y_{g2}]$. We conclude that there exists $M \in G$ such that $M \cdot x = y$. $\square$

On the other hand, $J[\frac{1}{\pi}]$ is invariant by the action of elements of $\mathcal{O}[\frac{1}{\ell}]^\times$. Let us denote by $D_\ell \in \mathcal{M}_{2g}(\mathbb{Q}_\ell)$ the matrix image of $D := \begin{pmatrix} u & 0 \\ 0 & Id_{g-1} \end{pmatrix} \in \mathcal{O}[\frac{1}{\ell}]^\times$, where u is a uniformizer at ℓ which splits in $\mathcal{O}_0$.

Lemma 4.6. *Let $A \in \mathcal{M}_{2g}(\mathbb{Z}_\ell)$. A is left divisible by D_ℓ in $\mathcal{M}_{2g}(\mathbb{Z}_\ell)$ if and only if $\overline{A}$ is left divisible by $\overline{D}_\ell$ in $\mathcal{M}_{2g}(\mathbb{F}_\ell)$, where $\overline{\cdot}$ is the canonical projection.*

Proof. First, note that $\ell D^{-1} \in \mathcal{O}$, and $\det(D_\ell) = \ell$. It is clear that the divisibility in $\mathcal{M}_{2g}(\mathbb{F}_\ell)$ is necessary. Let us prove the converse. By assumption, there exists $Y \in \mathcal{M}_{2g}(\mathbb{F}_\ell)$ such that $\overline{A} = \overline{D}_\ell Y$. Taking any representative for the class Y leads to $X, R \in \mathcal{M}_{2g}(\mathbb{Z}_\ell)$, such that $A = D_\ell X + \ell R$. Thus $D_\ell^{-1} A = X + \ell D_\ell^{-1} R$, but by the adjugate matrix formula we see that $\ell D_\ell^{-1} \in \mathcal{M}_{2g}(\mathbb{Z}_\ell)$. Hence $D_\ell^{-1} A \in \mathcal{M}_{2g}(\mathbb{Z}_\ell)$ which concludes the proof. $\square$

Our goal is now to act on β_ℓ by D_ℓ and by elements of G , until we reach an element γ_ℓ such that $v_\ell(\det(\gamma_\ell)) = 0$. Indeed, this action preserves the fact that I is generated locally almost everywhere by α , and acting by D_ℓ^{-1} allows us to decrease step by step the ℓ -valuation of the determinant of the generator β_ℓ .

Proposition 4.7. *There exists $\delta_\ell \in \iota(\mathcal{O}[\frac{1}{\ell}]^\times)$ such that $\delta_\ell J_\ell = \mathcal{M}_{2g}(\mathbb{Z}_\ell)$.*

Proof. By Lemma 4.6, checking whether β_ℓ is divisible by D_ℓ reduces to checking it on $\mathcal{M}_{2g}(\mathbb{F}_\ell)$. And since we are now over a field, a matrix $A \in \mathcal{M}_{2g}(\mathbb{F}_\ell)$ is left divisible by $\overline{D_\ell}$ if and only if $\text{Im}(A) \subset \text{Im}(\overline{D_\ell})$. Using the Hermite normal form and the fact that $\det(D_\ell) = \ell$, we see that $\text{rk}(\overline{D_\ell}) = 2g - 1$. It implies that $\text{Im}(\overline{D_\ell})$ is a hyperplane, say $v^\perp$, the orthogonal of a vector v .

From now on we suppose that $v_\ell(\det(\beta_\ell)) > 0$, else there is nothing to prove. Thus $\overline{\beta_\ell}$ is singular, and $\text{Im}(\overline{\beta_\ell})$ is contained in a hyperplane, say $w^\perp$. Note that v and w are well defined up to scaling. Hence Lemma 4.5 ensures that there exists $M_1 \in G$ such that $M_1 v = w$. Thus $\text{Im}(\overline{\beta_\ell}) \subset w^\perp = (M_1 v)^\perp = M_1^{-1t}(v)^\perp$, and finally $\text{Im}(M_1^t \overline{\beta_\ell}) \subset \text{Im}(\overline{D_\ell})$. Noticing that G is stable under transposition, we rewrite the previous equality as $\text{Im}(M_1 \overline{\beta_\ell}) \subset \text{Im}(\overline{D_\ell})$ for $M_1 \in G$. Thus there exists $\beta_\ell^{(0)} \in \mathcal{M}_{2g}(\mathbb{Z}_\ell)$ such that $M_1 \beta_\ell = D_\ell \beta_\ell^{(0)}$, which leads to $v_\ell(\det \beta_\ell^{(0)}) = v_\ell(\det \beta_\ell) - 1$. Repeating this process $v_\ell(\det(\beta_\ell))$ times leads to a matrix $M \in \mathcal{M}_{2g}(\mathbb{Q}_\ell)$ such that $M \beta_\ell = \gamma_\ell \in \text{GL}_{2g}(\mathbb{Z}_\ell)$. Then by taking $\delta_\ell := \ell^r M$, we have $\delta_\ell J_\ell = \gamma_\ell \mathcal{M}_{2g}(\mathbb{Z}_\ell) = \mathcal{M}_{2g}(\mathbb{Z}_\ell)$.

It remains to prove that $\delta_\ell \in \iota(\mathcal{O}[\frac{1}{\ell}]^\times)$. First notice that δ_ℓ is constructed as a product of elements of G , ℓ^r , and inverses of D_ℓ . But $G \subset \mathcal{O}_m^\times = \iota(\mathcal{O}^\times)$, and $D_\ell = \iota(D)$. Since we saw that $D \in \mathcal{O}[\frac{1}{\ell}]^\times$, it concludes the proof. $\square$

Proposition 4.8. *Algorithm 3 takes as input a right $\mathcal{O}$ -ideal J and an element $u \in \mathcal{O}_0$ such that its norm $\ell := \text{Nrd}(u)$ is a small prime and outputs a $\delta_\ell \in \iota(\mathcal{O}[\frac{1}{\ell}]^\times)$ as in Proposition 4.7. Moreover that algorithm runs in polynomial time in the size of the inputs and in $\log(p)$.*

Proof. The correctness follows from Proposition 4.7 and its proof, noting that the chosen ℓ -adic precision ensures that all the computations of valuations and reductions modulo ℓ are performed correctly. The other operations, derived from $\mathbb{Z}_\ell$ -linear algebra at polynomial precision, lead to a polynomial time algorithm. $\square$

4.3. Local to global and conclusion. In this section we describe how to finally recover the generator of I . Since I is in particular a $\mathbb{Z}$ -lattice, we will apply the following local-global principle:

Theorem 4.9. *[Voi21, Corollary 9.4.7] Let M, N two R -lattices for R a Dedekind domain. Then $M = N$ if and only if $M \otimes R_{\mathfrak{p}} = N \otimes R_{\mathfrak{p}}$ for all primes $\mathfrak{p}$ of R .*

Theorem 4.10. *On input a right $\mathcal{M}_g(\mathcal{O}_0)$ -ideal I , Algorithm 4 outputs a generator of I . If $\mathcal{O}_0$ contains a (given) quadratic order $\mathcal{Q}$ of small discriminant Δ , and under GRH, this algorithm runs in polynomial time in the size of the input and in $\log(p)$.*

Proof. Let us prove the correctness first. Suppose that ℓ is a small prime and $u \in \mathcal{O}_0$ such that $\text{Nrd}(u) = \ell$. Applying Algorithm 2 for $S = \{\ell\}$, we can compute $\alpha \in I[\frac{1}{\ell}]$ such that $I[\frac{1}{\ell}] = \alpha \mathcal{O}[\frac{1}{\ell}]$. Thus if $J := \alpha^{-1} I$ then for every prime $q \neq \ell$, we have the completion $J \otimes \mathbb{Z}_q = \mathcal{O} \otimes \mathbb{Z}_q$, since ℓ is invertible in $\mathbb{Z}_q$. Now using Algorithm 3, we can compute $\delta_\ell \in \iota(\mathcal{O}[\frac{1}{\ell}]^\times)$ such that $\mathcal{M}_{2g}(\mathbb{Z}_\ell) = \delta_\ell \iota(J \otimes \mathbb{Z}_\ell)$. Denoting $\delta = \iota^{-1}(\delta_\ell)$ yields to $\delta \in \mathcal{O}[\frac{1}{\ell}]^\times$ and $\mathcal{O} \otimes \mathbb{Z}_\ell = \delta(J \otimes \mathbb{Z}_\ell) = (\delta J) \otimes \mathbb{Z}_\ell$. Since $\delta \in \mathcal{O}[\frac{1}{\ell}]^\times$, we notice that $\delta J[\frac{1}{\ell}] = \delta \mathcal{O}[\frac{1}{\ell}] = \mathcal{O}[\frac{1}{\ell}]$, and for every prime $q \neq \ell$ we still have $\delta J \otimes \mathbb{Z}_q = \mathcal{O} \otimes \mathbb{Z}_q$.

Putting everything together we have that $\delta J \otimes \mathbb{Z}_q = \mathcal{O} \otimes \mathbb{Z}_q$ for every prime q . Then by Theorem 4.9, $\mathcal{O} = \delta J = \delta \alpha^{-1} I$, and finally $I = \alpha \delta^{-1} \mathcal{O}$.

Algorithm 3: PRINCIPALIZEAT ℓ

Input: A fractional right $\mathcal{O}$ -ideal J given by n generators, an element $u \in \mathcal{O}_0$ such that its norm $\ell := \text{Nrd}(u)$ is a small prime
Output: A matrix $\delta_\ell \in \iota(\mathcal{O}[\frac{1}{\ell}]^\times)$ such that $\delta_\ell J \otimes \mathbb{Z}_\ell = \mathcal{M}_{2g}(\mathbb{Z}_\ell)$

- 1 Compute $r = v_\ell(d)$ where d is the denominator of J ;
 // so that $\ell^r J \otimes \mathbb{Z}_\ell \subset \mathcal{M}_{2g}(\mathbb{Z}_\ell)$
- 2 Compute $e = v_\ell([\mathcal{O} : dJ])$;
 // we have $e = v_\ell([\mathcal{M}_{2g}(\mathbb{Z}_\ell) : \ell^r J \otimes \mathbb{Z}_\ell])$
- 3 Perform the following ℓ -adic computations at absolute precision $e + 1$;
- 4 Compute the images of i, j, k in $\mathcal{M}_2(\mathbb{Z}_\ell)$ by Hensel lifting ;
- 5 Deduce the images $\alpha_1, \dots, \alpha_n$ of the generators of $\ell^r J$ in $\mathcal{M}_{2g}(\mathbb{Z}_\ell)$;
- 6 Compute their greatest common divisor β_ℓ with a Hermite Normal Form ;
 // Here we know that $\ell^r J \otimes \mathbb{Z}_\ell \simeq \beta_\ell \mathcal{M}_{2g}(\mathbb{Z}_\ell)$
- 7 Construct D_ℓ thanks to u ;
- 8 Compute $v \in \mathbb{F}_\ell^{2g}$ such that $\text{Im}(\overline{D}_\ell) = v^\perp$;
- 9 Denote $\beta_\ell^{(-1)} := \beta_\ell$ and $h := v_\ell(\det(\beta))$;
- 10 **for** i_v **in** $\text{range}(h)$ **do**
 - 11 Compute $w^{(i_v)}$ such that $\text{Im}(\overline{\beta}_\ell^{(i_v-1)}) \subset w^{(i_v)\perp}$;
 - 12 Compute M_{i_v} such that $M_{i_v} v = w^{(i_v)}$ with Lemma 4.5 ;
 - 13 Compute $\beta_\ell^{(i_v)}$ such that $M_{i_v} \beta_\ell^{(i_v-1)} = D_\ell \beta_\ell^{(i_v)}$;
- 14 **end**
- 15 **return** $\ell^r D_\ell^{-1} M_{h-1} D_\ell^{-1} M_{h-2} \dots D_\ell^{-1} M_0$

We now analyse the complexity of the algorithm, assuming that we know a quadratic order $\mathcal{Q}$ of small discriminant Δ embedded in $\mathcal{O}_0$. The cost of finding a small prime ℓ and $u \in \mathcal{Q}$ such that $\text{Nrd}(u) = \ell$ is polynomial in Δ thanks to Cornacchia's algorithm. By assumption $\Delta = O(\log(p))$, therefore the first step is polynomial in $\log(p)$. Finally we proved in Propositions 4.4 and 4.8 that the steps 2 and 3 run in polynomial time under GRH, hence Algorithm 4 also does. $\square$

In our presentation of Algorithm 4 COMPUTEGENERATOR, for clarity we choose to use $S = \{\ell\}$, with ℓ a small prime which is a norm in $\mathcal{O}_0$. But for efficiency reasons we might be interested in using several small different primes as in Algorithm 2, allowing some trade-off. This would be useful for working with 1-dimensionals ideals of powersmooth norm N as in our implementation, instead of smooth norm ideals, ensuring that the N -torsion is accessible.

Let us explain the small changes needed to make it work. Firstly, Step 2 would be a call to Algorithm 2 with $S = \{\ell_1, \dots, \ell_n\}$. Then for every ℓ_i , we would compute the δ_{ℓ_i} iteratively, calling Algorithm 3 on the corresponding u_i , and replacing J by $\delta_{\ell_{i-1}} J$ each time. Denoting δ_i for the preimage of δ_{ℓ_i} , the result would be $\alpha(\delta_n \dots \delta_1)^{-1}$. Recall that, as explained in Section 2.1 and in [CEIV26], Theorem 4.10 leads to a solution of the PIP for arbitrary maximal orders in $\mathcal{M}_g(\mathcal{B}_{p,\infty})$.

Remark 4.11. A solution of the Principal Ideal Problem with usual cryptographic settings comes as a corollary of Theorem 4.10. Indeed for a prime $p \equiv 3 \pmod{4}$ and

the maximal order $\mathcal{O}_0 = \mathbb{Z} + i\mathbb{Z} + \frac{i+j}{2}\mathbb{Z} + \frac{1+k}{2}\mathbb{Z}$, it is easy to find elements $u \in \mathcal{O}_0$ of small prime norm. It follows from the fact that $\mathbb{Z}[i] \subset \mathcal{O}_0$, and thus that every prime ℓ such that $\ell \equiv 1 \pmod{4}$ will be a norm as a sum of squares.

Algorithm 4: COMPUTEGENERATOR

Input: A right $\mathcal{O}$ -ideal I given by generators

Output: An element $\gamma \in \mathcal{O}$ such that $I = \gamma\mathcal{O}$

- 1 Compute an element $u \in \mathcal{Q} \subset \mathcal{O}_0$ such that its norm $\ell := \text{Nrd}(u)$ is a small prime
 - 2 Call PRINCIPALIZEALMOSTEVERYWHERE($I, \{\ell\}$) to compute $\alpha \in I[\frac{1}{\ell}]$ such that $I[\frac{1}{\ell}] = \alpha\mathcal{O}[\frac{1}{\ell}]$ and denote $J = \alpha^{-1}I$;
 - 3 Call PRINCIPALIZEATL(J, u) to compute δ_ℓ such that $\delta_\ell J \otimes \mathbb{Z}_\ell = \mathcal{M}_{2g}(\mathbb{Z}_\ell)$;
 - 4 Recover the preimage of δ_ℓ in $\mathcal{O}[\frac{1}{\ell}]^\times$, denoted δ ;
 - 5 return $\alpha\delta^{-1}$
-

4.4. Consequences for abstract and standard representations. As explained in Proposition 3.9, our solution to the PIP allows us to recover isomorphisms between endomorphism rings, and the distinguished maximal order $\mathcal{M}_g(\mathcal{O}_0) \subset \mathcal{M}_g(\mathcal{B}_{p,\infty})$. We still rely on the GRH, since Algorithm 4 does.

Proposition 4.12. *With the notation of Proposition 3.9. There exists an algorithm that computes an isomorphism $\text{End}(M) \rightarrow \mathcal{M}_g(\mathcal{O}_0)$. Under the GRH, this algorithm is (randomised) polynomial in $\log(p)$.*

Remark that if we start with a refined abstract representation on $\text{End}(A)$, up to conjugating φ by the outer automorphism we can assume that φ is refined, i.e. respects the $\mathbb{F}_{p^2}$ -algebra structure. Then we apply Remark 3.4, to recover the associated Hermitian module $(\mathcal{O}_0^g, S)$, where S is the Gram matrix of the Hermitian form. The following theorem sums up the translations provided by Section 3, Proposition 4.12, and our solution to the PIP. An illustration of these can be found in Appendix A, Figure 1.

Theorem 4.13. *Under GRH, there are polynomial time algorithms to convert between the abstract module representation, the refined abstract endomorphism ring representation, the standard endomorphism representation and the standard module representation.*

5. SWITCHING BETWEEN EXPLICIT AND ABSTRACT REPRESENTATIONS

In this section we explain how to switch between explicit and abstract representations. It will be crucial in Section 6.3 to prove that from (M, H_M) we can efficiently recover the corresponding (A, λ_A) together with the explicit action of $\text{End}(A)$ over A , and an unpolarized isomorphism $A \simeq E_0^g$.

By an explicit representation of a principally polarized abelian variety $A/\mathbb{F}_q$, we mean that we are given some theta null point of level 4 on it. This is too specific for our purpose: we could be agnostic on the exact model as long as it allows us to do the basic operations we need (sampling points, computing a basis of the ℓ -torsion for small ℓ , computing pairings, computing isogenies from generators of their kernels, testing if two ppavs are isomorphic and if so computing an isomorphism), but for

the sake of simplicity we stick to the theta null point, for which there exist explicit algorithms in the literature for all of the above, see [Rob21].

By an explicit representation of a morphism $\varphi : A \rightarrow B$, defined over a finite extension k of $\mathbb{F}_q$, we mean that we have an explicit representation of A and B over k , and that we have a data structure that contains a bound M of the size of φ (as defined in Section 5.1), along with an algorithm such that given any point $P \in A(k)$ outputs $\varphi(P)$ in polynomial time in M (in terms of arithmetic over k).

Our main goal in this section is to develop algorithms to switch between abstract representations and explicit representations, both for modules and endomorphism rings.

Remark 5.1. We will need slightly different variants of an explicit representation of a morphism φ . A *sufficiently explicit* representation is a representation where we can evaluate φ only on points $P \in A(\mathbb{F}_q)$ of small prime torsion ℓ (more formally the evaluation $\varphi(P)$ is polynomial in ℓ and $\log q$.)

A *fully explicit* representation of φ allows to evaluate φ on points $P \in A(R)$ for R a local Artinian algebra over $\mathbb{F}_q$ (in time polynomial in $\log \deg R$), not only for points defined over fields. Being able to evaluate φ on such infinitesimal point allows us to recover the action of φ on differentials.

The HD representation ([Rob25, Section 5]) of a polarized isogeny φ between ppavs allows, using the theta isogeny algorithm of [LR22], to evaluate it on infinitesimal points. We then recover in polynomial time a fully explicit representation of φ , as defined above, from a sufficiently explicit representation.

It is not obvious how to extend this to general morphisms. Remarkably, we will see that if $A/\mathbb{F}_{p^2}$ is maximal, then a sufficiently explicit representation of generators φ_i of $M = \text{Hom}(A, E_0)$ allows to recover the Hermitian module (M, H_M) abstractly, and then our higher dimensional Clapots algorithm allows to recover a fully explicit representation of the φ_i , all in polynomial time.

5.1. From explicit to abstract representations. In this section we explain how to recover an abstract module structure from generators of $\text{Hom}(A, B)$ given by explicit isogenies. This generalizes previous algorithms to recover the endomorphism ring of a supersingular elliptic curve to higher dimension and to modules. Our method works for general abelian varieties. As a particular case, when A is maximal supersingular over $\mathbb{F}_{p^2}$, applying the general method to $\text{Hom}(A, E_0)$ or $\text{End}(A) = \text{Hom}(A, A)$ will allow us to recover the abstract module and endomorphism ring representations from explicit ones.

Definition 5.2. For $A, B/k$ explicit principally polarized abelian varieties over a finite field k , an explicit representation of $\text{Hom}_k(A, B)$ is a list $(\gamma_1, \dots, \gamma_r)$ of morphisms, each given via an explicit representation, and such that they generate $\text{Hom}_k(A, B)$ as an abelian group.

A complete representation of $\text{Hom}_k(A, B)$ is an explicit representation as above, as well as a basis of their lattice of relations. In the case where $B = A$, for a complete representation of $\text{End}_k(A)$ we also require explicit matrices that compute the composition.

We remark that a complete representation is both an abstract and an explicit representation, in such a way that we can express an abstract morphism $\varphi \in \text{Hom}_k(A, B)$ in terms of the explicit generators γ_i , so as to obtain an explicit representation of φ . As in Remark 5.1, we also have notions of sufficiently and fully

complete representations of $\text{Hom}_k(A, B)$. We will prove in Theorem 5.8 that given an explicit (resp. sufficiently explicit) representation of $\text{Hom}_k(A, B)$, we can recover a complete representation (resp. sufficiently complete).

Recall that if A/k is an abelian variety over a finite field, we have a well defined notion of trace on $\text{End}(A)$, which can be computed by embedding $\text{End}(A)$ into $\mathcal{M}_{2g}(T_\ell A)$ where $T_\ell A$ is any Tate module, and taking the trace of the associated matrix (which is independent of ℓ , and belongs to $\mathbb{Z}$), see [Mum70, p. 180–182].

Definition 5.3. Let (A, λ_A) and (B, λ_B) be principally polarized abelian varieties over a finite field k . We define a bilinear form on $\text{Hom}_k(A, B)$ via $(\varphi_1 | \varphi_2) = \text{Tr}(\lambda_A^{-1} \circ \varphi_2^\vee \circ \lambda_B \circ \varphi_1)$ where Tr is the (reduced rational) trace from $\text{End}(A)$ to $\mathbb{Z}$, and associated norm $N(\varphi) = \sqrt{(\varphi | \varphi)}$.

Example 5.4. Let $\varphi: (A, \lambda_A) \rightarrow (B, \lambda_B)$ be an n -isogeny between g -dimensional ppavs, which means that $\lambda_A^{-1} \circ \varphi^\vee \circ \lambda_B \circ \varphi = [n]$. Then $N(\varphi) = \sqrt{2gn}$, whereas $\deg \varphi = n^g$. More generally, if $\varphi: A \rightarrow B$ is an isogeny, then by the AM-GM inequality, $\deg \varphi \leq N(\varphi)^{2g}/(2g)^g$. In particular, we can recover $\deg \varphi$ from a bound N on its size in polynomial time (in N and $\log p$) by combining the CRT and Weil pairings and DLPs on points of small order. The main advantage of N compared to the degree is that it gives a well behaved notion of size for morphisms that may not be isogenies. In particular if a morphism $\varphi: A \rightarrow B$ vanishes over $A[m]$ for any $m > N(\varphi)$, then by submultiplicativity (proved in the next proposition) $\varphi = 0$.

Proposition 5.5. *The bilinear form $(\cdot | \cdot)$ is symmetric and positive definite. It satisfies $|(\varphi_1 | \varphi_2)| \leq N(\varphi_1)N(\varphi_2)$, $N(\varphi_1 + \varphi_2) \leq N(\varphi_1) + N(\varphi_2)$, and if $\varphi_1: A \rightarrow B$, $\varphi_2: B \rightarrow C$, $N(\varphi_2 \circ \varphi_1) \leq N(\varphi_1)N(\varphi_2)$.*

Proof. We can embed $\text{Hom}_k(A, B)$ into $\text{End}_k(A \times B)$ as well as $\text{Hom}_k(A, B) \times \text{Hom}_k(B, C)$ into $\text{End}_k(A \times B \times C)$ in a way compatible with composition. If we take the product polarizations, then these embeddings respect the bilinear forms by definition of the product polarization. So we reduce to proving this statement for endomorphism rings. Tensoring with $\mathbb{R}$, arguing as in [Mum70, Theorem 6 p. 208] we reduce to the case of a ring $\mathcal{M}_g(K)$ where $K = \mathbb{R}, \mathbb{C}, \mathbb{H}$ and the Rosati involution is given by the conjugate transpose $M \mapsto M^*$. So the scalar product is $(M_1, M_2) \mapsto \text{Tr}(M_1^* M_2)$ which is symmetric positive definite, and the associated norm is the Frobenius norm which is submultiplicative, as we wanted. $\square$

Lemma 5.6. *Suppose that we have a sufficiently explicit representation of a morphism $\varphi: (A, \lambda_A) \rightarrow (B, \lambda_B)$. Then it holds for the polarized dual $\tilde{\varphi}: B \rightarrow A$.*

Proof. We want to evaluate $\tilde{\varphi}(P)$ for $P \in B[\ell]$ and ℓ a small prime. We sample a basis $(e_1, \dots, e_{2g})$ of $A[\ell]$. By compatibility of the polarized Weil pairings with morphisms [Mum70, p. 186], we have $e_{\lambda_A}(\tilde{\varphi}(P), e_i) = e_{\lambda_B}(P, \varphi(e_i))$. Since we can compute $\varphi(e_i)$ by assumption on φ , then Weil pairings and DLPs allow to express $\tilde{\varphi}(P)$ in the basis of the e_i in polynomial time in ℓ . $\square$

Lemma 5.7. *Suppose that we have a sufficiently explicit representation of two morphisms $\varphi_1, \varphi_2: (A, \lambda_A) \rightarrow (B, \lambda_B)$. Then we can compute $(\varphi_1 | \varphi_2)$ in polynomial time in their size.*

Proof. We use a CRT method as in Schoof's algorithm. Let ℓ_i be a small prime. By Lemma 5.6, we can compute the action of $\tilde{\varphi}_2 \circ \varphi_1$ on a basis of $A[\ell_i]$, the trace of the associated matrix gives us $(\varphi_1 | \varphi_2)$ modulo ℓ_i . Since $(\varphi_1 | \varphi_2) \leq N(\varphi_1)N(\varphi_2)$

we recover the scalar product using sufficiently many small primes, in polynomial time in $N(\varphi_1), N(\varphi_2)$. $\square$

Theorem 5.8. *If we have explicit representations of generators $\varphi_1, \dots, \varphi_m \in \text{Hom}_k(A, B)$ (as an abelian group), then we can compute the abstract $\mathbb{Z}$ -module structure of $\text{Hom}_k(A, B)$ in polynomial time in the size of the base field, the number of generators, and their size. Given an explicit representation of some other $\varphi \in \text{Hom}_k(A, B)$, we can also recover a decomposition $\varphi = \sum m_i \varphi_i$ of φ .*

If we also have explicit representations of generators of $\text{Hom}_k(B, C)$, then we can compute the abstract structure of the composition map $\text{Hom}_k(A, B) \otimes \text{Hom}_k(B, C) \rightarrow \text{Hom}_k(A, C)$.

In particular, if we have explicit representations of generators γ_i of $\text{End}_k(A)$ (resp. of $\text{End}_k(B)$, resp. of both), then we can compute the ring structure of $\text{End}_k(A)$ via the multiplication map $\text{End}_k(A) \otimes \text{End}_k(A) \rightarrow \text{End}_k(A)$, along with the right $\text{End}_k(A)$ -module structure of $\text{Hom}_k(A, B)$ (resp. its left $\text{End}_k(B)$ -module, resp. its $(\text{End}_k(B), \text{End}_k(A))$ -bimodule structure).

Proof. Let $M = \text{Hom}_k(A, B)$. Since the φ_i are generators of M by assumption, and the bilinear form from Proposition 5.5 is positive definite, $\psi \in M \mapsto (\psi | \varphi_i) \in \mathbb{Z}^m$ is an embedding, which we can explicitly compute by Lemma 5.7. Applying this to $\psi = \varphi_i$ allows to recover the Gram matrix $(\varphi_i | \varphi_j)$. Applying this to the composition of two explicit morphisms, this also allows to recover the ring and module structures. $\square$

Example 5.9. Recall that $\varphi \in \text{Hom}_k(A, B) \mapsto \tilde{\varphi} \in \text{Hom}_k(B, A)$ is an isomorphism of abelian groups. If we have a sufficiently explicit representation of generators γ_i of $\text{Hom}_k(A, B)$, Lemma 5.6 also gives us sufficiently explicit representation of the generators $\tilde{\gamma}_i$ of $\text{Hom}_k(B, A)$. In particular, using Theorem 5.8, given an explicit representation of $\text{End}(B)$ we can recover the full endomorphism $\gamma_j^\vee \circ \gamma_i \in \text{End}(B)$ rather than just its trace. As an application, this allows to recover the abstract representation of the Rosati involution on $\text{End}(A)$ (using $B = A$), or the Hermitian form associated with a module $M = \text{Hom}_k(A, E_0)$ (using $B = E_0$).

Remark 5.10. Given (A, λ_A) , to recover the action of $\text{End}(A)$ on the differentials, we need a fully explicit representation. If we are only given a sufficiently explicit representation of $\text{End}(A)$, by Section 5.1 we can still build an abstract representation of $\text{End}(A)$, but not a refined representation. Hence, using Section 3 to recover the module representation, we obtain both the module (M, H_M) associated with A , and the one associated with its Galoisian conjugate A^σ .

Then to distinguish between the two, we need to use HD-clapotis to compute the abelian variety associated with each module, and test which one is isomorphic to A . In the end we obtain, in a very circumvoluted manner, a way to turn a sufficiently explicit representation of $\text{End}(A)$ into an Hermitian module representation. We will see in Example 6.8 that (M, H_M) allows to furthermore recover a *fully* explicit representation.

5.2. The refined abstract representation of $\text{End}(A)$. Let us highlight the link between our refined representation and the work of [Mon26] to get ride of the ambiguity carried by an abstract endomorphism ring representation.

Proposition 5.11. *Fix ω_0 a global differential on E_0 . Then the morphism $\Phi : \varphi \in M = \text{Hom}_{\mathbb{F}_{p^2}}(A, E_0) \mapsto \varphi^* \omega_0$ induces an $\mathbb{F}_{p^2}$ -isomorphism $M/\mathfrak{p}M \simeq H^0(A, \Omega_{A/k}^1)$.*

Proof. Since $\Omega_{E_0/\mathbb{F}_{p^2}}$ is of dimension 1, the differential $\varphi^*\omega_0$ is non zero precisely when φ is not a smooth morphism. But this is the case if and only if $\varphi^\vee$ factors through the infinitesimal subgroup $E_0[\pi_p]$, hence the kernel of Φ is $\mathfrak{p}M$, and we have a monomorphism $M/\mathfrak{p}M \rightarrow \Omega_{A/k}^1$. Since both modules have rank $2g$ over $\mathbb{F}_p$ this is an isomorphism. We remark that since the action of $\mathcal{O}_0$ onto ω_0 gives an isomorphism $\mathcal{O}_0/\mathfrak{p}\mathcal{O}_0 \simeq \Omega_{E_0/\mathbb{F}_{p^2}} \simeq \mathbb{F}_{p^2}$ and our morphisms φ are defined over $\mathbb{F}_{p^2}$, Φ is indeed compatible with the $\mathbb{F}_{p^2}$ structure on both M and $\Omega_{A/k}^1$. $\square$

Remark 5.12. The isomorphism above is a special case of a more general isomorphism of Dieudonné modules $\mathbb{D}(A[p]) \simeq \mathbb{D}(E_0[p]) \otimes_{\mathcal{O}_0} M$. Here we endow $\mathbb{D}(E_0[p])$ with its natural $(\mathbb{Z}_q\{F, V\}, \mathcal{O}_0)$ -bimodule structure coming from the natural action of $\mathcal{O}_0$ on E_0 , where $\mathbb{Z}_q\{F, V\}$ is the usual non commutative Dieudonné ring.

Indeed, taking a presentation $\mathcal{O}_0^m \rightarrow \mathcal{O}_0^n \rightarrow M \rightarrow 0$ of M , we have by the module action $0 \rightarrow A \rightarrow E_0^n \rightarrow E_0^m$, and since the Dieudonné module is a contravariant equivalence of categories on p -divisible group, we obtain $\mathbb{D}(E_0^m[p]) \rightarrow \mathbb{D}(E_0^n[p]) \rightarrow \mathbb{D}(A[p]) \rightarrow 0$. On the other hand, taking tensor products we have $\mathbb{D}(E_0[p]) \otimes_{\mathcal{O}_0} \mathcal{O}_0^m \rightarrow \mathbb{D}(E_0[p]) \otimes_{\mathcal{O}_0} \mathcal{O}_0^n \rightarrow \mathbb{D}(E_0[p]) \otimes_{\mathcal{O}_0} M \rightarrow 0$, whence the conclusion using the natural isomorphism $\mathbb{D}(E_0^m[p]) \simeq \mathbb{D}(E_0[p])^m \simeq \mathbb{D}(E_0[p]) \otimes_{\mathcal{O}_0} \mathcal{O}_0^m$.

Now by [Oda69, Corollary 5.11], we have a canonical isomorphism of $\mathbb{D}(A[p])[F]$ and of the global differentials Ω_A^1 .

5.3. From abstract to explicit representations. In this section, we give a general framework to turn an abstract endomorphism representation $\text{Hom}_k(A, B)$ into an explicit (and even full) representation. This relies on being able to construct suitable double paths, as introduced in [DLRW24].

Usually, a double path refers to a couple $\varphi_1, \varphi_2 : (A_1, \lambda_1) \rightarrow (A_2, \lambda_2)$ of two (explicit) isogenies of polarized coprime degrees D_1, D_2 . But we can extend this definition, by also considering polarized $(d_1, \dots, d_r)$ -isogenies:

Definition 5.13. For $i \in \llbracket 1, r \rrbracket$, let (A_i, λ_i) be principally polarized abelian varieties. Let $\varphi : A_1 \times \dots \times A_r \rightarrow B$ be an isogeny, where (B, λ_B) is another ppav. We say that φ is a $(d_1, \dots, d_r)$ -isogeny between the ppav $(A_1 \times \dots \times A_r, \lambda_1 \otimes \dots \otimes \lambda_r)$ and (B, λ_B) , for $d_i \in \mathbb{Z}_{\geq 1}$, when:

$$\tilde{\varphi} \circ \varphi = \text{diag}([d_1], \dots, [d_r]) = \begin{pmatrix} [d_1] & 0 & 0 \\ 0 & \ddots & 0 \\ 0 & 0 & [d_r] \end{pmatrix} \in \text{End}(A_1 \times \dots \times A_r).$$

This notion of $(d_1, \dots, d_r)$ -isogenies naturally extend to similitudes between Hermitian modules. Note that when $r = 1$ or $d_1 = \dots = d_r$, we recover the notion of d -isogenies.

Definition 5.14. With the notation of Definition 5.13, we say that a pair (φ_1, φ_2) consisting of φ_1 a $(d_1, \dots, d_r)$ -isogeny and φ_2 a $(e_1, \dots, e_r)$ -isogeny, both from $A_1 \times \dots \times A_r$ to B , is a *double path*, when the nonnegative integers $\text{lcm } d_i$ and $\text{lcm } e_i$ are coprime.

To emphasize when the double path is given by d -isogenies, we will use the terminology *scalar double path* (since in that case $\tilde{\varphi}_i \circ \varphi_i \in \mathbb{Z}$).

The splitting lemma, at the core of the Clapotis algorithm in dimension 1, gives a nice way to construct a scalar double path between principally polarized abelian varieties, see [Rob25, Propositions 6.4 and 6.11]:

Lemma 5.15. *Suppose that we have an explicit description of an endomorphism $\gamma : A_1 \rightarrow A_1$, such that γ factorizes as $\gamma = \tilde{\varphi}_2 \circ \varphi_1$ for two n_i -isogenies $\varphi_1, \varphi_2 : A_1 \rightarrow A_2$ where n_1 is coprime to n_2 (A_1 and A_2 principally polarized). Then we can efficiently compute an explicit representation of A_2 along with explicit representations of φ_1, φ_2 , and their duals.*

Such double paths play a central role in this work. As an initial illustration, the following shows that they enable the computation of complete representations of homomorphism sets, only from their group structure.

Proposition 5.16. *Let $(A_i, \lambda_i), (B_i, \mu_i)$ be principally polarized abelian varieties, for $i \in \{1, 2\}$. Suppose that we have:*

- *An abstract representation of $\text{Hom}_k(A_1, B_1)$ (as a $\mathbb{Z}$ -module),*
- *Two scalar double paths, $(u_1, u_2) : (A_1, \lambda_1) \rightarrow (A_2, \lambda_2)$ and $(v_1, v_2) : (B_1, \mu_1) \rightarrow (B_2, \mu_2)$,*
- *A sufficiently complete representation³ of $\text{Hom}_k(A_2, B_2)$,*
- *And the abstract representation of the four maps $\text{Hom}_k(A_1, B_1) \rightarrow \text{Hom}_k(A_2, B_2), \varphi \mapsto v_j \circ \varphi \circ \tilde{u}_i$.*

Then we can compute a sufficiently complete representation of $\text{Hom}_k(A_1, B_1)$, in polynomial time in the input.

Proof. Given $P \in A_1(\mathbb{F}_q)$, and $\varphi \in \text{Hom}_k(A_1, B_1)$, we want to compute $\varphi(P)$. We can find a multiple N of the order of P by point counting.

Let d_1, d_2 be the polarized degrees of u_1, u_2 . Since they are coprime, taking suitable gcds, we can write $N = N_1 N_2$ where N_1, N_2 are coprime, N_1 is coprime to d_2 , and N_2 is coprime to d_1 . By the CRT we can write $P = P_1 + P_2$ where P_1 (resp. P_2) is of order dividing N_1 (resp. N_2). We now explain how to compute $\varphi(P_1)$, the case $\varphi(P_2)$ being symmetric.

Let e_1, e_2 be the polarized degrees of v_1, v_2 . Splitting further, we can decompose $P_1 = Q_1 + Q_2$, where this time Q_1 is of order coprime to e_2 , and Q_2 of order coprime to e_1 . Once again, we only describe how to compute $\varphi(Q_1)$, the other cases can be treated similarly.

We know that Q_1 is of order coprime to both d_2 and e_2 . In particular we can first compute $S \in A_2$ such that $S = u_2(Q_1/d_2)$, and hence $\tilde{u}_2(S) = Q_1$. We then compute $T := v_2 \circ \varphi \circ \tilde{u}_2(S) = v_2(\varphi(Q_1))$, and we recover $\varphi(Q_1) = \tilde{v}_2(T)/e_2$, by coprimality with e_2 . $\square$

Remark 5.17. The same techniques hold if, rather than a double path between A_1, A_2 and B_1, B_2 , we can construct explicit embeddings $i_1 : A_1 \rightarrow A_2, i_2 : B_1 \rightarrow B_2$ along with explicit sections, and where we can make $\text{Hom}_k(A_2, B_2)$ explicit. A typical example, that we highlight below, is to take powers of abelian varieties: $A_2 = A_1^r, B_2 = B_1^r$, for some integer $r \geq 1$. Suppose that we want to evaluate an abstract morphism $\varphi : A_1 \rightarrow B_1$ on a point $P \in A_1$. Denote by ι_B the diagonal embedding $B_1 \hookrightarrow B_1^r$, and by π_A (resp. π_B) the projection on the first factor $A_1^r \twoheadrightarrow A_1$ (resp. $B_1^r \twoheadrightarrow B_1$). By assumption on $\text{Hom}(A_1^r, B_1^r)$, we can evaluate the map $\iota_B \circ \varphi \circ \pi_A$ on the point $(P, \dots, P) \in A_1^r$, to compute the corresponding $T \in B_1^r$. Then $\pi_B(T)$ gives us $\varphi(P)$.

³By Theorem 5.8 it suffices to have an explicit representation of $\text{Hom}_k(A_2, B_2)$.

The Zarhin's trick allows us to compute a scalar double path, from a general double path, at the cost of quadrupling the dimension. To lighten notation, we omit the polarizations in the following. When considering a product of ppav, the underlying polarization will always be implicitly given by the product polarization.

Lemma 5.18. *With the same notation as in Definition 5.14. A double path $(\varphi_1, \varphi_2): \prod_i^r A_i \rightarrow B$ induces a scalar double path: $\prod_i^r A_i^4 \rightarrow B^4$.*

Proof. Let us focus on φ_1 , denoting $D = \text{lcm}(d_i)$. Using a diagonal embedding $\varphi_1 \text{Id}_4$, we obtain a $(d_1, \dots, d_g, \dots, d_1, \dots, d_g)$ -isogeny $\prod_i^r A_i^4 \rightarrow B^4$. For every $i \in \llbracket 1, r \rrbracket$, denote by d'_i the cofactor $d'_i := D/d_i$. Using the Zarhin's trick, we can compute a $(d'_1, \dots, d'_g, \dots, d'_1, \dots, d'_g)$ -isogeny $\prod_i^r A_i^4 \rightarrow \prod_i^r A_i^4$. Hence by composition we obtain, a D -isogeny $\Phi_1: \prod_i^r A_i^4 \rightarrow B^4$. The same construction for φ_2 leads to an E -isogeny $\Phi_2: \prod_i^r A_i^4 \rightarrow B^4$, for $E = \prod_i^r e_i$. By definition of the double path (φ_1, φ_2) , D and E are coprime integers. $\square$

We saw in Remark 5.1 that the HD-representation induces fully explicit representations of polarized isogenies. Therefore in Proposition 5.16, if we begin with a *fully* complete representation of $\text{Hom}_k(A_2, B_2)$, then using an HD-representation (which is universal [Rob25, Corollary 5.22]) of the scalar double paths, we deduce a *fully* complete representation of $\text{Hom}_k(A_1, B_1)$. However, this applies only for scalar double paths, since we need an HD representation. We now extend this.

Proposition 5.19. *Proposition 5.16 holds for general double paths (u_1, u_2) and (v_1, v_2) , and in that setting, a fully complete representation of $\text{Hom}_k(A_2, B_2)$ leads to a fully complete representation of $\text{Hom}_k(A_1, B_1)$.*

Proof. By Remark 5.17, it suffices to give a fully complete representation of $\text{Hom}_k(A_1^4, B_1^4)$. Using Lemma 5.18, we recover scalar double paths $A_1^4 \rightarrow A_2^4$ and $B_1^4 \rightarrow B_2^4$, and then we can apply Proposition 5.16 with HD representation, leading to the desired representation. $\square$

Example 5.20. We highlight the case where $A_2 = B_2 = E_0^g$, and $A_1 = B_1 = A$. Suppose that we fixed a fully complete representation of $\text{End}(E_0^g)$ seen as $\mathcal{M}_g(\mathcal{O}_0)$ acting on E_0^g . Then, assuming the construction of double paths $E_0^g \rightarrow A$, Proposition 5.19 enables us to recover a fully complete representation of $\text{End}(A)$.

Remark 5.21. To apply Proposition 5.16 for a point $P \in A_1(R)$ where R is a finite Artinian algebra over $k = \mathbb{F}_q$, we need to know a multiple N of the order of P . In practice, apart from the case where R is a field, we only need the case where $R = \mathbb{F}_q[\epsilon]/(\epsilon^2)$ in Section 5.2, and P is above 0_{A_1} , i.e. gives a tangent vector of A_1 at 0_{A_1} , in which case we can take $N = p$.

We briefly explain the general case. Up to a precomputation only involving R , we may assume that we know an explicit isomorphism $R = \prod R_i$ of R with local Artinian algebras, and reduce to the case where (R, m) is local, with residue field k' . By definition of the formal group Γ associated with A_1 (the formal completion of the zero section in A_1), the kernel of the reduction map $A_1(R) \rightarrow A_1(k')$ is equal to $\text{Hom}(\text{Spec } R, \Gamma)$ (morphisms of formal groups, with R given the discrete topology). Since Γ is a formal Lie group, we have that $\text{Hom}(\text{Spec } R, \Gamma) = \Gamma[m]$, since m consists of the nilpotent elements of R . Since A is smooth, the reduction map is surjective. Hence the reduction map gives an exact sequence

$$0 \rightarrow \Gamma[m] \rightarrow A_1(R) \rightarrow A_1(k') \rightarrow 0.$$

A bound on $\#A_1(k')$ is given by point counting. Finally, $\Gamma[m]$ is a p -group, and if $P \in \Gamma[m]$, $[p] \cdot P \in m^{p^h}$ where $h \geq 1$ is the height of Γ . This gives a bound, depending only on the nilpotency index of m (which is smaller than the dimension of R as a $\mathbb{F}_q$ -vector space), of the order of P .

Example 5.20 motivates us to investigate the construction of such double paths, which is precisely the goal of the next section. Additional applications of this framework will be presented later in Section 6.3.

Remark 5.22. If $\phi : (A_1, \lambda_1) \rightarrow (A_2, \lambda_2)$ is an isogeny between maximal supersingular abelian varieties and (M_1, H_1) the unimodular module associated with (A_1, λ_1) , then ϕ corresponds to a monomorphism $\psi : M_2 \rightarrow M_1$, and we can recover the image of M_2 in M_1 as the elements of M_1 that are zero on $\text{Ker } \phi$ [Rob24, Proposition 4.24]. In particular if we have an explicit representation of ϕ and of M_1 , and that furthermore the degree d of ϕ is smooth and the d -torsion is accessible on A_1 , we can recover M_2 by computing the action of ϕ and generators of M_1 on $A_1[d]$ (as usual thanks to the Weil pairings, DLPs in μ_ℓ for divisors ℓ of d and linear algebra).

Thanks to Theorem 6.9, we can refine this to the case where ϕ is an N -isogeny where N is smooth but the N -torsion may not be accessible. We split ϕ into a product of powersmooth isogenies: $\phi = \phi_m \circ \cdots \circ \phi_1$. We recover the kernel of $\phi_1 : A_1 \rightarrow A_{2,1}$ as above, this gives the submodule $M_2 \subset M_{2,1} \subset M_1$, and we then build a double path to the codomain of ϕ_1 from the knowledge of $M_{2,1}$, to find the explicit module action of $M_{2,1}$ on $A_{2,1}$. We can evaluate the reminder of the isogeny $\phi_m \circ \cdots \circ \phi_2$ on any point of order prime to $\deg \phi_1$, which is enough to recover it. This allows us to proceed recursively to find the chain $M_2 = M_{2,m} \subset \cdots \subset M_{2,1} \subset M_1$. We refer to [CDK+25, § 5.2] for more details.

6. CONSTRUCTING DOUBLE PATHS

In this section we show how to build a double path $E_0^g \rightarrow A$, given an abstract module representation (M, H_M) of (A, λ_A) a maximal supersingular abelian variety of dimension g . Unless stated otherwise, we assume $g \geq 2$. Because we want to apply the theory in the commutative case and not only the maximal supersingular case, in this section we let R be either a maximal quadratic imaginary order or a maximal quaternionic order, and K its fraction field. In both case, we recall that R carries a canonical unimodular Hermitian form defined by $H_0(x, y) = x\bar{y}$. We denote by $r_{\mathbb{Z}}$ the rank of R as a $\mathbb{Z}$ module, being either 2 or 4. Proposition 6.1 lies at the heart of our approach. Corollaries 6.3 will help us improve KLPT² in Section 7.

6.1. The Poincaré decomposition for modules.

Proposition 6.1. *Let (M, H_M) be a unimodular Hermitian module over R (torsion free finitely generated as always). Let $V = M \otimes_{\mathbb{Z}} \mathbb{Q}$, and $H = H_M \otimes_{\mathbb{Z}} \mathbb{Q}$ the induced Hermitian form on V . Let $\iota_1 : M_1 \hookrightarrow M$ be a primitive submodule. Let V_1 be the K -span of M_1 inside V , and $M_1^{\sharp}$ the R -orthogonal of M_1 in V_1 . Let $V = V_1 \oplus^{\perp} V_2$ be the orthogonal decomposition given by V and V_1 , and denote by $M_2 = M \cap V_2$ the primitive submodule of M induced by V_2 . Then $M_1^{\sharp}/M_1 \simeq M_2^{\sharp}/M_2$.*

Proof. This follows from the polarized Poincaré decomposition for abelian varieties ([BL04, § 5.3, § 12.1]), and the anti-equivalence of categories Theorem 2.5. For

convenience, we give a pure module-theoretic proof. We will prove that $M/(M_1 + M_2) \simeq M_1^\sharp/M_1$. By symmetry, we deduce $M_2^\sharp/M_2 \simeq M/(M_1 + M_2) \simeq M_1^\sharp/M_1$. Note that M_2 is, by construction, a primitive submodule of M . Indeed, if $m \in M \setminus M_2$ and $n \in \mathbb{Z} \setminus \{0\}$ are such that $n \cdot m \in M_2$, then $\frac{1}{n}n \cdot m = m \in M_2$ since V_2 is a K -vector space, yielding a contradiction.

Using the natural isomorphism $M^\sharp \simeq M^\vee$ induced by H , we have a natural restriction map $i_1^\sharp : M = M^\sharp \rightarrow M_1^\sharp$, dual to the inclusion $i_1 : M_1 \rightarrow M$, and whose kernel is given by the elements of M that are orthogonal to M_1 , which is precisely M_2 by definition.

On the other hand since M_1 is primitive in M , M/M_1 is torsion free, so projective under our hypothesis that R is maximal. It follows that M splits as $M_1 \oplus M/M_1$, and that $i_1^\sharp$ is surjective. So we obtain an isomorphism $M/M_2 \simeq M_1^\sharp$, which induces $M/(M_1 + M_2) \simeq M_1^\sharp/M_1$. $\square$

Corollary 6.2. *With the notations above, suppose that M is of even rank $g = 2g_0$, that M_1 is of rank g_0 and that $M_1^\sharp = \frac{1}{e}M_1$. Then the natural map $(M_1 \oplus M_2, \frac{1}{e}H_M | M_1 \oplus H_M | M_2) \rightarrow (M, H_M)$ is an e -similitude between unimodular modules.*

Proof. By Proposition 6.1, $M_2^\sharp/M_2 \simeq M_1^\sharp/M_1 \simeq (\mathbb{Z}/e\mathbb{Z})^{r_z g_0}$ as a $\mathbb{Z}$ -module. Since M_2 is of rank $g - g_0 = g_0$, it follows that $M_2^\sharp = \frac{1}{e}M_2$. So $M_1 \oplus M_2$ is unimodular for the product polarization $\frac{1}{e}H_M | M_1 \oplus H_M | M_2$, and the claim follows. $\square$

Corollary 6.3. *If $g = 2$ and $x \in M$ is a primitive element of norm $q_M(x) = H_M(x, x)$, then we obtain a $q_M(x)$ -similitude $Rx \oplus M_2 \rightarrow M$ where M_2 is a rank 1 module.*

6.2. A higher dimensional Clapotis algorithm. Our algorithm can be seen as a generalisation of Clapotis to higher dimension. In Clapotis we had a rank 1 module Hermitian module, and we needed to sample two elements x_1, x_2 of coprime norm $q_M(x) = H_M(x, x)$. Here we explain how a similar strategy as Clapotis applies in higher dimension, provided that we can sample two distinct orthogonal basis of coprime “total degree”.

Definition 6.4. Let (M, H_M) be a unimodular Hermitian module of rank g . An orthogonal full submodule of M is a submodule $M' \hookrightarrow M$, given by an orthogonal basis $M' = \bigoplus_{i=1}^g R e_i$ such that $H_M(e_i, e_j) = 0$ for $i \neq j$.

Notice that for any orthogonal full submodule $\iota : M' \hookrightarrow M$, we have that $\iota^* H_M = q_M(e_1)H_0 \oplus \cdots \oplus q_M(e_g)H_0$, so $M'^\sharp \simeq \sum_i \frac{1}{q_M(e_i)} R$, and in particular $\text{Card}(M'^\sharp/M') = \prod q_M(e_i)^{r_z}$.

For technical reasons precised below Theorem 6.5, we introduce the notion of *pseudo double path*, being a double path as in Definition 5.14, but allowing the integers $\text{lcm } d_i$ and $\text{lcm } e_i$ to share a smooth cofactor inducing accessible torsion.

Theorem 6.5 (Higher dimensional Clapotis). *Let $M_1 = (e_1, \dots, e_g), M_2 = (e'_1, \dots, e'_g)$ be two orthogonal full submodules of the unimodular module (M, H_M) associated with (A, λ_A) . Let $D_1 = \text{lcm}(q_M(e_i)), D_2 = \text{lcm}(q_M(e'_i))$. Assume that the gcd, denoted d , of D_1, D_2 is smooth and the d -torsion is accessible on E_0 . Then we can efficiently build a pseudo double path $(E_0^g, \lambda_0^g) \rightarrow (A, \lambda_A)$.*

Proof. Denote $d_i := q_M(e_i)$. The inclusion $M_1 \rightarrow M$ induces, up to dualization, a $(d_1, \dots, d_g)$ -isogeny $E_0^g \rightarrow A$ on the geometric side. We now use the same trick

as in Lemma 5.18, to obtain a D_1 -isogeny $\Phi_1 : E_0^{4g} \rightarrow A^4$ and a D_2 -isogeny $\Phi_2 : E_0^{4g} \rightarrow A^4$.

Under our assumptions on the gcd of D_1, D_2 , we can compute efficiently the d -isogenies $\Phi'_1 : E_0^{4g} \rightarrow A_1$ and $\Phi'_2 : E_0^{4g} \rightarrow A_2$ factoring $\Phi_1 = \Phi'_1 \circ \Phi''_1, \Phi_2 = \Phi'_2 \circ \Phi''_2$. Then $\Phi = \widetilde{\Phi}_2 \circ \Phi_1$ factors as $E_0^{4g} \rightarrow A_1 \rightarrow A^4 \rightarrow A_2 \rightarrow E_0^{4g}$. We can compute Φ which is given by explicit endomorphisms of E_0 , along with Φ'_1, Φ'_2 , so we can find the action of the middle term $\widetilde{\Phi}_2'' \circ \Phi''_1$ on enough torsion points. We can then apply the splitting Lemma 5.15 to efficiently recover Φ''_1 and Φ''_2 , since their polarized degrees are coprime. By composition we deduce an explicit representation of the $\Phi_i = \Phi''_i \circ \Phi'_i : E_0^{4g} \rightarrow A^4$ (the splitting lemma does not apply directly to $\Phi = \widetilde{\Phi}_2 \circ \Phi_1$, as D_1 and D_2 may share a smooth cofactor). This construction is efficient, since the splitting Lemma is proven to run in polynomial time in $\log(p)$.

We conclude by using the same techniques as in Remark 5.17, to recover in polynomial time the pseudo double paths $E_0^g \rightarrow A$ given by the $(q_M(e_1), \dots, (q_M(e_g)))$ and $(q_M(e'_1), \dots, (q_M(e'_g)))$ -isogenies. $\square$

The reason we allow D_1, D_2 to not be coprime is that there exists positive definite even unimodular lattices arising from quadratic forms in (at least) 8 variables over $\mathbb{Z}$, see [CH87]. It means that we might end up with a quadratic form q_M such that $2 \mid q_M(x)$ for every $x \in M$, and thus constructing two coprime orthogonal full submodules of M might be impossible.

The above discussion shows that we boiled down the problem to computing two orthogonal full submodules $M_1, M_2 \hookrightarrow M$, inducing respectively a D_1 and D_2 -similitudes, where $D_i = 2^{u_i} D'_i$ for u_i polynomial in g and D'_i coprime odd integers. Therefore the following proposition will allow us to conclude. We refer to Section F for its proof, which makes crucial use of Proposition 6.1.

Proposition 6.6. *Let (M, H_M) be a unimodular Hermitian module, and D' be an odd integer bounded by a polynomial in p . We can efficiently compute an orthogonal full submodule $M_1 \hookrightarrow M$ inducing a D_1 -similitude $M^4 \rightarrow \mathcal{O}_0^{4g}$, where the $2^{v_2(D_1)}$ -torsion is accessible on E_0 and D_1 is coprime with D' .*

Remark 6.7. Note that technically, since the pseudo double paths constructed in Theorem 6.5 are not of coprime degrees, we cannot apply Proposition 5.19 directly. However, by the proof of Proposition 6.6, their gcd is of the form 2^n with 2^n bounded by a constant depending only on g (and not p), hence is in particular power-smooth. So we can adapt the proof of Proposition 5.16 as follows: we reduce to the case where we have a morphism $\varphi : A_1 \rightarrow B_1$, and 2^n -isogeny $u : A_1 \rightarrow E_0^r, v : B_1 \rightarrow E_0^r$, and an efficient representation of the morphism $\psi = v \circ \varphi \circ \tilde{u}$. We want to obtain an efficient representation of φ . We first remark that the proof of Proposition 5.16 works whenever we want to evaluate φ on a torsion point $P \in A_1$ of odd order N . This is already enough to deduce an efficient representation of φ whenever it is a d -isogeny, thanks to the HD representation. For the general case of a morphism, we need to “invert” u, v . We may assume that P is of order 2^m over a finite field $\mathbb{F}_q$. We first compute a basis of the 2^{m+2n} torsion on E_0^r , this can be done in polynomial time in $\log q$ because E_0 is maximal supersingular and $2^n = O(1)$. Using pairings and DLPs on this basis, we can find a point $P_1 \in E_0^r$ such that $\tilde{u}(2^n P_1) = P$. Then $\tilde{v} \circ \psi(P_1) = 2^n \varphi \circ \tilde{u}(P_1) = \varphi(P)$. (The same method would work for any pseudo double path where the gcd of the degrees is power-smooth.)

6.3. Applications to maximal supersingular abelian varieties. The construction provided by Proposition 6.6 enables us to leverage the full framework developed in Sections 6.2 and 5.3 to recover complete representations from the module-theoretic perspective. We begin with an illustrative example before stating the general result, which follows from the same line of argument.

Example 6.8. Following Example 5.20, by application of Proposition 6.6, Remark 6.7 and Theorem 6.5, the Hermitian module representation (M, H_M) of a ppav (A, λ_A) allows to recover a fully complete representation of $\text{End}(A)$.

Theorem 6.9. *Given Hermitian modules $(M_1, H_1), (M_2, H_2)$ we can construct in polynomial time the associated abelian varieties $(A_1, \lambda_1), (A_2, \lambda_2)$, and make explicit the isomorphism $\text{Hom}_{\mathbb{F}_{p^2}}(A_1, A_2) \simeq \text{Hom}_{\mathcal{O}_0}(M_2, M_1)$.*

Let us emphasize two additional corollaries to illustrate the versatility of Theorem 6.9. Corollary 6.10 is obtained by applying the theorem to the cases $(M_2, H_2) = (\mathcal{O}_0, H_0)$ and $(M_2, H_2) = (M, H_M)$. On the other hand, Corollary 6.11 follows from Theorem 6.9 applied to $(M_2, H_2) = (\mathcal{O}_0^g, H_0^g)$, together with the fact that our solution to the PIP enables the construction of an isomorphism in $\text{Hom}_{\mathcal{O}_0}(M, \mathcal{O}_0^g)$.

Corollary 6.10. *Given an Hermitian module representation (M, H_M) we can construct in polynomial time the associated abelian variety (A, λ_A) , i.e., compute the module action $M \cdot E_0$. Furthermore we can also evaluate efficiently the module action M on A along with the endomorphism action $\text{End}(A) = \text{End}_{\mathcal{O}_0}(M)$ on A .*

Corollary 6.11. *Given an Hermitian module representation (M, H_M) of a maximal ppav (A, λ_A) of dimension $g \geq 2$. We can compute, efficiently under GRH, an unpolarized isomorphism $A \simeq E_0^g$ (in both directions).*

The problem of computing isomorphisms between superspecial abelian varieties has so far only been partially addressed in the literature [Mon26; GSS25]. In [GSS25], the authors present an algorithm for computing isomorphisms between *given products* of elliptic curves, assuming that the corresponding endomorphism rings are known. In contrast, [Mon26] treats the case of arbitrary abelian *surfaces*, showing how to compute isomorphisms such as $A \rightarrow E_0^2$, using the IKO framework.

7. IMPROVEMENT TO KLPT2

The goal of this section is, given a rank 2 unimodular Hermitian module (M, H_M) , to deduce a smooth N -similitude $\mathcal{O}_0^2 \rightarrow M$. If the N -torsion on E_0 is accessible, this allows to compute the abelian surface A associated with M explicitly. This complements Section 6 where we instead compute A via a suitable double path provided by our higher dimensional Clapotis algorithm. This question was already solved in [CDK+25] (under some heuristics), using the IKO representation rather than the module representation. In this section, we explain how the module point of view, in particular Proposition 6.1 allows to improve on the bounds of [CDK+25], under the same Heuristics. For convenience, we assume that N is of the form ℓ^e . Our goal is as follows:

Problem 7.1. *Given (A_1, λ_1) and (A_2, λ_2) two principally polarized abelian surfaces, along with their associated module representation, compute an ℓ^e -isogeny $\varphi: (A_1, \lambda_1) \rightarrow (A_2, \lambda_2)$, for ℓ a small prime.*

By the anti-equivalence of categories, this reduces to:

Problem 7.2. Given (M_1, H_1) and (M_2, H_2) two u.p.d Hermitian modules of rank 2, compute an ℓ^e -similitude $\psi: (M_2, H_2) \rightarrow (M_1, H_1)$, for ℓ a small prime.

The idea is to find short smooth similitudes $\psi_i: (M_i, H_i) \rightarrow (I_i \oplus J_i, H_{i1} \oplus H_{i2})$ to direct sums of rank 1 projective $\mathcal{O}_0$ -modules I_i, J_i , and then to connect the direct sums using the 1-dimensional theory. This would lead to a composition of the form

$$(M_2, H_2) \xrightarrow{\psi_2} (I_2 \oplus J_2, H_{21} \oplus H_{22}) \xrightarrow{\text{1-dimensional computations}} (I_1 \oplus J_1, H_{11} \oplus H_{12}) \xrightarrow{\psi_1^\vee} (M_1, H_1) .$$

The 1-dimensional computations will be complete thanks to a call to KLPT [Wes21]. Therefore we mainly focus on the special case where we want to compute an ℓ^e -similitude $\psi: I_1 \oplus I_2 \hookrightarrow M$ splitting the module M .

7.1. A smooth splitting similitude. Let us present a heuristic algorithm to compute a smooth $\psi: I_1 \oplus I_2 \hookrightarrow M$. This will be done in two steps. The idea is to find an element $v \in M$ such that $H_M(v, v) = q_M(v)$ is the power of a small prime ℓ , in order to apply Corollary 6.3. For this, we first sample a primitive small element $x \in M$ of norm in $O(\sqrt{p})$. Then, we restrict to a particular submodule $\mathcal{O}_0 x + \mathcal{O}_0 y \subset M$ to find v . In both steps we will frequently use the tools developed in the KLPT² paper.

Without any loss of generality, thanks to our solution of the PIP, we assume that our M is given by a basis: $M = \mathcal{O}_0 b_1 + \mathcal{O}_0 b_2$, with given $b_1, b_2 \in M$. Then H_M is described by its Gram matrix $(H_M(b_i, b_j))_{i,j \in \{1,2\}}$.

Step 1: Find a good $x \in M$.

For efficiency reasons we first compute an element $x \in M$ of norm $q_M(x) \approx \sqrt{p}$. Writing $x = \nu_1 b_1 + \nu_2 b_2$, direct computations lead to the following equation: $q_M(x) = q_M(b_1) \text{Nrd}(\nu_1) + q_M(b_2) \text{Nrd}(\nu_2) + \text{Trd}(\nu_1 H_M(b_1, b_2) \bar{\nu}_2)$. Knowing that (M, H_M) is a unimodular Hermitian module, since it represents a principally polarized abelian, its Gram matrix has determinant $\det(\mathfrak{m}) = q_M(b_1)q_M(b_2) - \text{Nrd}(H_M(b_1, b_2)) = 1$. So, as in [CDK+25, Proposition 3.10], we can apply the usual Minkowski bound to argue that, using the Hermite–Korkine–Zolotarev lattice reduction algorithm we find an $x \in M$ such that $q_M(x) < \frac{3}{2}\sqrt{p}$. This step is polynomial in $\log(p)$ (the logarithm of the discriminant of the lattice), and theoretically efficient since we work at fixed dimension. Here we can suppose that x is primitive, otherwise we can find a smaller element in $x\mathbb{Q} \cap M$.

Step 2: Reduce to a particular submodule.

First of all, since x is primitive then as in the proof of Proposition 6.1, we can decompose $M \simeq \mathcal{O}_0 x \oplus M'$. Since M' is of rank 1, it is isomorphic to an $\mathcal{O}_0$ -ideal, so we can find by lattice reduction [DLRW24, Lemma 12] a vector $\tilde{y} \in M'$ such that the inclusion $\mathcal{O}_0 \tilde{y} \hookrightarrow M'$ is of index $q_M(\tilde{y}) = \alpha \approx \sqrt{p}$. Then if we denote by $y \in M$ any lift of $\tilde{y}$ we obtain a similitude $\psi_y: \mathcal{O}_0 x \oplus \mathcal{O}_0 \tilde{y} \hookrightarrow M$ of index α , given by $\psi_y(\mu_1 x, \mu_1 \tilde{y}) = \mu_1 x + \mu_2 y$. Now to recover $v \in M$ such that $q_M(v) = \ell^e$ for a small ℓ , it suffices to find $\mu_1, \mu_2 \in \mathcal{O}_0$ such that $q_M(\psi_y(\mu_1 x, \mu_2 \tilde{y})) = \ell^e$. Hence it suffices to solve the below equation in $\mu_1, \mu_2 \in \mathcal{O}_0$ (compare with [CDK+25, Equation (10)]):

$$(7.1) \quad \ell^e = q_M(x) \text{Nrd}(\mu_1) + q_M(y) \text{Nrd}(\mu_2) + \text{Trd}(\mu_1 H_M(x, y) \bar{\mu}_2).$$

Step 3: Find v of smooth norm.

Now we want to find $v \in \mathcal{O}_0 x + \mathcal{O}_0 y$, such that $q_M(v) = \ell^e$ for a small prime ℓ . As before we represent the Hermitian submodule $(\mathcal{O}_0 x + \mathcal{O}_0 y, H_M)$ by its Gram matrix $\mathbf{m}_y$. Note that the congruence class $\{u^* \mathbf{m}_y u\}$ does not depend on the chosen lift of $\tilde{y}$. In particular we see that the determinant of $\mathbf{m}_y$ and its top left entry are independant of the chosen lift y . Now by [CDK+25, Lemma 2.10 and Corollary 2.11] we have: $\alpha = \deg(\psi_y) = \det(\mathbf{m}_y)^2$. Using [CDK+25, Section 3.3] we can reduce the matrix $\mathbf{m}_y$ to another matrix $g = \begin{pmatrix} s & r \\ \bar{r} & t \end{pmatrix}$, such that $\text{Nrd}(r) \leq s^2(p+5)/8$. In addition reducing $\mathbf{m}_y$ preserves the determinant and the top left entry of the matrix, so $\sqrt{\alpha} = \det(g) = st - \text{Nrd}(r) = q_M(x)t - \text{Nrd}(r)$. All that leads to the estimate:

$$t \leq q_M(x)(p+5)/8 + \sqrt{\alpha}/q_M(x) \approx q_M(x)p \approx p^{3/2}.$$

Now following the proof of [CDK+25, Theorem 3.13] we can heuristically find μ_1 and μ_2 such that $q_M(\psi_y(x\mu_1, \tilde{y}\mu_2)) = \ell^e$, if $\ell^e - 2tp \text{Nrd}(r) q_M(x)^2 > 0$. Combined with the previous bounds, we obtain the approximate sufficient condition: $\ell^e > p^{11/2}$.

Consequently we can heuristically compute, in polylogarithmic time, a lift $y \in M$ of $\tilde{y}$ and $\mu_1, \mu_2 \in \mathcal{O}_0$ such that $q_M(x\mu_1 + y\mu_2) = \ell^e > p^{11/2}$. Moreover, by denoting $v := x\mu_1 + y\mu_2$, we found an element in $v \in M$ such that $q_M(v) = O(p^{11/2})$. By Corollary 6.3, the addition map induces an ℓ^e -similitude $\mathcal{O}_0 v \oplus \mathcal{O}_0 v^\perp \hookrightarrow M$ between unimodular Hermitian modules. We summarize the above method in Algorithm 5.

Algorithm 5: SPLITTINGMODULE

Input: M an left f.p.t.f $\mathcal{O}_0$ -module of rank 2, and H_M an u.p.d Hermitian form over M given as its Gram matrix

Output: A primitive left f.p.t.f $\mathcal{O}_0$ -module of rank 1 $M_1 \subset M$, such that the addition map $M_1 \oplus M_1^\perp \hookrightarrow M$ induces an ℓ^e -similitude, for ℓ a small prime.

- 1 Use a lattice reduction algorithm to compute $x \in M$ such that $q_M(x) \approx \sqrt{p}$;
 - 2 Compute $\tilde{y} \in M' = M/\mathcal{O}_0 x$ such that $\text{Card}(M'/\mathcal{O}_0 \tilde{y}) \approx \sqrt{p}$;
 - 3 Compute a lift $y \in M$ of $\tilde{y}$, $\mu_1, \mu_2 \in \mathcal{O}_0$ such that $q_M(\mu_1 x + \mu_2 y) = \ell^e \approx p^{11/2}$;
 - 4 Return $\mathcal{O}_0 v$, where $v = \mu_1 x + \mu_2 y$.
-

7.2. KLPT2 for the module representation. We can now solve Problem 7.2, using the same notations and the subscript $i \in \{1, 2\}$. Previous paragraph allows (heuristically) for the computation of two vectors $v_i \in M_i$ and ℓ^{e_i} -similitudes $\mathcal{O}_0 v_i \oplus M_i' \hookrightarrow M_i$, with $\ell^{e_i} \approx p^{11/2}$. Thus by Example 2.8 we recover the corresponding ℓ^{e_i} -isogenies $\varphi_i: A_i \rightarrow E_0 \times E_i$. Now the 1-dimensional KLPT algorithm allows to compute a smooth ℓ^{e_3} -isogeny $\varphi_3: E_1 \rightarrow E_2$, for $\ell^{e_3} = O(p^{4.5})$, see [DKLPW20, Section 5.2] for the estimation of ℓ^{e_3} . Since $p^{4.5} \gg p$, we can compute an endomorphism $\varphi_3': E_0 \rightarrow E_0$ of degree ℓ^{e_3} by solving a norm equation in $\mathcal{O}_0$. Finally, by denoting $\Phi: E_0 \times E_1 \rightarrow E_0 \times E_2$ the diagonal isogeny (φ_3', φ_3) , the composition $\widehat{\varphi}_2 \circ \Phi \circ \varphi_1: A_1 \rightarrow A_2$ gives an $\ell^{e_1+e_2+e_3}$ -isogeny for $\ell^{e_1+e_2+e_3} = O(p^{31/2})$.

Theorem 7.3. *Algorithm 6 is correct. Under the heuristics from [CDK+25, Theorem 3.13], it outputs an ℓ^e -isogeny for $\ell^e = O(p^{31/2})$, in polynomial-time.*

Algorithm 6: MODULEKLPT²

Input: Two Hermitian unimodular f.p.t.f $\mathcal{O}_0$ -modules of rank 2, (M_i, H_i) .

Output: An ℓ -isogeny $A_1 \rightarrow A_2$, where A_i is the abelian variety associated with M_i .

- 1 For $i \in \{1, 2\}$ compute a smooth ℓ -similitude of the form $\mathcal{O}_0 v_i \oplus \mathcal{O}_0 v_i^\perp \rightarrow M_i$ given by Algorithm 5 SPLITTINGMODULE ;
 - 2 Recover the two corresponding ℓ -isogeny $\varphi_i: A_i \rightarrow E_0 \times E_i$;
 - 3 Compute the dual isogeny $\widehat{\varphi}_2: E_0 \times E_2 \rightarrow A_2$;
 - 4 Use 1-dimensional KLPT to compute an ℓ -isogeny $\varphi_3: E_1 \rightarrow E_2$ and an endomorphism $\varphi'_3: E_0 \rightarrow E_0$ of the same degree;
 - 5 Return the composition $\widehat{\varphi}_2 \circ (\varphi'_3, \varphi_3) \circ \varphi_1: A_1 \rightarrow A_2$;
-

Proof. The correctness follows from the above discussion. Using the same notation, since we used arguments of [CDK+25, Section 3] to compute φ_1 and φ_2 in polynomial time and with the estimated size, our result inherit their heuristics (for Step 3 of Algorithm 5). It is clear that the other operations are at most polynomial. $\square$

Remark 7.4. If $g > 2$, we can as above sample a primitive $x \in M$, and invoke Proposition 6.1 to reduce to a submodule $M_1 \oplus_\perp M_2$ where $M_1 = \mathcal{O}_0 x$. The problem is that now M_2 is of rank 2, and $M_2 \mid H_M$ may not be a multiple of a principal polarization on M_2 . This is the key difference with the case $g = 2$ where M_2 is of rank 1 and every polarization is a multiple of the canonical principal one. Thus it is not obvious how to extend KLPT² to $g > 2$ by recursive calls to smaller dimension, unless we extend it to also treat non principally polarized abelian varieties.

REFERENCES

- [AAA+25] M. A. Aardal, G. Adj, D. F. Aranha, A. Basso, I. A. C. Martínez, J. Chávez-Saab, M. C.-R. Santos, P. Dartois, L. D. Feo, M. Duparc, J. K. Eriksen, T. B. Fouotsa, D. L. G. Filho, B. Hess, D. Kohel, A. Leroux, P. Longa, L. Maino, M. Meyer, K. Nakagawa, H. Onuki, L. Panny, S. Patranabis, C. Petit, G. Pope, K. Reijnders, D. Robert, F. R. Henríquez, S. Schaeffler, and B. Wesolowski. *SQIsign - Version 2.0. Technical report*. Available at <https://csrc.nist.gov/Projects/pqc-dig-sig/round-2-additional-signatures>. 2025 (cit. on p. 1).
- [ABIGW25] S. Anni, G. Bisson, A. Iezzi, E. L. García, and B. Wesolowski. “On the computation of endomorphism rings of abelian surfaces over finite fields”. In: *arXiv preprint arXiv:2503.08925* (2025). DOI: [10.48550/arXiv.2503.08925](https://doi.org/10.48550/arXiv.2503.08925) (cit. on p. 4).
- [BDF+24] A. Basso, P. Dartois, L. D. Feo, A. Leroux, L. Maino, G. Pope, D. Robert, and B. Wesolowski. “SQIsign2D-West The Fast, the Small, and the Safer”. In: *Lecture Notes in Computer Science*. Vol. 15486. Lecture Notes in Computer Science. Kolkata, India: ASIACRYPT, Dec. 2024, pp. 339–370. DOI: [10.1007/978-981-96-0891-1_11](https://doi.org/10.1007/978-981-96-0891-1_11). URL: <https://hal.science/hal-04603556> (cit. on pp. 1, 10).

- [BL04] C. Birkenhake and H. Lange. *Complex Abelian Varieties*. Springer Berlin, Heidelberg, 2004. DOI: [10.1007/978-3-662-06307-1](https://doi.org/10.1007/978-3-662-06307-1) (cit. on p. 30).
- [BHJ22] W. Bley, T. Hofmann, and H. Johnston. “Computation of lattice isomorphisms and the integral matrix similarity problem”. English. In: *Forum Math. Sigma* 10 (2022). Id/No e87, p. 36. ISSN: 2050-5094. DOI: [10.1017/fms.2022.74](https://doi.org/10.1017/fms.2022.74) (cit. on p. 18).
- [BPTW25] K. de Boer, A. Page, R. Toma, and B. Wesolowski. *Average hardness of SIVP for module lattices of fixed rank*. Preprint, arXiv:2511.13659. 2025. URL: <https://arxiv.org/abs/2511.13659> (cit. on pp. 15, 16).
- [BCE+25] G. Borin, M. Corte-Real Santos, J. K. Eriksen, R. Invernizzi, M. Mula, S. Schaeffler, and F. Vercauteren. “Qlapoti: Simple and efficient translation of quaternion ideals to isogenies”. In: *International Conference on the Theory and Application of Cryptology and Information Security*. Springer. 2025, pp. 174–205. DOI: [10.1007/978-981-95-5113-2_6](https://doi.org/10.1007/978-981-95-5113-2_6) (cit. on p. 1).
- [CD23] W. Castryck and T. Decru. “An efficient key recovery attack on SIDH”. In: *Annual International Conference on the Theory and Applications of Cryptographic Techniques*. Springer. 2023, pp. 423–447. DOI: [10.1007/978-3-031-30589-4_15](https://doi.org/10.1007/978-3-031-30589-4_15) (cit. on p. 1).
- [CDK+25] W. Castryck, T. Decru, P. Kutas, A. Laval, C. Petit, and Y. B. Ti. “KLPT²: Algebraic Pathfinding in Dimension Two and Applications”. In: *Advances in Cryptology – CRYPTO 2025*. Ed. by Y. Tauman Kalai and S. F. Kamara. Cham: Springer Nature Switzerland, 2025, pp. 167–200. ISBN: 978-3-032-01855-7. DOI: [10.1007/978-3-032-01855-7_6](https://doi.org/10.1007/978-3-032-01855-7_6) (cit. on pp. 2, 3, 5, 7, 8, 14, 30, 33–36).
- [CDS20] W. Castryck, T. Decru, and B. Smith. “Hash functions from superspecial genus-2 curves using Richelot isogenies”. In: *Journal of Mathematical Cryptology* 14.1 (2020), pp. 268–292. DOI: [10.1515/jmc-2019-0021](https://doi.org/10.1515/jmc-2019-0021) (cit. on p. 1).
- [CEIV26] W. Castryck, J. K. Eriksen, R. Invernizzi, and F. Vercauteren. “The Principal Ideal Problem for endomorphism rings of superspecial abelian varieties”. 2026. URL: <https://ia.cr/2026/454> (cit. on pp. 5, 9, 10, 17, 18, 22).
- [CLG09] D. Charles, K. Lauter, and E. Goren. “Cryptographic hash functions from expander graphs”. In: *Journal of Cryptology* 22.1 (2009), pp. 93–113. ISSN: 0933-2790. DOI: [10.1007/s00145-007-9002-x](https://doi.org/10.1007/s00145-007-9002-x) (cit. on p. 1).
- [CME+25] C. Cheviguard, G. Mureau, T. Espitau, A. Pellet-Mary, H. Pliatsok, and A. Wallet. “A reduction from Hawk to the principal ideal problem in a quaternion algebra”. English. In: *EUROCRYPT 2025*. Cham: Springer, 2025, pp. 154–183. ISBN: 978-3-031-91123-1; 978-3-031-91124-8. DOI: [10.1007/978-3-031-91124-8_6](https://doi.org/10.1007/978-3-031-91124-8_6) (cit. on p. 18).
- [Coh93] H. Cohen. *A course in computational algebraic number theory*. English. Vol. 138. Grad. Texts Math. Berlin: Springer-Verlag, 1993. ISBN: 3-540-55640-0 (cit. on p. 18).

- [Coh12] H. Cohen. *Advanced topics in computational number theory*. Vol. 193. Springer Science & Business Media, 2012. DOI: [10.1007/978-1-4419-8489-0](#) (cit. on p. 45).
- [CH87] P. J. Costello and J. S. Hsia. “Even unimodular 12-dimensional quadratic forms over $\mathbb{Q}(\sqrt{5})$ ”. In: *Advances in Mathematics* 64.3 (1987), pp. 241–278. ISSN: 0001-8708. DOI: [10.1016/0001-8708\(87\)90009-0](#) (cit. on p. 32).
- [DEF+25] P. Dartois, J. K. Eriksen, T. B. Fouotsa, A. Herlédan Le Merdy, R. Invernizzi, D. Robert, R. Rueger, F. Vercauteren, and B. Wesolowski. “PEGASIS: Practical Effective Class Group Action using 4-Dimensional Isogenies”. In: *Advances in Cryptology – CRYPTO 2025*. Ed. by Y. Tauman Kalai and S. F. Kamara. Cham: Springer Nature Switzerland, 2025, pp. 67–99. ISBN: 978-3-032-01855-7. DOI: [10.1007/978-3-032-01855-7_3](#) (cit. on pp. 1, 3).
- [DEIV26] P. Dartois, J. K. Eriksen, R. Invernizzi, and F. Vercauteren. “qt-Pegasis: simpler and faster effective class group actions”. In: *Annual International Conference on the Theory and Applications of Cryptographic Techniques*. Springer, 2026, pp. 636–665. DOI: [10.1007/978-3-032-25327-9_22](#) (cit. on pp. 1, 3).
- [DLRW24] P. Dartois, A. Leroux, D. Robert, and B. Wesolowski. “SQISignHD: New Dimensions in Cryptography”. In: *Advances in Cryptology – EUROCRYPT 2024*. Ed. by M. Joye and G. Leander. Vol. 14651. Lecture Notes in Computer Science. Springer, Heidelberg, 2024, pp. 3–32. ISBN: 978-3-031-58716-0. DOI: [10.1007/978-3-031-58716-0_1](#) (cit. on pp. 1, 27, 34).
- [DKLPW20] L. De Feo, D. Kohel, A. Leroux, C. Petit, and B. Wesolowski. “SQISign: Compact Post-quantum Signatures from Quaternions and Isogenies”. English. In: *Advances in Cryptology – ASIACRYPT 2020*. Ed. by S. Moriai and H. Wang. Lecture Notes in Computer Science. Springer, 2020, pp. 64–93. ISBN: 9783030648367. DOI: [10.1007/978-3-030-64837-4_3](#) (cit. on pp. 1, 35).
- [DD08] L. Dembélé and S. Donnelly. “Computing Hilbert modular forms over fields with nontrivial class group”. English. In: *Algorithmic number theory. 8th international symposium, ANTS-VIII Banff, Canada, May 17–22, 2008 Proceedings*. Berlin: Springer, 2008, pp. 371–386. ISBN: 978-3-540-79455-4. DOI: [10.1007/978-3-540-79456-1_25](#) (cit. on p. 18).
- [EHLMP18] K. Eisenträger, S. Hallgren, K. Lauter, T. Morrison, and C. Petit. “Supersingular Isogeny Graphs and Endomorphism Rings: Reductions and Solutions”. In: *Advances in Cryptology – EUROCRYPT 2018*. Ed. by J. B. Nielsen and V. Rijmen. Cham: Springer International Publishing, 2018, pp. 329–368. ISBN: 978-3-319-78372-7. DOI: [10.1007/978-3-319-78372-7_11](#) (cit. on pp. 1, 3, 4).
- [EPSV24] J. Eriksen, L. Panny, J. Sotáková, and M. Veroni. “Deuring for the people: Supersingular elliptic curves with prescribed endomorphism ring in general characteristic”. In: *Contemporary Mathematics* 796 (2024), pp. 339–373. DOI: [10.1090/conm/796/16008](#) (cit. on pp. 10, 43).

- [FP85] U. Fincke and M. Pohst. “Improved methods for calculating vectors of short length in a lattice, including a complexity analysis”. English. In: *Math. Comput.* 44 (1985), pp. 463–471. ISSN: 0025-5718. DOI: [10.2307/2007966](https://doi.org/10.2307/2007966) (cit. on p. 16).
- [FMP23] T. B. Fouotsa, T. Moriya, and C. Petit. “M-SIDH and MD-SIDH: Countering SIDH Attacks by Masking Information”. In: *Advances in Cryptology – EUROCRYPT 2023*. Ed. by C. Hazay and M. Stam. Cham: Springer Nature Switzerland, 2023, pp. 282–309. ISBN: 978-3-031-30589-4. DOI: [10.1007/978-3-031-30589-4_10](https://doi.org/10.1007/978-3-031-30589-4_10) (cit. on p. 10).
- [GSS25] P. Gaudry, J. Soumier, and P.-J. Spaenlehauer. “Computing Isomorphisms between Products of Supersingular Elliptic Curves”. In: *arXiv preprint arXiv:2503.21535* (2025) (cit. on pp. 6, 8, 15, 33).
- [Gra84] D. R. Grayson. “Reduction theory using semistability”. English. In: *Comment. Math. Helv.* 59 (1984), pp. 600–634. ISSN: 0010-2571. DOI: [10.1007/BF02566369](https://doi.org/10.1007/BF02566369). URL: <https://eudml.org/doc/139993> (cit. on p. 15).
- [Gra86] D. R. Grayson. “Reduction theory using semistability. II”. English. In: *Comment. Math. Helv.* 61 (1986), pp. 661–676. ISSN: 0010-2571. DOI: [10.1007/BF02621937](https://doi.org/10.1007/BF02621937). URL: <https://eudml.org/doc/140073> (cit. on p. 16).
- [HS07] G. Hanrot and D. Stehlé. “Improved analysis of Kannan’s shortest lattice vector algorithm”. English. In: *Advances in cryptology – CRYPTO 2007. 27th annual international cryptology conference, Santa Barbara, CA, USA, August 19–23, 2007. Proceedings*. Berlin: Springer, 2007, pp. 170–186. ISBN: 978-3-540-74142-8. DOI: [10.1007/978-3-540-74143-5_10](https://doi.org/10.1007/978-3-540-74143-5_10) (cit. on p. 16).
- [HW25] A. Herlédan Le Merdy and B. Wesolowski. “Unconditional Foundations for Supersingular Isogeny-Based Cryptography”. In: *Theory of Cryptography Conference*. Springer. 2025, pp. 266–297. DOI: [10.1007/978-3-032-12296-4_9](https://doi.org/10.1007/978-3-032-12296-4_9) (cit. on pp. 1, 3, 10).
- [IR93] G. Ivanyos and L. Rónyai. “Finding maximal orders in semisimple algebras over $\mathbb{Q}$ ”. In: *computational complexity* 3 (1993), pp. 245–261. URL: <https://api.semanticscholar.org/CorpusID:260640> (cit. on p. 17).
- [IRS12] G. Ivanyos, L. Rónyai, and J. Schicho. “Splitting full matrix algebras over algebraic number fields”. In: *Journal of Algebra* 354.1 (2012), pp. 211–223. DOI: [10.1016/j.jalgebra.2012.01.008](https://doi.org/10.1016/j.jalgebra.2012.01.008) (cit. on pp. 4, 15–17).
- [JKP+18] B. W. Jordan, A. G. Keeton, B. Poonen, E. M. Rains, N. Shepherd-Barron, and J. T. Tate. “Abelian varieties isogenous to a power of an elliptic curve”. In: *Compositio Mathematica* 154.5 (2018), pp. 934–959. DOI: [10.1112/S0010437X17007990](https://doi.org/10.1112/S0010437X17007990) (cit. on pp. 2, 6, 10–12).
- [Kir16] M. Kirschmer. “Definite quadratic and hermitian forms with small class number”. In: *Habilitation, RWTH Aachen University* (2016). URL: <https://www.math.rwth-aachen.de/~Markus.Kirschmer/papers/herm.pdf> (cit. on p. 15).

- [KNRR21] M. Kirschmer, F. Narbonne, C. Ritzenthaler, and D. Robert. *Spanning the isogeny class of a power of an ordinary elliptic curve over a finite field. Application to the number of rational points of curves of genus ≤ 4* . 2021. arXiv: [2004.08315 \[math.NT\]](https://arxiv.org/abs/2004.08315). URL: <https://arxiv.org/abs/2004.08315> (cit. on p. 45).
- [KV10] M. Kirschmer and J. Voight. “Algorithmic enumeration of ideal classes for quaternion orders”. English. In: *SIAM J. Comput.* 39.5 (2010), pp. 1714–1747. ISSN: 0097-5397. DOI: [10.1137/080734467](https://doi.org/10.1137/080734467) (cit. on p. 18).
- [KLPT14] D. Kohel, K. Lauter, C. Petit, and J.-P. Tignol. “On the quaternion-isogeny path problem”. In: *LMS Journal of Computation and Mathematics* 17.A (2014), pp. 418–432. DOI: [10.1112/S1461157014000151](https://doi.org/10.1112/S1461157014000151) (cit. on pp. 1, 2, 9, 18).
- [KMM+25] S. Kunzweiler, L. Maino, T. Moriya, C. Petit, G. Pope, D. Robert, M. Stopar, and Y. B. Ti. “Radical 2-isogenies and cryptographic hash functions in dimensions 1, 2 and 3”. In: *Public-Key Cryptography – PKC 2025*. Vol. 15676, Lecture Notes in Computer Science. Springer, 2025, pp. 265–299. DOI: [10.1007/978-3-031-91826-1_9](https://doi.org/10.1007/978-3-031-91826-1_9) (cit. on p. 1).
- [Ler25] A. Leroux. “Verifiable Random Function from the Deuring Correspondence and Higher Dimensional Isogenies”. In: *Advances in Cryptology - EUROCRYPT 2025*. 2025. DOI: [10.1007/978-3-031-91098-2_7](https://doi.org/10.1007/978-3-031-91098-2_7) (cit. on pp. 1, 10).
- [LR22] D. Lubicz and D. Robert. “Fast change of level and applications to isogenies”. In: *Research in Number Theory (ANTS XV Conference)* 9.1 (2022). DOI: [10.1007/s40993-022-00407-9](https://doi.org/10.1007/s40993-022-00407-9) (cit. on p. 24).
- [MMPPW23] L. Maino, C. Martindale, L. Panny, G. Pope, and B. Wesolowski. “A direct key recovery attack on SIDH”. In: *Annual International Conference on the Theory and Applications of Cryptographic Techniques*. Springer, 2023, pp. 448–471. DOI: [10.1007/978-3-031-30589-4_16](https://doi.org/10.1007/978-3-031-30589-4_16) (cit. on p. 1).
- [Mon26] M. Montessinos. “Equivalent computational problems for superspecial abelian surfaces”. In: *arXiv preprint arXiv:2601.18050* (2026). DOI: [10.48550/arXiv.2601.18050](https://doi.org/10.48550/arXiv.2601.18050) (cit. on pp. 4–6, 9, 26, 33).
- [Mum70] D. Mumford. *Abelian Varieties*. *Tata Institute of Fundamental Research Studies in Mathematics*, no. 5. Oxford University Press, 1970 (cit. on p. 25).
- [Oda69] T. Oda. “The first de Rham cohomology group and Dieudonné modules”. In: *Annales scientifiques de l’École Normale Supérieure*. Vol. 2. 1. 1969, pp. 63–135 (cit. on p. 27).
- [Pag14] A. Page. “An algorithm for the Principal Ideal Problem in indefinite quaternion algebras”. English. In: *LMS J. Comput. Math.* 17A (2014), pp. 366–384. ISSN: 1461-1570. DOI: [10.1112/S1461157014000321](https://doi.org/10.1112/S1461157014000321) (cit. on p. 18).
- [PR23] A. Page and D. Robert. *Introducing Clapoti(s): Evaluating the isogeny class group action in polynomial time*. Cryptology ePrint Archive, Paper 2023/1766. 2023. URL: <https://eprint.iacr.org/2023/1766> (cit. on p. 1).

- [PW24] A. Page and B. Wesolowski. “The supersingular endomorphism ring and one endomorphism problems are equivalent”. In: *Advances in Cryptology (Eurocrypt 2024)*. Springer. 2024, pp. 388–417. DOI: [10.1007/978-3-031-58751-1_14](https://doi.org/10.1007/978-3-031-58751-1_14) (cit. on p. 3).
- [PPS25] L. Panny, C. Petit, and M. Stopar. “KLaPoTi: An asymptotically efficient isogeny group action from 2-dimensional isogenies”. In: *IACR Communications in Cryptology 2.3* (2025). DOI: [10.62056/ahp2wakrz](https://doi.org/10.62056/ahp2wakrz) (cit. on pp. 1, 3).
- [PT18] P. Pollack and E. Treviño. “Finding the Four Squares in Lagrange’s Theorem.” In: *Integers* 18 (2018), A15. URL: <https://api.semanticscholar.org/CorpusID:203588112> (cit. on p. 5).
- [Rei03] I. Reiner. *Maximal Orders*. Oxford University Press, 2003. ISBN: 9780198526735 (cit. on pp. 18, 43).
- [Rob21] D. Robert. “Efficient algorithms for abelian varieties and their moduli spaces”. HDR thesis. Université Bordeaux, 2021. URL: <https://www.math.u-bordeaux.fr/~damienrobert/pro/publications/academic/hdr.pdf> (cit. on p. 24).
- [Rob23] D. Robert. “Breaking SIDH in polynomial time”. In: *Annual International Conference on the Theory and Applications of Cryptographic Techniques*. Springer. 2023, pp. 472–503. DOI: [10.1007/978-3-031-30589-4_17](https://doi.org/10.1007/978-3-031-30589-4_17) (cit. on p. 1).
- [Rob24] D. Robert. *The module action for isogeny based cryptography*. Cryptology ePrint Archive, Paper 2024/1556. 2024. URL: <https://eprint.iacr.org/2024/1556> (cit. on pp. 1–4, 6, 8, 10–12, 30, 44, 45).
- [Rob25] D. Robert. “On the efficient representation of isogenies (a survey)”. In: *Number-Theoretic Methods in Cryptology – NuTMiC 2024*. Ed. by A. Dąbrowski, J. Pieprzyk, and J. Pomykała. Vol. 14966. Lecture Notes in Computer Science. Springer Nature Switzerland, 2025, pp. 3–84. DOI: https://doi.org/10.1007/978-3-031-82380-0_1 (cit. on pp. 24, 27, 29, 45).
- [SW14] U. Shapira and B. Weiss. “A volume estimate for the set of stable lattices”. English. In: *C. R., Math., Acad. Sci. Paris* 352.11 (2014), pp. 875–879. ISSN: 1631-073X. DOI: [10.1016/j.crma.2014.08.019](https://doi.org/10.1016/j.crma.2014.08.019) (cit. on p. 15).
- [Voi21] J. Voight. *Quaternion Algebras*. Springer Cham, 2021. ISBN: 978-3-030-56692-0. DOI: [10.1007/978-3-030-56694-4](https://doi.org/10.1007/978-3-030-56694-4) (cit. on pp. 11, 14, 17, 21).
- [Wes21] B. Wesolowski. “The supersingular isogeny path and endomorphism ring problems are equivalent”. In: *FOCS 2021*. IEEE, 2021, pp. 1100–1111. DOI: [10.1109/FOCS52979.2021.00109](https://doi.org/10.1109/FOCS52979.2021.00109) (cit. on pp. 1–4, 34).
- [Wes22] B. Wesolowski. “Understanding and improving the Castryck-Decru attack on SIDH”. 2022. URL: <https://hal.science/hal-04557845> (cit. on p. 4).

APPENDIX A. A SUMMARY OF THE PROVIDED CONVERSIONS

In this section we illustrate the conversions provided through our work, section by section. The dashed arrows indicate trivial translations, where the starting point

is already more structured than the ending one. We would like to emphasize that the abstract module representation allows to compute every other representation in polynomial time, potentially under GRH. We begin by the results from Section 3.




FIGURE 1. Summary of Section 3 translations

We continue with results from Section 5. Note that the explicit (hence the complete) representation of $\text{End}(A)$ inherently contains an explicit representation of the polarized abelian variety (A, λ_A) .




FIGURE 2. Summary of Section 5 translations

APPENDIX B. EXPLICIT EXAMPLE

We present an experimental result obtained from our code, available at:

<https://gitlab.inria.fr/module-representation/PIP-quaternion-mat>.

We fix the prime $p = 2^{31} - 1$. We work with SageMath version 10.8 using Python 3.12.5. In the file `TestsPrincipalize.sage` (where the seed is fixed at 12345), the ideal to principalize is given by the two following generators:

$$\begin{pmatrix} -107 + 133/2 i + 7/2 j - 21 k & -76 - 17 i + 8 k \\ 47/2 - 167/2 i - 93/2 j + 65/2 k & -73/2 + 126 i + 40 j + 33/2 k \end{pmatrix}$$

and

$$\begin{pmatrix} -107/2 + 195/2 i + 1/2 j - 11/2 k & -77/2 + 2 i + 21 j - 5/2 k \\ 135/2 + 101/2 i + 75/2 j - 25/2 k & -13 - 47/2 i - 53/2 j + 15 k \end{pmatrix}.$$

Then our algorithm output a generator of the form $\begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix}$, where $x \in \mathcal{O}_0$ is:

$$\begin{aligned} & -46730783585866828646136107206482198199245953093776405796899252978 \\ & -48825422605014945677722426096850970410908167898189669861575082117/2 \, i \\ & + 9337186162012436912667941462130059756799956950052870966124091229/2 \, j \\ & + 592123978896402418727409736127151116360250499937526932543298788 \, k. \end{aligned}$$

Note that we check the outputs using linear algebra. Finally, our algorithm relies on KLPT as a subroutine, and we used the already implemented version of KLPT from [EPSV24].

APPENDIX C. MORITA EQUIVALENCE

In this section we explain some consequences of the Morita equivalence between $\mathcal{O}_0$ and $\mathcal{M}_g(\mathcal{O}_0)$. The interested reader may refer to [Rei03] for a more comprehensive exposition.

Let M be a right $\mathcal{O}_0$ -module. We denote by Λ the ring $\text{Hom}_{\mathcal{O}_0}(M, M)$. Remark that M is canonically equipped with a left Λ -module structure. By [Rei03, Corollary 16.9], if M is a free module with $g \geq 2$ generators, then we have a *Morita equivalence* between $\mathcal{O}_0$ and Λ . Moreover, if we have a free $\mathcal{O}_0$ -basis of M , we can explicitly compute an isomorphism $\Lambda \simeq \mathcal{M}_g(\mathcal{O}_0)$ (Equation (16.17) of the same book). We denote by N the left $\mathcal{O}_0$ -module $\text{Hom}_{\mathcal{O}_0}(M, \mathcal{O}_0)$, and by $\mathcal{M}_{\mathcal{O}_0}$ (resp. $\mathcal{M}_{\Lambda}$) the category of right $\mathcal{O}_0$ -modules (resp. right Λ -modules). Keeping this notations, we highlight a consequence of the Morita equivalence.

Theorem C.1. [Rei03, Theorem 16.14]

The following functors $S: \mathcal{M}_{\Lambda} \rightarrow \mathcal{M}_{\mathcal{O}_0}$ and $T: \mathcal{M}_{\mathcal{O}_0} \rightarrow \mathcal{M}_{\Lambda}$ are inverse equivalences of categories.

$$\begin{aligned} S &= \cdot \otimes_{\Lambda} M \simeq \text{Hom}_{\Lambda}(N, \cdot), \\ T &= \cdot \otimes_{\mathcal{O}_0} N \simeq \text{Hom}_{\mathcal{O}_0}(M, \cdot). \end{aligned}$$

Moreover they induce an isomorphism of lattices:

$$\begin{array}{ccc} \varphi & : & L_1 \rightarrow L_2 \\ & & J \mapsto J \cdot M \end{array}$$

Where L_1 is the lattice of right ideals of Λ , L_2 is the lattice of right $\mathcal{O}_0$ -submodules of M , and $J \cdot M$ denotes the canonical left action of $J \subset \Lambda = \text{Hom}_{\mathcal{O}_0}(M, M)$ over elements of M .

It follows that if $M = \mathcal{O}_0^g$, then φ induces a lattice isomorphism between the right submodules of $\mathcal{O}_0^g$ and the right ideals of $\mathcal{M}_g(\mathcal{O}_0)$, which recovers Proposition 3.5, establishing equivalence between Problems 4.1 and 4.2. Let us give more details to facilitate the readability of the description of our algorithm.

If $M = m_1 \mathcal{O}_0 + \dots + m_r \mathcal{O}_0$, then M is exactly the module generated by the columns of the matrix $C = (m_1 \dots m_r) \in \mathcal{M}_{g \times r}(\mathcal{O}_0)$. Write $r = qg + r$ the Euclidean division of r by g , and let us denote by I the right ideal of $\mathcal{M}_g(\mathcal{O}_0)$ generated by

$$(m_1 \dots m_g), (m_{g+1} \dots m_{2g}), \dots, (m_{qg+1} \dots m_r \, 0 \dots 0).$$

Since we work with right modules, if α is such that $I = \alpha \mathcal{M}_g(\mathcal{O}_0)$ then α is obtained as a linear combination of the columns m_i . In other words, the module generated by the columns of α is the same than the one generated by the columns of C , which is M . Thus the columns of α give a generating set of M with g elements. Since M is free of rank g , it is a basis. Reciprocally, if I is given by generators $\alpha_1, \dots, \alpha_r$ we consider the module M generated by the columns of the α_i . Then computing a basis $v_1, \dots, v_g \in \mathcal{O}_0^g$ of M leads to a principal generator of I given by the matrix $(v_1 \cdots v_g) \in \mathcal{M}_g(\mathcal{O}_0)$.

APPENDIX D. COMPUTING EXPLICIT ISOMORPHISMS $A \simeq E_0^g$

In this section we present an application of our HD-Clapotis algorithm. We detail the case of computing an effective isomorphism $A \simeq E_0^g$, given the Hermitian module representation (M, H_M) of a polarized abelian variety (A, λ_A) . We assume that we fixed an effective representation of $\mathcal{M}_g(\mathcal{O}_0)$ acting on E_0^g . To lighten notation, we assume in the following that the double paths computed are N -isogenies.

Following the proof of Theorem 6.5, up to dualization, we can compute, for $i \in \{1, 2\}$, $\psi_i: M \rightarrow \mathcal{O}_0^g$ a d_i -similitude with d_1 and d_2 coprime integers. Our solution to the PIP leads to the computation of an isomorphism $\psi: \mathcal{O}_0^g \rightarrow M$, that does not preserves the Hermitian structures. It yields the following diagram

$$(\mathcal{O}_0^g, \psi^* H_M) \xrightarrow{\psi} (M, H_M) \begin{array}{c} \xrightarrow{\psi_1} \\ \xrightarrow{\psi_2} \end{array} (\mathcal{O}_0^g, \text{Id}) .$$

On the one hand, we can compute the polarized isogenies $\varphi_1, \varphi_2: (E_0^g, \lambda_0^g) \rightarrow (A, \lambda_A)$ corresponding to the similitudes ψ_1, ψ_2 by applying the Clapoti(s) approach from [Rob24, Proposition 4.10].

On the other hand, we can compute the matrix representations of the compositions $\psi'_i := \psi \circ \psi_i: \mathcal{O}_0^g \rightarrow \mathcal{O}_0^g$ for $i \in \{1, 2\}$. Therefore, seeing them as element of $\text{End}(E_0^g) \simeq \mathcal{M}_g(\mathcal{O}_0)$ we can evaluate them as (unpolarized) isogenies $\varphi'_i: E_0^g \rightarrow E_0^g$.

The following diagram describes the isogenies that we can evaluate, the dashed arrow representing the isomorphism $\varphi: E_0^g \rightarrow A$ we want to recover.

$$\begin{array}{ccccc} & & \varphi'_1 & & \\ & \nearrow & & \searrow & \\ E_0^g & \xrightarrow{\varphi_1} & A & \xrightarrow{\varphi} & E_0^g \\ & \searrow & & \nearrow & \\ & & \varphi'_2 & & \end{array}$$

We have the commutativity of both up and down part of this diagram: $\varphi \circ \varphi_i = \varphi'_i$. Now to evaluate φ on points of A , for simplicity we look at the action of φ on a point of order ℓ^e . Let $P \in A[\ell^e]$ for a fixed prime ℓ . Since d_1 and d_2 are coprime, there exists $i \in \{1, 2\}$ such that $\gcd(\ell, d_i) = 1$.

Without any loss of generality, we may assume that $i = 1$. Therefore to compute $\varphi(P)$, we first compute $Q = \widehat{\varphi}_1(P) \in E_0^g$. Recall that an explicit representation of φ_1 always leads to an explicit representation of $\widehat{\varphi}_1$. Then we can compute $\varphi'_1(Q) = d_1 \varphi(P) \in A$. Finally since d_1 and ℓ are coprime, we recover $\varphi(P) = d_1^{-1} \widehat{\varphi}_1(Q)$.

APPENDIX E. THE COMMUTATIVE CASE

Let R be a primitive quadratic imaginary suborder of $\text{End}(E_0)$ for some elliptic curve $E_0/\mathbb{F}_q$. We suppose that we know how to evaluate the endomorphisms in R explicitly on E_0 .

Typical examples are when E_0 is ordinary or supersingular over $\mathbb{F}_p$ and $R = \text{End}(E_0)$, or E_0 is maximal over $\mathbb{F}_{p^2}$ and R is a primitive suborder of $\mathcal{O}_0 = \text{End}(E_0)$. We remark that when $\mathbb{Z}[\pi_q]$ is a non trivial suborder of R , then the isogeny division algorithm allows to compute any endomorphism on R explicitly, see [Rob25].

We also have an (anti)-equivalence of categories $A \mapsto \text{Hom}_R(A, E_0)$ between R -oriented abelian varieties A isogeneous to E_0^g , and torsion free R -modules of rank g (under some mild extra conditions on A if p is inert in R). A principal polarization on A also corresponds to a unimodular Hermitian R -form. We refer to [Rob24] for more details.

Assume first for simplicity that R is a maximal order. Then most of the results of Section 6.3 apply directly to this case. First, a torsion free R -module M of rank g is projective, and isomorphic to $R^{g-1} \oplus I$, where I is an invertible ideal of R (called the Steinitz class of M). The isomorphism class of I uniquely determines the isomorphism class of M . Computing I from M is standard using a pseudo-basis of M , along with the computation of an explicit isomorphism $M \simeq R^{g-1} \oplus I$, see [Coh12, Chapter 1].

If R is a non maximal quadratic imaginary order, things become more tricky. Already in dimension 1 we do not know how to efficiently convert the isogeny associated with the conductor ideal $\mathfrak{f} \subset R$ (which corresponds to the ascending isogeny in the volcano) into an isogeny when $f = N(\mathfrak{f})$ contains a large prime factor. We cannot apply Clapotis directly because each element $x \in \mathfrak{f}$ has norm divisible by f . We need to add the extra assumption that M is projective over R , not only torsion free. In that case, M is also isomorphic to $R^{g-1} \oplus I$ for an invertible ideal I of R , as in the maximal case.

The Hermitian Poincaré decomposition theorem still holds, because it holds for principally polarized abelian varieties and we have a suitable anti-equivalence of categories. We give below a direct proof.

Proposition E.1. *Let R be an imaginary quadratic order, and let (M, H_M) be a unimodular Hermitian module over R (torsion free finitely generated as always, but we do not assume M projective here). Let $V = M \otimes_{\mathbb{Z}} \mathbb{Q}$, and $H = H_M \otimes_{\mathbb{Z}} \mathbb{Q}$ the induced Hermitian form on V . Let $\iota_1: M_1 \hookrightarrow M$ be a primitive submodule. Let V_1 be the K -span of M_1 inside V , and $M_1^\sharp$ the R -orthogonal of M_1 in V_1 . Let $V = V_1 \oplus^\perp V_2$ be the orthogonal decomposition given by V and V_1 , and denote by $M_2 = M \cap V_2$ the primitive submodule of M induced by V_2 . Then $M_1^\sharp/M_1 \simeq M_2^\sharp/M_2$.*

Proof. The proof is the same as in Proposition 6.1, except for the argument that $i_1^\sharp$ is surjective. Instead, we rely on biduality: for any lattice N in $V = M \otimes \mathbb{Q}$, $N^{\sharp\sharp} = N$ [KNRR21, p. 6]. Let $U_1 = i_1^\sharp(M) \subset M_1^\sharp$ be the image of M by $i_1^\sharp$, we want to prove that $U_1 = M_1^\sharp$, or equivalently by biduality that $U_1^\sharp = M_1$. Since we know that $U_1 \subset M_1^\sharp$, it suffices to prove that $U_1^\sharp \subset M_1$. But if $v_1 \in U_1^\sharp$, then $H_M(v_1, i_1^\sharp(m)) = H_M(v_1, m) \in R$ for all $m \in M$, so (using that $M^\sharp = M$), $v_1 \in V_1 \cap M = M_1$, as we wanted. $\square$

Remark E.2. When R is not maximal, the proof of Proposition 6.1 gives the following stronger statement: since M/M_1 is torsion free it is projective, hence we get a splitting $M \simeq M_1 \oplus M/M_1$. When R is not maximal, M/M_1 may not be projective anymore, hence we may not get a splitting as above.

Now if (M, H_M) is a unimodular *projective* Hermitian R -module corresponding to (A, λ_A) , A is isomorphic without the polarization to $E_0^{g-1} \times E_I$, where $E_I = I \cdot E_0$. Our higher dimensional Clapotis algorithm (see Section 6.2) works just as well in this setting, because it only relies on the Poincaré decomposition theorem and the fact that the module is locally free when it is projective, and allow us to build a double path $E_0^g \rightarrow A$.

As in Section 6.3, we obtain applying Proposition 5.16:

Proposition E.3. *Given an abstract representation of a projective module (M, H_M) associated with A R -isogenous to E_0^g , we can recover A explicitly, along with the explicit action of the module $\text{Hom}_R(A, E_0)$ and the endomorphism ring $\text{End}_R(R) = \text{End}_R(M)$ on A . Given an isomorphism $M \simeq R^{g-1} \oplus I$, we can also explicitly build the (non polarized) isomorphism $A \simeq E_0^{g-1} \times E_I$.*

Of course, given an explicit representation of $\text{Hom}_R(A, E_0)$, we can also build the Hermitian module (M, H_M) explicitly, this is a special case of Section 5.1.

The main difference with the maximal supersingular case is that in the commutative case, $(\text{End}_R(M), r_M)$ does not give enough information to recover M , hence A . This is already clear in dimension 1, where all ideals I have a canonical Hermitian R -form $H_I = \frac{1}{N(I)} H_R$ where $H_R(x, y) = x\bar{y}$, yet their polarized endomorphism rings are all given by $(R, \cdot)$. The advantage of the module point of view is thus that it allows to treat both cases uniformly.

APPENDIX F. SAMPLING SUITABLE ORTHOGONAL BASIS

To prove Proposition 6.6 we first state few usefull lemmas. First of all, we will rely on the Schwartz-Zippel Lemma:

Lemma F.1. *Let R be an integral domain, and $I \subset R$ be a finite set of elements. Let $Q \in R[x_1, \dots, x_n]$ be a non-zero polynomial of degree d for, whose vanishing set is denoted $Z(P)$. Then $\#Z(P) \cap I^n \leq d \#I^{n-1}$.*

Indeed this lemma will help us to sample elements of M , with norm coprime to some fixed integer. Before using it, we need to be sure that the quadratic form given by q_M is not zero modulo odd primes.

Lemma F.2. *Let (M, H_M) be a unimodular Hermitian module of rank g . Let $M_1 \hookrightarrow M$ be a free submodule of rank r , equipped with the induced Hermitian form. We denote $N = \text{Card}(M_1^\sharp/M_1)$, and $n_2 = v_2(N)$ its 2-adic valuation. For every odd prime ℓ not dividing N , there exists $y_\ell \in M_1$ such that $q_{M_1}(y_\ell) \not\equiv 0 \pmod{\ell}$. Also there exists $y_2 \in M_1$ such that $q_{M_1}(y_2) \not\equiv 0 \pmod{2^{n_2+2}}$.*

Proof. To avoid any confusion, we denote by q_{M_1} (resp. H_{M_1}) the quadratic (resp. Hermitian) form induced on M_1 . We introduce the $\mathbb{Z}$ -bilinear map $H_{\mathbb{Z}} = \text{Trd } H_{M_1}$, and the following notation $M_1^* = \{v \in M_1 \otimes \mathbb{Q} \mid \forall m_1 \in M_1, H_{\mathbb{Z}}(v, m_1) \in \mathbb{Z}\}$, and similarly $\mathcal{O}_0^* = \{v \in \mathcal{B}_{p,\infty} \mid \forall m \in \mathcal{O}_0, \text{Trd}(v\bar{m}) \in \mathbb{Z}\}$. We work by contraposition here.

First suppose that ℓ be an odd prime distinct from p , such that $q_{M_1} \equiv 0 \pmod{\ell}$. Then the $\mathbb{Z}$ -bilinear map $H_{\mathbb{Z}}$ is also zero modulo ℓ . Thus we have $\frac{1}{\ell}M_1 \subset M_1^*$, and there exists an element of order ℓ in M_1^*/M_1 . Now we link M_1^* and $M_1^\sharp$ as follows. First notice that $M_1^* = \{v \in M_1 \otimes \mathbb{Q} \mid \forall m_1 \in M_1, H_M(v, m_1) \in \mathcal{O}_0^*\}$. Moreover at the cost of working locally we may assume that $\mathcal{O}_0^* = \pi/p\mathcal{O}_0$ (where $\pi\bar{\pi} = p$). We thus have $M^* = \pi/pM^\sharp$. Let $m \in M_1^*$ such that $m \notin M_1$ and $\ell m \in M_1$. Writing $m = \pi/pm'$ with $m' \in M_1^\sharp$ leads to the fact that $\ell m' \in M_1$. To see that $m' \notin M_1$, remark that it would imply $m \in \frac{1}{p}M_1 \cap \frac{1}{\ell}M_1$, and since $\ell \neq p$, $m \in M_1$ which is a contradiction. So we have an element of order $\ell \in M_1^\sharp/M_1$, and $\ell \mid N$. This reasoning also gives the last assertion of the proposition.

On the other hand if $q_{M_1} \equiv 0 \pmod{p}$ the same argument implies that $\frac{1}{p}M \subset M^*$, and hence $\frac{1}{\pi}M \subset M^\sharp$. And since $\frac{1}{\pi}M/M \simeq \mathcal{O}_0^g/\pi\mathcal{O}_0^g$ has elements of order p , so do $M^\sharp/M$. $\square$

Lemma F.3. *We keep the notations of Lemma F.2, and D' be an odd integer, coprime with N , and bounded by a polynomial in p . Suppose that we can efficiently compute an $y_0 \in M_1$ such that $q_{M_1}(y_0) \not\equiv 0 \pmod{2^{n_2+2}}$. Then we can efficiently sample an element $y \in M_1$ such that $q_{M_1}(y)$ is coprime with D' , and that $q_{M_1}(y) \not\equiv 0 \pmod{2^{n_2+2}}$.*

Proof. Notice that in order to find $y \in M_1$ with norm coprime to D' and non zero modulo 2^{n_2+2} , it suffices to work locally at each prime divisor ℓ_i of D' . Indeed if, for all $\ell_i \mid D'$, we manage to find an $y_i \in M_1$ such that $q_M(y_i) \not\equiv 0 \pmod{\ell_i}$, then we can combine them using the CRT to recover some $y \in M_1$ such that $q_{M_1}(y)$ is coprime to $\prod \ell_i$, and hence coprime to D' . Remark that for any (odd) prime $\ell_i \mid D'$, $q_{M_1} \not\equiv 0 \pmod{\ell_i}$ by Lemma F.2, since N and D' are coprime. On the other hand, the 2-adic condition on $q_{M_1}(y)$ can be handle thanks to the y_0 given by assumption. So now we can focus only on the ℓ_i dividing D' .

Let B be a positive integer, that will be use as a bound on the primes ℓ_i for the next step. Now for each $\ell_i \leq B$ dividing D' , we can compute $q \pmod{\ell_i}$ and construct in time polynomial in B elements $y_i \in M_1$ such that $q_{M_1}(y_i) \not\equiv 0 \pmod{\ell_i}$.

For other prime factors $\ell'_i > B$ of D' that we do not want to compute for efficiency reasons, we proceed differently. Let s_B be the number of primes $\ell'_i > B$ dividing D' . The idea is now to sample random elements in $y' \in M_1/D'M_1$ and hope that one gives a non zero value mod ℓ'_i for every $i \in \llbracket 1, s_B \rrbracket$. To ensure that we can efficiently find such an y' , we rely on Lemma F.1. Indeed it implies for an $\ell'_i > B$, the reduction $\overline{q_{M_1}}$ of q_{M_1} over $R_i = \mathbb{Z}/\ell'_i\mathbb{Z}$ has at most $2\ell_i^{4r-1}$ roots. This implies that the proportion of isotropic elements of $\overline{q_{M_1}}$ in R_i^{4r} is at most $2/\ell'_i < 2/B$. Therefore the probability $\mathbb{P}_i$ of picking at random an element $v_i \in M_1/\ell'_i M_1$ such that $\overline{q_{M_1}}(v_i) \not\equiv 0 \pmod{\ell'_i}$ is at least $1 - 2/B$. Since the CRT gives a ring isomorphism and that q_{M_1} is polynomial, the elements $v \in M_1/\prod \ell'_i M_1$ with norm coprime with $\prod \ell'_i$ are in bijection with the tuples $(v_1, \dots, v_{s_B}) \in \prod M_1/\ell'_i M_1$ such that $\overline{q_{M_1}}(v_i) \not\equiv 0 \pmod{\ell'_i}$ for all $i \in \llbracket 1, s_B \rrbracket$. Now by independence of the events, the probability of finding a suitable tuple $(v_1, \dots, v_{s_B}) \in \prod M_1/\ell'_i M_1$ equals $\prod \mathbb{P}_i$. Hence the probability $\mathbb{P}$ of finding a good $y' \in M_1/\prod \ell'_i M_1$ is at least $(1 - 1/B)^{s_B}$. We can now bound $s_B \leq \log(D')/\log(B)$, leading to $\mathbb{P} \geq (1 - 1/B)^{\log(D')/\log(B)}$. By setting the bound B to be $\log(p)^2$ and using that $\log(D') \leq C \log(p)$, we have that $\mathbb{P} \geq (1 - \frac{1}{\log(p)})^{2C \log(p)/\log(\log(p))}$. Using the substitution $x = 1/\log(p)$, and

computing the first order expansion of $x \mapsto \exp(\frac{-\log(1-x^2)}{x \log(x)})$ near $x = 0$, we eventually obtain $\mathbb{P} \geq (1 - \frac{1}{\log(p) \log(\log(p))} + o_{p \rightarrow \infty}(\frac{1}{\log(p)}))^C$. Therefore we only need a constant number of retries to sample an $y' \in M_1$ such that $q_M(y') \not\equiv 0 \pmod{\prod \ell'_i}$.

Wrapping up, we can construct in polynomial time in $\log(p)$ an $y \in M_1$ such that $\gcd(q_M(y), D') = 1$ and $v_2(q_M(y)) < n_2 + 2$. $\square$

Proof. of Proposition 6.6.

To construct the desired orthogonal full submodule $M_1 \hookrightarrow M$, we in fact compute the elements $e_1, \dots, e_g \in M$ such that $M_1 = Re_1 \oplus \dots \oplus Re_g$, $i \neq j \Rightarrow H_M(e_i, e_j) = 0$ and $q_M(e_i) = 2^{\epsilon_i} d_i$, where d_i is an odd integer coprime with D' , and $\sum_{i=1}^g \epsilon_i$ is bounded by a polynomial in g (and independant with p). We proceed by iteration over the e_i .

First, we need to sample an element $e_1 \in M$ such that $q_M(e_1) = 2^{\epsilon_1} d_1$, with d_1 odd and coprime to D' . Such an e_1 is precisely given by Lemma F.3, with $\epsilon_1 \in \{0, 1\}$ since M is unimodular.

Second, suppose that we have $(e_1, \dots, e_{r-1})$ satisfying the above conditions, and let $M_1 = \bigoplus_{i=1}^{r-1} Re_i$ seen as a submodule of (M, H_M) (that we can assume to be primitive). Then we can write $N := \text{Card}(M_1^\sharp/M_1) = \prod_{i=1}^{r-1} (q_M(e_i)^{r_z}) = 2^{n_2} d$, where d is odd and coprime with D' . Denoting M_2 the orthogonal of M_1 as constructed in Proposition 6.1, then since $M_2^\sharp/M_2 \simeq M_1^\sharp/M_1$, $\text{Card}(M_2^\sharp/M_2)$ is also coprime with D' . Hence we can apply Lemma F.3 over M_2 to compute e_r , and then we iterate over $M'_1 = M_1 \oplus Re_r$.

In the end we obtain a D_1 -similitude with $v_2(D_1) \leq r_{\mathbb{Z}} g^2$, which leads to accessible torsion. (Recall that $r_{\mathbb{Z}}$ is the rank of R as a $\mathbb{Z}$ -module, hence either 2 or 4.) The coprimality condition follows directly from the constructions of the e_i . $\square$