

# LOCAL-TO-GLOBAL RIGIDITY OF BRUHAT-TITS BUILDINGS

MIKAEL DE LA SALLE AND ROMAIN TESSERA

ABSTRACT. A vertex-transitive graph  $X$  is called local-to-global rigid if there exists  $R$  such that every other graph whose balls of radius  $R$  are isometric to the balls of radius  $R$  in  $X$  is covered by  $X$ . Let  $d \geq 4$ . We show that the 1-skeleton of an affine Bruhat-Tits building of type  $\tilde{A}_{d-1}$  is local-to-global rigid if and only if the underlying field has characteristic 0. For example the Bruhat-Tits building of  $\mathrm{SL}(d, \mathbf{F}_p((t)))$  is not local-to-global rigid, while the Bruhat-Tits building of  $\mathrm{SL}(d, \mathbf{Q}_p)$  is local-to-global rigid.

A vertex-transitive graph  $X$  is called local-to-global rigid (LG-rigid) if there exists  $R$  such that every other graph whose balls of radius  $R$  are isometric to the balls of radius  $R$  in  $X$  is covered by  $X$ . This notion was introduced by Benjaminin and Georgakopoulos and investigated in [BE, G, ST15]. It follows from these works that in many cases, Cayley graphs of finitely presented groups are LG-rigid: for instance all Cayley graphs of torsion-free lattices in simple Lie groups, or Cayley graphs of torsion-free groups of polynomial growth. We also proved ([ST15]) that every finitely presented group which is not a quotient of a Burnside group admits LG-rigid Cayley graphs. On the other hand, in [ST15], we constructed many examples of such graphs which are not LG-rigid: e.g. a Cayley graph of  $F_2 \times F_2 \times \mathbf{Z}/2\mathbf{Z}$ . In this article we investigate LG-rigidity for 1-skeletons of Bruhat-Tits buildings.

By *non-archimedean local field* we will mean a locally compact discrete valuation field (not necessarily commutative). If  $K$  is a non-archimedean local field and  $d \geq 3$ , we denote by  $X_d(K)$  the Bruhat-Tits building of type  $\tilde{A}_{d-1}$  constructed from  $K$ . Our main result characterizes, for  $d \geq 4$ , the fields for which  $X_d(K)$  is LG-rigid.

**Theorem 0.1.** *Let  $K$  be a non-archimedean local field. If  $K$  has positive characteristic and  $d \geq 3$ , then  $X_d(K)$  is not LG-rigid. By contrast, if  $K$  has characteristic 0 and  $d \geq 4$ , then  $X_d(K)$  is LG-rigid.*

Let us discuss the proof of Theorem 0.1. There is a natural locally compact Hausdorff topology on the isomorphism classes of non-archimedean local fields where two fields are  $R$ -close if their residue rings  $\mathcal{O}/\pi^R\mathcal{O}$  are isomorphic. For example,  $\mathbf{Q}_p[p^{1/R}]$  and  $\mathbf{F}_p((t))$  are  $R$ -close. Indeed, the elements of their rings of integers have a unique representation as a formal series  $\sum_{n \geq 0} a_n t^n$  with  $a_n \in \{0, 1, \dots, p-1\}$  and as a formal series  $\sum_{n \geq 0} a_n (p^{1/R})^n$  with  $a_n \in \{0, 1, \dots, p-$

---

*Date:* December 8, 2015.

1} respectively. These two rings have therefore “the same elements”, but the operations are different (without carry for  $F_p((t))$  and with carry for  $\mathbf{Q}_p[p^{1/R}]$ ). However when  $R$  becomes large, the difference becomes smaller and smaller as the carry is sent at distance  $R$ , and in particular  $\mathbf{Q}_p[p^{1/R}]$  and  $\mathbf{F}_p((t))$  are  $R$ -close.

Similarly there is a locally compact Hausdorff topology on the isometry classes of vertex-transitive locally finite graphs where two graphs are  $R$ -close if they have the same balls of radius  $R$ . The idea of the proof of Theorem 0.1 is simple and can be summarized as follows.

- (i) For these topologies, the map  $K \mapsto X_d(K)$  is a homeomorphism on its image.
- (ii) A non-archimedean local field is isolated if and only if it has characteristic 0.
- (iii) If  $d \geq 4$ ,  $\{X_d(K), K \text{ non-archimedean local field}\}$  is open in the set of large-scale simply connected graphs.

The first statement in Theorem 0.1 is immediate from the continuity of  $X_d$  in (i) and from (ii). The second statement follows from (i–iii) and our work [ST15].

We prove in Corollary 2.2 that the map  $K \mapsto X_d(K)$  is continuous (actually 1-Lipschitz for the natural distances). Since it is injective by a deep Theorem of Tits and clearly proper, (i) follows. We could not find a direct proof of (i); for example we could not decide whether  $X_d$  is isometric. The point (ii) is classical, at least as far as commutative fields are concerned [K47, D84]; its simplest illustration is, as recalled above, that  $\mathbf{F}_p((t))$  is the limit of  $\mathbf{Q}_p[p^{1/R}]$  as  $R \rightarrow \infty$ . We recall this in §1. The meaning of (iii) is made precise in Corollary 2.7; it is proved as a consequence of other deep results of Tits which give a local characterization of the graphs  $X_d(K)$  among graphs with a special kind of labelling of the vertices called a *geometry of type  $\tilde{A}_{d-1}$*  and from our Proposition 2.3 where we show that such a labelling can be recovered locally.

The reason why we allow non-commutative fields is not to make the exposition hard to follow: even if we were only interested in commutative fields, we would have to work with non-commutative fields in the proof of the second statement. Indeed, we do not know of a direct proof showing that  $\{X_d(K), K \text{ commutative}\}$  is open (this is true *a posteriori* because the set of commutative non-archimedean local fields is open in the space of all commutative non-archimedean local fields).

Let us state two consequences. The following is a consequence of [ST15, Corollary 1.6] and of Theorem 0.1 (or rather of (i) and of the convergence of  $\mathbf{Q}_p[p^{1/R}]$  to  $\mathbf{F}_p((t))$ ).

**Proposition 0.2.** *Let  $N \in \mathbf{N}$ , then for all but finitely many  $R \in \mathbf{N}$ , the building  $X_d(\mathbf{Q}_p[p^{1/R}])$  does not admit any discrete group of isometry  $\Gamma$  such that*

- *the cardinality of the vertex set of the quotient graph  $X(\mathbf{Q}_p[p^{1/R}])/\Gamma$  is at most  $N$ ;*
- *for all vertex  $x$  of  $X_d(\mathbf{Q}_p[p^{1/R}])$ , the stabilizer of  $x$  in  $\Gamma$  has cardinality at most  $N$ .*

In particular, for all but finitely many  $R \in \mathbf{N}$ ,  $X_d(\mathbf{Q}_p[p^{1/R}])$  is not a Cayley graph. For  $d = 3$ ,  $X_d(\mathbf{F}_p((t)))$  turns out to have a group of isometries acting simply transitively on its vertex set [CMSZ93]. Hence  $X_d(\mathbf{F}_p((t)))$  can be seen as a Cayley graph of this group. In particular, Theorem 0.1 yields to new examples of Cayley graphs of finitely presented groups which are not LG-rigid.

**Theorem 0.3.** *There exists a Cayley graph  $X$  of some finitely presented group, and for each  $R > 0$ , a 2-simply connected vertex transitive graph  $Y_R$  which is  $R$ -locally  $X$ , but is not even quasi-isometric to  $X$ . In particular  $X$  is not LG-rigid.*

**Acknowledgements.** *We thank Sylvain Barré, Laurent Berger, Gaëtan Chenevier, Gabriel Dospinescu, Frédéric Haglund, Pierre Pansu, Mikael Pichot, Vincent Pilloni and Sandra Rozensztajn for useful discussions.*

## 1. NON-ARCHIMEDEAN LOCAL FIELDS

**Definition 1.1.** A non-archimedean local field is a (not necessarily commutative) field which is locally compact for a discrete valuation.

If  $K$  is a non-archimedean local field with a discrete valuation  $v: K \rightarrow \mathbf{Z} \cup \{\infty\}$ , we will always assume that the image of  $v$  is  $\mathbf{Z} \cup \{\infty\}$ , and we will denote its ring of integers  $\mathcal{O} = \{x \in K, v(x) \geq 0\}$  (or  $\mathcal{O}^K$  if we need to keep track of  $K$ ) and  $\mathfrak{m} = \{x \in K, v(x) \geq 1\}$  the unique prime ideal in  $\mathcal{O}$ . Denote by  $\pi$  a uniformizer of  $K$ , that is a generator of the  $\mathcal{O}$ -module  $\mathfrak{m}$ . For an integer  $R \geq 1$  denote by  $\mathcal{O}_R$  the ring  $\mathcal{O}/\pi^R \mathcal{O}$ .

A field  $K$  with a discrete valuation  $v$  is locally compact if and only if it is complete and the residue field  $\mathcal{O}/\pi \mathcal{O}$  is finite.

Two non-archimedean local fields are  $R$ -close if their residue rings  $\mathcal{O}_R$  are isomorphic.

**Theorem 1.2.** *The distance*

$$d(K, K') = \inf\{e^{-R}, K \text{ and } K' \text{ are } R\text{-close}\}$$

*defines a locally compact Hausdorff topology on the isomorphism classes (as topological fields) of non-archimedean local fields.*

*For this topology, a set of non-archimedean local fields is relatively compact if and only if the cardinality of the residue field is bounded on this set.*

*A field is isolated if and only if it has characteristic 0.*

The topology is Hausdorff because a non-archimedean local field is determined as a topological field by the sequence of its residual rings. For the rest of the proof of Theorem 1.2 we will need an explicit description of all non-archimedean local fields. We refer to [W74, Chapter 1] for the statements for which we do not provide a precise reference. If  $p$  is a prime number and  $q = p^f$  for some  $f \geq 1$ , we denote by  $\mathbf{Q}_q$  the totally unramified extension of degree  $f$  of  $\mathbf{Q}_p$ .

Let  $K$  be a *commutative* non-archimedean local field with valuation  $v$ . Its residual field is isomorphic to  $\mathbf{F}_{p^f}$  for a prime number  $p$  called the *residual characteristic of  $K$*  and an integer  $f \geq 1$  called the *absolute residual degree of  $K$* . The value  $e = v(p) \in \mathbf{N} \cup \infty$  is called the *absolute ramification index*. If  $K$  has characteristic 0 (*i.e.*  $e < \infty$ ), then  $K$  is isomorphic to a totally ramified commutative extension of  $\mathbf{Q}_{p^f}$  of degree  $e$  (hence  $K$  is isomorphic to  $\mathbf{Q}_{p^f}[X]/(P)$ , where  $P$  is an Eisenstein polynomial of degree  $e$  with coefficients in  $\mathcal{O}^{\mathbf{Q}_q}$ ). We will use that, for every prime number  $p$  and every integers  $f, e \geq 1$ , there are finitely many [R00, §3.1.6], and at least one (for example  $\mathbf{Q}_{p^f}[p^{1/e}]$ ), commutative non-archimedean local fields with residual characteristic  $p$ , absolute ramification index  $e$  and absolute residual degree  $f$ . If  $K$  has positive characteristic (*i.e.*  $e = \infty$ ),  $K$  is isomorphic to  $\mathbf{F}_{p^f}((t))$ .

**Lemma 1.3.** *If  $K$  is a local field of residual field  $\mathbf{F}_q$  and absolute ramification index  $e$ , then  $K$  and  $\mathbf{F}_q((t))$  are  $e$ -close.*

*Proof.* Note that  $K$  is a totally ramified commutative extension of  $\mathbf{Q}_q$  of degree  $e$ . Hence we can write  $K = \mathbf{Q}_q[X]/(P)$  where  $q = p^f$  and  $P = X^e + a_{e-1}X^{e-1} + \dots + a_0$  is an Eisenstein polynomial of degree  $e$  with coefficients in  $\mathbf{Z}_q = \mathcal{O}^{\mathbf{Q}_q}$ . Recall that  $P$  satisfies  $a_i \in p\mathbf{Z}_q$  for all  $0 \leq i \leq e-1$ , and  $a_0 = bp$  where  $b$  is invertible in  $\mathbf{Z}_q$ . Note that  $\mathcal{O}_e^K = \mathbf{Z}_q[X]/(J)$  where  $J = (X^e) + (P)$ . It follows that in  $\mathcal{O}_e^K$  we have

$$p(b + a_1X + \dots + a_{e-1}t^{e-1}) = 0,$$

from which we deduce that  $p = 0$ . On the other hand, modulo  $p$ , one has  $P = X^e$ . So finally we deduce that

$$\mathcal{O}_e^K \simeq \mathbf{Z}_q[X]/((X^e) + (p)) \simeq (\mathbf{Z}_q[X]/(p))/(X^e) \simeq \mathbf{F}_q[X]/(X^e) \simeq \mathcal{O}_e^{\mathbf{F}_q((t))},$$

and we are done.  $\square$

Let us now move to non-commutative fields. If  $K$  is a non-archimedean local field with center  $L$ , the residual field  $\mathbf{k}$  of  $K$  is an extension of the residual field  $\mathbf{l}$  of  $L$ ; denote by  $d$  the degree of  $\mathbf{k}/\mathbf{l}$  ( $d$  is called the *residual degree of  $K/L$* ). The Galois group  $\text{Gal}(\mathbf{k}/\mathbf{l})$  is cyclic of order  $d$  with generator the Frobenius automorphism. Moreover, if  $\pi$  is a uniformizer of  $K$ , the conjugation  $x \in K \mapsto \pi^{-1}x\pi$  belongs to  $\text{Gal}(K/L)$ . Its image  $\alpha$  in  $\text{Gal}(\mathbf{k}/\mathbf{l})$  does not depend on the choice of the uniformizer because  $\mathbf{k}$  is commutative, and corresponds to the  $r$ -th power of the Frobenius for some  $r \in \mathbf{Z}/d\mathbf{Z}$ . For convenience we shall call  $r$  the *Hasse invariant* of  $K$ . The next lemma states that  $r$  is a generator of  $\mathbf{Z}/d\mathbf{Z}$ , and that the triple  $(L, d, r)$  with  $d \geq 1$  and  $r$  a generator of  $\mathbf{Z}/d\mathbf{Z}$  determines a unique non-commutative field  $K$ . Observe that the case where  $K$  is commutative corresponds to  $d = 1$ , in which case  $r = 0$  (which is a generator of the trivial group  $\mathbf{Z}/\mathbf{Z}$ ).

**Lemma 1.4.** ([H32],[W74, Chapter 1 (p20–22), Chapter XII]) *Let  $K, L, d, r$  as above. Then  $r$  is a generator of  $\mathbf{Z}/d\mathbf{Z}$ . Conversely, for every commutative non-archimedean local field  $L$ , every integer  $d \geq 1$  and every generator  $r$  of  $\mathbf{Z}/d\mathbf{Z}$ , there is a unique non-archimedean local field  $K$  with center  $L$ , residual degree  $d$  over  $L$  and Hasse invariant  $r$ . It has degree  $d^2$  over  $L$  and can be described as follows. It contains a maximal commutative extension  $K_1$  of  $L$  of degree  $d$  which is unramified. Moreover,  $L$  has a uniformizer  $\pi$  and  $K$  has a uniformizer  $x$  such that  $x^d = \pi$ ,  $(1, x, \dots, x^{d-1})$  forms a basis of  $K$  as a  $K_1$ -vector space, and for all  $a \in K_1$ ,  $x^{-1}ax = f^r(a)$ , where  $f$  is the unique automorphism of  $K_1$  inducing the Frobenius automorphism of  $\text{Gal}(\mathbf{k}_1/\mathbf{l})$ .*

We will say that a non-archimedean local field  $K$  has type  $(p, f, e, d, r)$  if its center  $L$  has residual characteristic  $p$ , absolute ramification index  $e$  and absolute residual degree  $f$ , and if the extension  $K/L$  has residual degree  $d$  and Hasse invariant  $r$ . From the preceding discussion we conclude that for every prime number  $p$ , every integers  $f, d \geq 1$ , every  $e \in \mathbf{N} \cup \{\infty\}$  and every  $r \in (\mathbf{Z}/d\mathbf{Z})^*$ , the number of fields of type  $(p, f, e, d, r)$  is finite and nonzero. As we have seen, there exists a unique field of type  $(p, f, \infty, d, r)$ . If  $q = p^f$ , it can be concretely defined as the quotient of the  $\mathbf{F}_q((t))$ -algebra freely generated by  $\mathbf{F}_{q^d}((t))$  and by an element  $x$ , by the ideal generated by the following relations:  $x^d = t$ , and  $ax = xf(a)$ , for all  $a \in \mathbf{F}_{q^d}((t))$ , where  $f$  is the automorphism of  $\mathbf{F}_{q^d}((t))$  uniquely defined by  $f(t) = t$ , and  $f(z) = z^{q^r}$  for all  $z \in \mathbf{F}_{q^d}$ .

**Lemma 1.5.** *Let  $K$  (resp.  $F$ ) be respectively a field of type  $(p, f, e, d, r)$  (resp. the field of type  $(p, f, \infty, d, r)$ ). Then  $K$  and  $F$  are  $ed$ -close: i.e. the residue rings  $\mathcal{O}_{ed}^K$  and  $\mathcal{O}_{ed}^F$  are isomorphic.*

*Proof.* With the notation of Lemma 1.4,  $K$  is isomorphic to the quotient  $\tilde{K}$  of the ring freely generated by  $K_1$  and  $x$ , by the ideal generated by the relations  $x^d = \pi$ , and  $ax = xf^r(a)$ , for all  $a \in K_1$ , where  $f$  induces the Frobenius on the residue field. Indeed, one clearly has a morphism of  $L$ -algebras from  $\tilde{K}$  to  $K$  that is the identity on  $K_1$  and on  $x$ . The fact this is an isomorphism follows by comparing the dimensions over  $L$ . We deduce that the residue ring  $\mathcal{O}_{ed}^K = \mathcal{O}^K/(x^{ed}) = \mathcal{O}^K/(\pi^e)$  is isomorphic to the quotient of the ring freely generated by  $\mathcal{O}^{K_1}/(\pi^e)$  and  $x$ , by the ideal generated by the relations  $x^d = \pi$ , and  $ax = x\tilde{f}^r(a)$ , where  $\tilde{f}$  is the unique (by Hensel's Lemma) automorphism of the ring  $\mathcal{O}^{K_1}/(\pi^e)$  which induces the Frobenius on the residue field. The same description of course applies to  $\mathcal{O}_{ed}^F$ , and we conclude by Lemma 1.3 because  $K_1$  has absolute residual degree  $fd$  and absolute ramification index  $e$ .  $\square$

We can now prove Theorem 1.2. To do so we show that the balls of radius  $\frac{1}{2}$  are compact (here  $\frac{1}{2}$  could be any number in  $[\frac{1}{e}, 1)$ ), and that their only accumulation points are the fields of positive characteristic. Since two fields are at distance less than  $\frac{1}{2}$  if and only if they have the same residue field, it amounts to investigating,

for every finite field  $\mathbf{F}_q$ , the set of fields having  $\mathbf{F}_q$  as residue field. This set contains exactly the fields of type  $(p, f, e, d, r)$  for  $q = p^{fd}$  and  $k \in \mathbf{N} \cup \{\infty\}$ . This determines  $p$  and forces  $f, d$  to take only finitely many values. Therefore (since there are finitely many fields of each type) a sequence of such fields either has a stationary subsequence, or a subsequence of type  $(p, f, e_n, d, r)$  for a sequence  $e_n \rightarrow \infty$ , which converges to the field of type  $(p, f, \infty, d, r)$  by Lemma 1.5. This shows that the set of fields with residue field  $\mathbf{F}_q$  is compact, and that the fields with characteristic 0 are isolated. Conversely, every nonarchimedean local field  $K$  of characteristic  $p > 0$  is the field of type  $(p, f, \infty, d, r)$  for some  $f, d, r$ . As we discussed there is a sequence of fields of type  $(p, f, n, d, r)$ , and it converges as  $n \rightarrow \infty$  to  $K$  by Lemma 1.5.

## 2. BUILDINGS

**2.1. Graphs.** In this paper “a graph” means a connected, locally finite, simplicial graph without multiple edges and loops. It is called vertex-transitive if its isometry group acts transitively on the set of vertices. A graph  $Y$  is  $R$ -locally  $X$  if every ball of radius  $R$  around a vertex in  $Y$  is isometric to a ball of radius  $R$  around a vertex in  $X$ . This defines a locally compact Hausdorff topology on the isomorphism classes of transitive graphs, for example for the distance

$$\inf\{e^{-R}, Y \text{ is } R\text{-locally } X\}.$$

A set of vertex-transitive graphs is relatively compact if and only if the degree is bounded on this set.

**2.2. Classical buildings of type  $\tilde{A}_{d-1}$ .** Let  $d \geq 2$ . Let us recall the description of the building of  $GL(d, K)$  (the building  $\tilde{A}_{d-1}(K, v)$ ) associated to a nonarchimedean local field  $K$  with discrete valuation  $v$ , see [R85, Chapter 9] for details.

An  $\mathcal{O}$ -lattice in  $K^d$  is a finitely generated  $\mathcal{O}$ -submodule which generates  $K^d$  as a  $K$ -vector space. Such a module is free of rank  $d$ , *i.e.* of the form  $\mathcal{O}v_1 + \dots + \mathcal{O}v_d$  for a basis  $(v_1, \dots, v_d)$  of  $K^d$ . By the invariance property  $a\mathcal{O} = \mathcal{O}a = \pi^k\mathcal{O}$  for any  $a \in K^*$  and  $k \in \mathbf{Z}$  with  $v(a) = k$ , we see that if  $L$  is an  $\mathcal{O}$ -lattice and  $a \in K^*$ ,  $aL$  is also a lattice, so that it makes sense to talk about lattices modulo homothety.

The building  $\tilde{A}_{d-1}(K, v)$  is a simplicial complex of dimension  $d - 1$ . Its 1-skeleton, that we denote by  $X_d(F)$  (or  $X$  for short if there is no ambiguity) is described as follows. The vertices of  $X$  are the  $\mathcal{O}$ -lattices in  $K^d$  modulo homothety. There is an edge between two different vertices  $x$  and  $y$  if there are representatives  $L_1$  and  $L_2$  of  $x$  and  $y$  such that  $\pi L_1 \subset L_2 \subset L_1$ . This is the vertex transitive graph  $X_d(F)$  we are interested in.

**2.3. Continuity of  $K \mapsto X_d(K)$ .** A lattice modulo homothety  $x$  has a unique representative, denoted by  $L(x)$ , contained in  $\mathcal{O}^d$  but not in  $\mathfrak{m}^d$ . There is an edge between two different vertices  $x$  and  $y$  if and only if  $\pi L(x) \subset L(y) \subset L(x)$  or  $\pi L(y) \subset L(x) \subset L(y)$ .

The following Lemma expresses that the ball of radius  $R$  around  $\mathcal{O}^d$  in  $X$  is entirely described in terms of the ring  $\mathcal{O}_R$ .

**Lemma 2.1.** *A lattice modulo homothety  $x$  belongs to the ball of radius  $R$  around  $o$  if and only if  $\pi^R \mathcal{O}^d \subset L(x)$ .*

*Moreover the map  $\bar{L}: x \mapsto L(x) \bmod \pi^R \mathcal{O}^d$  is a bijection between the ball of radius  $R$  around  $\mathcal{O}^3$  in  $X$  and the  $\mathcal{O}_R$ -submodules of  $(\mathcal{O}_R)^d$  not contained in  $(\pi \mathcal{O}_R)^d$ .*

*Lastly two different vertices  $x$  and  $y$  in the ball of radius  $R$  around  $\mathcal{O}^d$  in  $X$  are adjacent if and only if  $\pi \bar{L}(x) \subset \bar{L}(y) \subset \bar{L}(x)$  or  $\pi \bar{L}(y) \subset \bar{L}(x) \subset \bar{L}(y)$ .*

*Proof.* It is immediate that  $\pi^R \mathcal{O}^d \subset L(x)$  if  $d(x, o) \leq R$ . The converse follows by applying, for any lattice  $L \subset \mathcal{O}$ , the invariant factor decomposition over  $\mathcal{O}$ -modules ([T37]) to  $\mathcal{O}^d/L$ , which provides a basis  $v_1, \dots, v_d$  for the  $\mathcal{O}$ -module  $\mathcal{O}^d$  and integers  $n_1 \leq \dots \leq n_d$  such that  $\pi^{n_1} v_1, \dots, \pi^{n_d} v_d$  is a basis for  $L$ . For the lattice  $L(x)$ , we have  $n_1 = 0$ , and if  $\pi^R \mathcal{O}^d \subset L(x)$ , we have  $n_d \leq R$ . If  $x_k$  is the equivalence class of the lattice  $\bigoplus_{1 \leq i \leq d} \mathcal{O} \pi^{\min(k, n_i)} v_i$  then  $x_k$  and  $x_{k+1}$  are adjacent in  $X_d$ ,  $x_0 = o$  and  $x_R = x$ , which shows that  $d(x, y) \leq R$ .

Since  $L \mapsto L \bmod \pi^R \mathcal{O}^d$  is a bijection between the lattices  $L$  such that  $\pi^R \mathcal{O}^d \subset L \subset \mathcal{O}^d$  and the  $\mathcal{O}_R$ -submodules of  $\mathcal{O}_R^d$ , the second statement is immediate from the first.

The last statement is easy. □

We immediately deduce that if two local fields  $K, K'$  are  $R$ -close, then the graphs  $X_d(K)$  is  $R$ -locally  $X_d(K')$ .

**Corollary 2.2.** *The ball of radius  $R$  in  $X_d(K)$  does only depend (up to isometry) on the ring  $\mathcal{O}_R$ .*

*Proof.* By Lemma 2.1 the ball only depends on the pair  $\pi \mathcal{O}_R \subset \mathcal{O}_R$ . But  $\pi \mathcal{O}_R$  is determined by  $\mathcal{O}_R$  as its unique maximal ideal. □

**2.4.  $\{X_d(K)\}$  is open.** We start by recalling some material from [T81].

For an integer  $m \geq 2$ , a generalized  $m$ -gon is a connected bipartite graph of diameter  $m$  and girth  $2m$ , in which every vertex has degree at least 2.

A Coxeter diagram over  $I$  is a function  $M: I \times I \rightarrow \mathbf{N} \cup \{\infty\}$  such that for all  $i, j \in I$ ,  $M(i, i) = 1$  and  $M(i, j) = M(j, i) \geq 2$  if  $i \neq j$ . A symmetry of a Coxeter diagram  $M$  is a permutation  $\sigma$  of  $I$  satisfying  $M(i, j) = M(\sigma(i), \sigma(j))$  for all  $i, j \in I$ .

If  $I$  is a set, a geometry over  $I$  is a pair  $(X, \tau)$  where  $X$  is a graph and  $\tau: X \rightarrow I$  is a coloring of the vertices of  $X$  by labels in  $I$  satisfying that every pair of adjacent vertices have a different label. In a geometry, a complete subgraph is called a

flag, and its type is the subset of  $I$  defined as its image by  $\tau$ . The residue of a flag  $Z$  of type  $J \subset I$  is the geometry over  $I \setminus J$  given by  $(Y, \tau|_Y)$  where  $Y$  is the set of vertices in  $X \setminus Z$  adjacent to  $Z$ , with the same edges as in  $X$ . By convention the residue of the empty flag is  $(X, \tau)$ .

If  $M$  is a Coxeter diagram over a set  $I$ , a geometry of type  $M$  is a geometry  $(X, \tau)$  over  $I$  where for any subset  $J \subset I$ , the residue of any flag of type  $J$  is (1) nonempty if  $|I \setminus J| \geq 1$ , (2) connected if  $|I \setminus J| \geq 2$ , (3) a generalized  $M(i, j)$ -gon if  $J = I \setminus \{i, j\}$  for some  $i \neq j \in I$ . We say that a graph admits a geometry of type  $M$  if there exists  $\tau: X \rightarrow I$  such that  $(X, \tau)$  is a geometry of type  $M$ .

The example important for us is the Coxeter diagram  $\tilde{A}_{d-1}$  over  $\mathbf{Z}/d\mathbf{Z}$  given by where  $\tilde{A}_{d-1}(i, i) = 1$ ,  $\tilde{A}_{d-1}(i, j) = 3$  if  $i - j \in \{-1, 1\}$  and  $\tilde{A}_{d-1}(i, j) = 2$  otherwise. Then  $X_d(K)$  admits a geometry of type  $\tilde{A}_{d-1}$  (see [R85]). It is characterized by the following properties. The origin  $o$  is labelled by  $\tau(o) = 0$ , and if  $x$  and  $y$  are two adjacent vertices with representatives  $L_1$  and  $L_2$  such that  $\pi L_1 \subset L_2 \subset L_1$ , then  $L_1/\pi L_1$  is a vector space of dimension  $d$  over the finite field  $\mathcal{O}/\mathfrak{m}$  and the dimension (modulo  $d$ ) of the image of  $L_2$  inside it is equal to  $\tau(y) - \tau(x)$ .

A particular case of a theorem of Tits [T81, Theorem 1.3] characterizes the buildings of type  $\tilde{A}_{d-1}$  as the simply connected geometries of type  $\tilde{A}_{d-1}$ . This motivates the following result, which shows that for a large scale simply connected graph  $Y$ , admitting a geometry of type  $M$  is a local property.

**Proposition 2.3.** *Let  $M$  be a Coxeter diagram over a finite set  $I$ . Let  $X$  be a 3-simply connected graph. Assume that every ball of radius 3 in  $X$  is isomorphic to a ball of radius 3 in a graph admitting a geometry of type  $M$ . Then  $X$  admits a geometry of type  $M$ .*

*Remark 2.4.* More generally if  $X$  is  $k$ -simply connected, then  $X$  admits a geometry of type  $M$  if every ball of radius  $\lfloor \frac{k+3}{2} \rfloor$  in  $X$  is isometric to a ball of the same radius in a graph admitting geometry of type  $M$ .

If  $X$  is a graph and  $x \in X$ , let  $V(x)$  be the graph with vertex set  $\{x' \in X, d(x, x') \leq 1\}$  and same edges as in  $X$ . A *germ of geometry of type  $M$*  at  $x$  is a coloring  $\tau: V(x) \rightarrow I$  such that  $(V(x), \tau)$  is a geometry over  $I$  and such that the conditions (1) (2) (3) hold for every flag in  $V(x)$  containing  $x$ . Denote by  $G(x)$  the set of all germs of geometry of type  $\tilde{A}_{d-1}$  at  $x$ . Observe that for a connected graph  $X$  and a map  $\tau: X \rightarrow I$ ,  $(X, \tau)$  is a geometry of type  $I$  if and only if the restriction of  $\tau$  to  $V(x)$  is a germ of geometry of type  $M$  for every  $x \in X$ . It is through this observation that we will construct a suitable labelling of a graph satisfying the local properties of Proposition 2.3.

We start by a lemma which implies that a germ of geometry of type  $M$  at  $x$  is characterized by its restriction to any flag of type  $I$ .

**Lemma 2.5.** *Let  $X$  be a graph and  $x \in X$ . If  $\tau \in G(x)$ ,  $G(x)$  consists of all maps of the form  $\sigma \circ \tau$  for a symmetry  $\sigma$  of  $M$ .*

*Proof.* It is clear that  $\sigma \circ \tau \in G(x)$  if  $\sigma$  is a symmetry of  $M$ .

To see the converse take  $\tau' \in G(x)$ . Let  $Z$  be a flag of type  $I$  containing  $x$  (such a flag exists by (1)). Then for every  $z_1 \neq z_2 \in Z \setminus \{x\}$ , by looking at the residue of  $Z \setminus \{z_1, z_2\}$  we obtain from condition (3) that  $M(\tau(z_1), \tau(z_2)) = M(\tau'(z_1), \tau'(z_2))$ . This implies that there exists a unique symmetry  $\sigma_Z$  of  $M$  such that  $\tau'(z) = \sigma_Z \circ \tau(z)$  for all  $z \in Z$ . We claim that  $\sigma_Z = \sigma_{Z'}$  for every pair of flags  $Z$  and  $Z'$  of cardinality  $d$  containing  $x$ . The claim is proved by downwards induction on the cardinality of  $Z \cap Z'$ . The case when  $|Z \cap Z'| = |I|$  or  $|Z \cap Z'| = |I| - 1$  is obvious. Assuming that the claim is valid when  $|Z \cap Z'| = k \leq |I| - 1$ , let us prove the claim when  $|Z \cap Z'| = k - 1$ . Pick  $z \in Z \setminus Z'$  and  $z' \in Z' \setminus Z$ . By (2) there is a path  $z_0, \dots, z_n$  contained in the residue of  $Z \cap Z'$  such that  $z_0 = z$  and  $z_n = z'$ . By (1) for each  $i = 0, \dots, n - 1$  there is a flag  $Z_i$  of type  $I$  containing  $(Z \cap Z') \cup \{z_i, z_{i+1}\}$ , and by induction hypothesis applied to  $Z_i$  and  $Z_{i+1}$  we have  $\sigma_{Z_i} = \sigma_{Z_{i+1}}$  for each  $0 \leq i \leq n - 1$ . This proves that  $\sigma_Z = \sigma_{Z'}$ . Hence  $\sigma_Z$  does not depend on  $Z$ , which proves the lemma.  $\square$

From it we deduce the following

**Lemma 2.6.** *Let  $X$  be a graph admitting a geometry of type  $M$  and  $x$  a vertex in  $X$ . For every germ  $\tau$  of geometry of type  $M$  at  $x$ , there exists  $\tilde{\tau}: X \rightarrow I$  which extends  $\tau$  and such that  $(X, \tilde{\tau})$  is a geometry of type  $M$ .*

*Proof.* Let  $\tau_0: X \rightarrow I$  such that  $(X, \tau_0)$  is a geometry of type  $I$ . By Lemma 2.5, there is a symmetry  $\sigma$  of the diagram  $M$  such that  $\tau$  coincides with  $\sigma \circ \tau_0$  on  $V(x)$ . Then  $\tilde{\tau} = \sigma \circ \tau_0$  extends  $\tau$  and satisfies that  $(X, \tilde{\tau})$  is a geometry of type  $M$ . This shows the existence.  $\square$

With these two lemmas we can proceed to the proof of Proposition 2.3. The argument is the same as for the proof of [ST15, Theorem C]. Since every ball of radius 3 in  $X$  is isometric to a ball in a graph admitting a geometry of type  $M$ ,  $G(x)$  is nonempty for every  $x \in X$ , and it follows from Lemma 2.6 that if  $x, x'$  are two neighbors in  $X$ , then for every  $\tau \in G(x)$ , there is  $\tau' \in G(x')$  which coincides with  $\tau$  on  $V(x) \cap V(x')$ . It is unique because by Lemma 2.5,  $\tau'$  is determined by its value on any complete subgraph with  $d$  vertices containing  $x'$  (and there is such a subgraph containing both  $x$  and  $x'$  by (1)). This defines a bijection that we denote  $F_{x,x'}: G(x) \rightarrow G(x')$ .

For every path  $\gamma = (x_0, \dots, x_n)$  of adjacent vertices, we can define a bijection  $F_\gamma: G(x_0) \rightarrow G(x_n)$  by composing the bijections  $F_{x_i, x_{i+1}}$  along  $\gamma$ . We claim that  $F_\gamma$  only depends on the endpoints  $x_0$  and  $x_n$ . Since  $X$  is 3-simply connected, we only have to check that  $F_\gamma$  is the identity of  $A(x_0)$  if  $\gamma$  is a path of length  $n \leq 3$  with  $x_0 = x_n$ . This property clearly holds if  $X$  admits a geometry of type  $M$ , and hence also in  $X$  because  $\cup_{k \leq n} V(x_k)$  (and all its edges) is contained in the ball of radius 3 around  $x_0$ , which is isometric to a ball of radius 3 in a graph admitting a geometry of type  $M$ .

It remains to fix a vertex  $x_0 \in X$  and  $\tau_0 \in G(x_0)$ . For every other vertex  $x$ , let  $\tau_x \in G(x)$  be the common value of  $F_\gamma(\tau_0)$  for all paths  $\gamma$  from  $x_0$  to  $x$ . Define  $\tau(x) = \tau_x(x)$ . Since for adjacent edges  $x, x'$ ,  $\tau_{x'} = F_{x,x'}(\tau_x)$  coincides with  $\tau_x$  on  $V(x) \cap V(x')$ , we see that  $\tau$  coincides with  $\tau_x$  on  $V(x)$ . In particular the restriction of  $\tau$  to  $V(x)$  belongs to  $G(x)$  for all  $x \in X$ . This means that  $(X, \tau)$  is a geometry of type  $\tilde{A}_{d-1}$ . This concludes the proof of Proposition 2.3.

We deduce the following

**Corollary 2.7.** *Let  $d \geq 4$ . Let  $Y$  be a 3-simply connected graph which is 3-locally  $X_d(K)$  for some non-archimedean local field  $K$ . Then  $Y$  is isometric to  $X_d(F)$  for a (unique up to isomorphism) non-archimedean local field  $F$ .*

*Proof.* From the discussion before Proposition 2.3,  $Y$  is isometric to the 1-skeleton of a locally finite building of type  $\tilde{A}_{d-1}$  (and this holds whenever  $d \geq 3$ ). By a theorem of Tits, this forces  $Y$  to be isometric to  $X_d(F)$  for some  $F$  if  $d \geq 4$ . Moreover,  $X_d(F)$  determines the projective space  $PG(d-1, F)$  up to collineation, which determines  $F$  up to isomorphism by the fundamental theorem of projective geometry. See [R85, page 137], or [T86, Corollaire 15] and [T74, Theorem 6.3].  $\square$

### 3. PROOF OF THEOREM 0.1

The map  $K \mapsto X_d(K)$  is continuous by Corollary 2.2. It is injective by the theorems of Tits recalled in the proof of Corollary 2.7. Let us show that it is proper, that is that if the cardinality of the residue field  $K_n$  goes to  $\infty$ , then the degree in  $X_d(K_n)$  also. But this holds because (Lemma 2.1) the degree in  $X_d(K)$  is equal to the number of linear subspaces of dimension  $\neq 0, d$  in  $^1(\mathcal{O}/\pi\mathcal{O})^d$ . This implies (i):  $K \mapsto X_d(K)$  is a homeomorphism on its image.

The first part of Theorem 0.1 follows from Theorem 1.2 and (i).

Let us now prove the second half of Theorem 0.1. Let  $d \geq 4$  and  $K$  be a field of characteristic 0. By Theorem 1.2 and (i), there exists  $R > 0$  such that if  $K'$  is another non-archimedean local field such that  $X_d(K')$  is  $R$ -locally  $K$ , then  $K'$  is isomorphic to  $K$ , and in particular  $X_d(K')$  is isometric to  $X_d(K)$ . By Corollary 2.7 this implies that if  $Y$  is a 3-simply connected graph which is  $\max(3, R)$ -locally  $X_d(K)$ , it is isometric to  $X_d(K)$ . By [ST15, Proposition 1.5]  $X_d(K)$  is LG-rigid.

### REFERENCES

- [Ba97] L. Babai. The growth rate of vertex-transitive planar graphs. In *Proceedings of the Eighth Annual ACM-SIAM Symposium on Discrete Algorithms (New Orleans, LA, 1997)*, pages 564–573. ACM, New York, 1997.
- [B13] I. Benjamini. *Coarse geometry and randomness*, volume 2100 of *Lecture Notes in Mathematics*. Springer, Cham, 2013. Lecture notes from the 41st Probability Summer School held in Saint-Flour, 2011.

---

<sup>1</sup>Precisely, if the residue field of  $K$  is  $\mathbf{F}_q$ , the degree equals  $\prod_{i=1}^d (q^i - 1)/(q - 1) = \prod_{i=1}^d (\sum_{v=0}^{i-1} q^v)$ , which is a strictly increasing function of  $q$ .

- [BE] I. Benjamini, D. Ellis. The structure of graphs which are locally indistinguishable from a lattice. Preprint.
- [CH] Y. Cornulier, P. de la Harpe. Metric geometry of locally compact groups. 228 pp. Book in progress. arXiv:1403.3796v3
- [CMSZ93] D. I. Cartwright, A. M. Mantero, T. Steger, A. Zappa, Groups acting simply transitively on the vertices of a building of type  $A_2$  I, *Geom. Dedicata* 47 (1993), no. 2, 143–166.
- [D84] P. Deligne. Les corps locaux de caractéristique  $p$ , limites de corps locaux de caractéristique 0. In *Representations of reductive groups over a local field*, Travaux en Cours, pages 119–157. Hermann, Paris, 1984.
- [F01] A. Furman. Mostow-Margulis rigidity with locally compact targets. *Geom. Funct. Anal.*, 11(1) (2001), 30–59.
- [G] A. Georgakopoulos. On covers of graphs by Cayley graphs. Preprint.
- [Gr93] M. Gromov. Asymptotic invariants of infinite groups. In G. Niblo and M. Roller (Eds.), *Geometric group theory II*, number 182 in LMS lecture notes. Camb. Univ. Press, 1993.
- [H32] H. Hasse. Theory of Cyclic Algebras Over An Algebraic Number Field. *Trans. Amer. Math. Soc* 34, no. 1 (1932), 171–171.
- [KL97] B. Kleiner, B. Leeb, Rigidity of quasi-isometries for symmetric spaces and Euclidean buildings, *Inst. Hautes Etudes Sci. Publ. Math.* no. 86 (1997), 115–197.
- [K47] M. Krasner. Théorie non abélienne des corps de classes pour les extensions finies et séparables des corps valués complets: approximation des corps de caractéristique  $p \neq 0$  par ceux de caractéristique 0; modifications de la théorie. *C. R. Acad. Sci. Paris*, 224:434–436, 1947.
- [KM08] B. Krön, R. G. Möller. Quasi-isometries between graphs and trees. *Journal of Combinatorial Theory, Series B* 98 (2008) 994–1013.
- [R00] A. M. Robert. *A course in  $p$ -adic analysis*, volume 198 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2000.
- [R85] M. Ronan. *Lectures on buildings*, volume 7 of *Perspectives in Mathematics*. Academic Press, Inc., Boston, MA, 1989.
- [ST15] M. de la Salle, R. Tessera. Characterizing a vertex-transitive graph by a large ball.
- [T37] O. Teichmüller. Der Elementarteilersatz für nichtkommutative Ringe, *S. Ber. Preuss. Akad. Wiss.* (1937), 169–177.
- [T74] J. Tits. *Buildings of spherical type and finite BN-pairs*. Lecture Notes in Mathematics, Vol. 386. Springer-Verlag, Berlin-New York, 1974.
- [T81] J. Tits. A local approach to buildings, in *The Geometric Vein (Coxeter Festschrift)*, Springer-Verlag, (1981), 317–322.
- [T86] J. Tits. Immeubles de type affine. In *Buildings and the geometry of diagrams (Como, 1984)*, volume 1181 of *Lecture Notes in Math.*, pages 159–190. Springer, Berlin, 1986.
- [T85] V.I. Trofimov. Graphs with polynomial growth, *Math. USSR Sb.*, 51, No. 2 (1985), 405–417.
- [W74] A. Weil. *Basic number theory*. Springer-Verlag, New York-Berlin, third edition, 1974. Die Grundlehren der Mathematischen Wissenschaften, Band 144.

UMPA, ENS-LYON, LYON, FRANCE  
*E-mail address:* mikael.de.la.salle@ens-lyon.fr

LABORATOIRE DE MATHÉMATIQUES, UNIVERSITÉ PARIS-SUD 11, ORSAY, FRANCE  
*E-mail address:* romtessera@gmail.com