

Leureux 1: Boundary amenability of groups acting on buildings

Def (Γ) T has property (A) if $\exists H$ Hilbert, $p_{1,2} \uparrow_{\infty}$
 $f: T \rightarrow H$

$$p(d(x,y)) \leq \|f(x) - f(y)\| \leq p_2(d(x,y))$$

Notation If T has property A then it satisfies the

Novikov conjecture.

Thm (Higson Roe) T has property A if $\exists K$ compact space
 $T \curvearrowright K$ which is topologically amenable i.e. $\exists p_2: K \rightarrow \text{Prob}(T)$

s.t. $\|P_n(\gamma x) - \gamma P_n(x)\| \xrightarrow{n \rightarrow \infty} 0$ uniformly ~~acting~~

Groups which have property A.

- 1) Amenable groups.
- 2) Hyperbolic groups (Adams)
- 3) linear groups (Guetner, Higson, Weinberger)
- 4) groups acting properly on f.-dim. CAT(0) cube complexes
 (Brodzki, Campbell, Guetner, Niblo, Wright)


The G acting properly on a locally finite building then it has property A. In fact its action on infinite boundary is amenable.

Buildings

Coxeter groups

$$W = \langle s \in S, (st)^{m_{st}} = 1 \rangle$$

$$m_{st} \in \mathbb{N}^* \cup \{\infty\}, m_{ss} = 1.$$

Examples  chamber
 Affine Coxeter groups.



Fact (Davis-Thoussang)

Every Coxeter group acts properly discontinuously on some CAT(0) space Σ .

Reflection conjugate of ~~SES~~ SES

Wall = $\{ \text{fixed points of a reflection} \}$

Chamber = connected component of $\Sigma \setminus \{ \text{walls} \}$

Panel = Intersection of two adjacent chambers

Def A building is a gluing of chambers along their panels, covered by apartments, such that

1) every apartment is $\cong \Sigma$

2) any two points are in some apartment

3) For any two apartments A, A' , \exists isom $A = A'$ fixing AAA' .


Example A tree without end points, $W = \mathbb{Z}/2\mathbb{Z} * \mathbb{Z}/2\mathbb{Z}$

Buildings associated to semi-simple groups over local fields
(e.g. $SL_2(\mathbb{Q}_p)$). class of Euclidean buildings.

Kac-Moody groups.

II Combinatorial boundary (joint with Caprace)

Fact For every chamber C , panel σ , there is a unique chamber adjacent to σ , and at a minimal distance from C , denoted $\text{proj}_\sigma(C)$.



$i_{ch} \text{ch}(X) \rightarrow \prod_{\sigma \text{ panel}} \text{link}(\sigma)$ is injective.


$C \mapsto (\sigma \mapsto \text{proj}_\sigma(C))$

Def $\mathcal{C}_{ch}(X) = i_{ch}(\text{ch}(X))$

is the combinatorial compactification.

Lemma 2

Ex



compactification of the apartment.


Hyperbolic case



$\mathcal{E}ch(X) \rightarrow \partial_\infty \mathbb{H}^2$

is surjective with cardinality 1 or 2
(2 for ~~the~~ a limit of a wall at infinity)

Tool Sectors :



$\bar{z} \in \mathcal{E}ch(X)$
 $x \in ch(X)$

$Q(x, \bar{z})$ is a "convex hull of x and \bar{z} ."

More precisely, if $C_n \rightarrow \bar{z}$, $Conv(x, C_n)$ converges pointwise
and $Q(x, \bar{z}) = \lim conv(x, C_n)$.

Th If $\bar{z} \in \mathcal{E}ch(X)$, $x, y \in ch(X)$

then $Q(x, \bar{z}) \cap Q(y, \bar{z}) \supseteq Q(z, \bar{z})$ for some z .

Co Every point in $\mathcal{E}ch(X)$ is in the boundary of
some apartment.

(because every sector is contained in an apartment).


Th (Caprace - L.) G amenable group acting properly on X , then G
is amenable $\Leftrightarrow G$ fixes a point (up to finite index).

III Boundary and amenability

Th $G \curvearrowright X$ properly. Then $\exists \mu_n: \mathcal{E}ch(X) \rightarrow Prob(G)$
such that $\|\mu_n(gx) - g\mu_n(x)\| \rightarrow 0$.


Basic construction

$G = \mathbb{F}_2 \curvearrowright T$ Cayley graph.



$$\rho_n(\xi) = \frac{\mathbb{1}_{[0, x_m]}}{n}$$

$$\rho_n(gx) = \mathbb{1}_{[g_0, g x_m]}$$



$$\|\rho_n(g\xi) - g\rho_n(\xi)\| \leq \frac{2d(0, g_0)}{n} \rightarrow 0$$

① Choose a base point $o \in \text{ch}(x)$


We construct $\rho_n(\xi)$ with support in $Q(o, \xi)$

\rightarrow reduce the problem to the construction of ρ_n in an apartment.

② (Janukiewicz) W embeds equivariantly in a finite product of trees

in fact $\mathcal{E}_{\text{ch}}(\Sigma) \hookrightarrow \mathcal{E}_{\text{ch}}(T_1 \times \dots \times T_\ell) = \mathcal{E}_{\text{ch}}(T_1) \times \dots \times \mathcal{E}_{\text{ch}}(T_\ell)$.

Choose $W_0 \triangleleft W$ torsion free, finite index $\dots \times \mathcal{E}_{\text{ch}}(T_\ell)$.



③ Use the basic construction on each T_i $i \in \{1, \dots, \ell\}$ and take the product. Beware that $\text{Supp}(\rho_n(\xi))$ should be included in $Q(o, \xi)$.