

# Fractional Brownian motion satisfies two-way crossing

Rémi Peyre\* (CNRS & Université de Lorraine)

*Institut Élie Cartan de Lorraine*

*Campus Aiguillettes*

*BP 70239*

*54506 Vandœuvre-lès-Nancy Cedex (France)*

May 10, 2016

## Abstract

We prove the following result: For  $(Z_t)_{t \in \mathbf{R}}$  a fractional Brownian motion with arbitrary Hurst parameter, for any stopping time  $\tau$ , there exist arbitrarily small  $\varepsilon > 0$  such that  $Z_{\tau+\varepsilon} < Z_\tau$ , with asymptotic behaviour when  $\varepsilon \searrow 0$  satisfying a bound of iterated logarithm type. As a consequence, fractional Brownian motion satisfies the “two-way crossing” property, which has important applications in financial mathematics.

*Keywords:* fractional Brownian motion; stopping time; law of the iterated logarithm; two-way crossing.

## 1 Introduction

### 1.1 Context

In this article, we consider a filtered probability space  $(\Omega, \mathbb{P}, \mathcal{B}, (\mathcal{B}_t)_{t \in \mathbf{R}})$ . Notation ‘ $\omega$ ’ will implicitly refer to eventualities of  $\Omega$ ; we will use it from time to time when needing to make the dependency on the random phenomenon perfectly clear. We consider a (bilateral) Brownian motion  $(W_t)_{t \in \mathbf{R}}$  whose increments are adapted to our filtered space, which means, for all  $t \in \mathbf{R}$ , for all  $u \leq 0$ ,  $(W_{t+u} - W_t)$  is  $\mathcal{B}_t$ -measurable, while for all  $v \geq 0$ ,  $(W_{t+v} - W_t)$  is independent from  $\mathcal{B}_t$ .

We fix once for all some arbitrary parameter  $H \in (0, 1)$  (so that, in the sequel, “absolute” constants may actually depend on  $H$ ) such that  $H \neq 1/2$ ; moreover, in all this article,  $(H - 1/2)$  may be referred to as  $\eta$ . Then we consider the fractional Brownian motion (fBm)  $(Z_t)_{t \in \mathbf{R}}$  driven by  $W$  with Hurst parameter  $H$ , which means that

$$Z_t := C_1 \int_{\mathbf{R}} ((t-s)_+^\eta - (-s)_+^\eta) dW_s, \quad (1)$$

(with the convention that  $0^r = 0 \forall r \in \mathbf{R}$ ), where

$$C_1 := \left( \frac{1}{2H} + \int_0^\infty ((1+s)^\eta - s^\eta)^2 ds \right)^{-1/2}. \quad (2)$$

Then, the properties of  $Z$  are well known: it is a centred Gaussian process whose increments are adapted to  $(\mathcal{B}_t)_t$ , with  $\text{Var}(Z_t - Z_s) = |t - s|^{2H}$  and  $Z_0 = 0$  a.s.; its trajectories are locally  $(H - \varepsilon)$ -Hölder with divergence in  $O(|t|^{H+\varepsilon})$  at infinity, etc. [see e.g. Nourdin, 2012, Chap. 2].

*Remark 1.* The integral in the r.h.s. of (1) should not be seen as an actual Itô integral, but rather as a compact writing for some complicated deterministic integral, which corresponds to the “Itô” formulation via (formal) integrating by parts. Namely: (here in the case  $t > 0$ ),

$$\begin{aligned}
C_1^{-1}Z_t &= \int_{\mathbf{R}} ((t-s)_+^\eta - (-s)_+^\eta) dW_s \\
&= \int_{-\infty}^0 ((t-s)^\eta - (-s)^\eta) dW_s + \int_0^t (t-s)^\eta dW_s \\
&= \left[ ((t-s)^\eta - (-s)^\eta) W_s \right]_{s=-\infty}^0 - \eta \int_{-\infty}^0 ((t-s)^{\eta-1} - (-s)^{\eta-1}) W_s ds \\
&\quad + \left[ (t-s)^\eta (W_s - W_t) \right]_{s=0}^t - \eta \int_0^t (t-s)^{\eta-1} (W_s - W_t) ds \\
&= t^\eta W_t - \eta \int_{-\infty}^t ((t-s)^{\eta-1} - (-s)_+^{\eta-1}) (W_s - \mathbf{1}_{s>0} W_t) ds, \quad (3)
\end{aligned}$$

where all the computations are licit (with absolutely converging integrals) because of the properties of regularity and slow divergence of the (ordinary) Brownian motion.

*Remark 2.* It has to be stressed that in all this article, actually we are not interested in the values themselves of the processes  $W$  and  $Z$ , but rather in their *increments*. This way, the fact that  $W_0, Z_0 = 0$  should be considered as a mere convention, completely unessential though convenient.

## 1.2 Main result

The main result of this article is the following

**Theorem 3.** *In the context of § 1.1, for  $\tau$  any stopping time adapted to  $(\mathcal{B}_t)_t$ , one has almost-surely:*

$$\liminf_{v \searrow 0} \frac{Z_{\tau+v} - Z_\tau}{(\log|\log v|)^{1/2} v^H} \leq -C_1 H^{-1/2}. \quad (4)$$

*Remark 4.* What is the intuitive meaning of Theorem 3? Well, we know that fractional Brownian motion does not have Markov nor martingale property, which means that the future trajectory of the fBm after a stopping time may differ qualitatively from what it would be at a fixed time. Our theorem says however that that future trajectory cannot differ too wildly from its ordinary behaviour, in the sense that some weak form of the law of the iterated logarithm (LIL) always has to remain true.

My motivation for proving Theorem 3 was its application to financial mathematics, which will be explained in § 6; but the theorem may also be interesting even on its own right.

*Remark 5.* It is interesting to compare (4) with the standard LIL for the fBm, which states [*cf.* Arcones, 1995, Cor. 3.1] that in the case  $\tau \equiv 0$ :

$$\liminf_{v \searrow 0} \frac{Z_v - Z_0}{(\log|\log v|)^{1/2} v^H} = -\sqrt{2}. \quad (5)$$

Indeed, as soon as  $H \neq 1/2$ , one has  $C_1 H^{-1/2} < \sqrt{2}$ ; therefore, provided the r-h.s. of (4) is sharp, Theorem 3 does allow for a deviation from the standard LIL behaviour of the fBm after a stopping time—though not a too wild one, as already said. I conjecture that the r-h.s. of (4) is sharp indeed, and that it would be attained for  $\tau$  being a hitting time, from above for  $H \geq 1/2$ , resp. from below for  $H \leq 1/2$ —for instance, for  $H \geq 1/2$ , I believe that the following stopping time would fit equality in Equation (4):

$$\tau := \inf\{t \geq 0 \mid Z_t = Z_0 + 1\}. \quad (6)$$

*Remark 6.* As the increments of  $Z$  are adapted to the filtration  $(\mathcal{B}_t)_t$ , obviously in Theorem 3 we may replace that filtration by the filtration generated by the increments of  $Z$ .

*Remark 7.* In this article we are only considering the case  $H \neq 1/2$ , but Theorem 3 is trivially valid for  $H = 1/2$  too, since then the fBm  $Z$  is nothing but the ordinary Brownian motion  $W$  itself, for which the result follows immediately from the Markov property and the local properties of oBm.

### 1.3 Outline of the paper

§§ 2–5 will be devoted to proving Theorem 3. In § 2, we will see how one can get rid of the notion of stopping time to get Theorem 3 back to a result on the trajectories of the fractional Brownian motion. In § 3, we will make the needed result on fBm’s trajectories more precise, by establishing a kind of law of the iterated logarithm for some variant of the fBm. Next, an issue will be that we have to control the probability of an event being a union over a continuous infinity of  $t$ ’s: that issue will be handled by § 4, in which we will use regularity estimates on the fBm to get our continuous union back to a finite union. Finally, after all these simplifications it will only remain to prove some estimates on Gaussian vectors, which will be the work of § 5.

Some technical results will be postponed to appendices. In particular, in Appendix A we will compute the precise expression of the “drift operator” appearing in Lemma 15 describing what the law of the fBm becomes when you condition it by a stopping time: this formula, though not actually required to prove our main result, looks indeed intrinsically worthy to be written down to my eyes. Also, in Appendices B and C we will prove two lemmas on resp. the supremum of Gaussian processes and the inverse of nearly diagonal matrices.

Finally in § 6 we will explain how Theorem 3 can be applied to yield some new results in financial mathematics.

## 2 Conditional future of the fractional Brownian motion

### 2.1 Preliminary definitions

To begin with, it will be convenient to set some notation for certain sets of trajectories:

**Definition 8** (Sets  $\mathcal{P}$  and  $\mathcal{F}$ ).

- We denote by  $\mathcal{P}$  [like “past”] the set of the (deterministic) paths  $(X_u)_{u \leq 0}$  such that:

1.  $X_0 = 0$ ;
2.  $X$  is locally  $(H - \varepsilon)$ -Hölder for all  $\varepsilon > 0$ ;
3.  $X_u \stackrel{u \rightarrow -\infty}{\asymp} O(|u|^{H+\varepsilon})$  for all  $\varepsilon > 0$ .

- Similarly, we denote by  $\mathcal{F}$  [like “future”] the space of the paths  $(X_v)_{v \geq 0}$  satisfying the analogues of conditions 1–3 for non-negative times.

*Remark 9.* With the notation of Definition 8, one has almost-surely that, for all  $t \in \mathbf{R}$ ,  $(Z(\omega)_{t+u} - Z(\omega)_t)_{u \leq 0} \in \mathcal{P}$  and  $(Z(\omega)_{t+v} - Z(\omega)_t)_{v \geq 0} \in \mathcal{F}$ . [cf. Nourdin, 2012, Prop. 1.6 & Prop. 2.2.3].

We also define a certain “drift operator”:

**Definition 10** (“Drift operator”  $\mathbf{D}$ ). Let  $\mathbf{D}: \mathcal{P} \rightarrow \mathcal{F}$  be the linear operator such that, for  $X \in \mathcal{P}$ :

$$(\mathbf{D}X)_v := \int_{-\infty}^0 K(u, v) X_u du, \quad (7)$$

where

$$K(u, v) := \frac{\eta}{\Pi(\eta)\Pi(-\eta)} \times \left\{ \begin{aligned} &\eta \int_{-\infty}^0 (\mathbf{1}_{s>u} \xi_{\eta-1}(s-u, v) \xi_{-\eta-1}(-s, s-u) - \xi_{\eta-1}(-u, v) \xi_{-\eta-1}(-s, -u)) ds \\ &- v(v-u)^{\eta-1} (-u)^{-\eta-1} \end{aligned} \right\}, \quad (8)$$

where  $\Pi(\cdot)$  is Euler’s pi function extrapolating the factorial, and where we denote, for  $r \in \mathbf{R}$ ,  $a, b > 0$ :

$$\xi_r(a, b) := (a+b)^r - a^r. \quad (9)$$

*Remark 11.* Note that, since  $H \in (0, 1)$ , the integrals in (8) and (7) do converge (absolutely) indeed, and  $\mathbf{D}$  is well-defined on the whole  $\mathcal{P}$  with values in  $\mathcal{F}$ .

*Remark 12.* The equations (7)-(8) defining  $\mathbf{D}$ , though interesting as such, shall not play an essential role in this article. What is really important to have in mind is the moral *meaning* of this operator: actually  $\mathbf{D}$  was defined so that, informally,

$$(\mathbf{D}X)_v = \mathbb{E}(Z_v \mid (Z_u)_{u \leq 0} = (X_u)_{u \leq 0}). \quad (10)$$

The formal meaning of (10) will be made clear by Lemma 15 below.

Finally we define a process called the “Lévy fractional Brownian motion”, which is a kind of unilateral version of the “classic” fBm:

**Definition 13** (Lévy fBm). If  $(W_v)_{v \geq 0}$  is a (unilateral) ordinary Brownian motion, then the process  $(Y_v)_{v \geq 0}$  defined by

$$Y_v := C_1 \int_0^v (v-s)^\eta dW_s \quad (11)$$

(interpreted *via* the same integration by parts trick as in (3)) (and where we recall that  $C_1$  is defined by (2)) is called a *Lévy fractional Brownian motion* (with Hurst parameter  $H$ )—or, more accurately, the law of this process (which (11) defines without ambiguity) is called “the law of the Lévy fBm”.

*Remark 14.* From the regularity properties of the oBm, it is easy to check that the trajectories of the Lévy fBm lie in  $\mathcal{F}$  a.s..

## 2.2 Conditioning lemma

Now we can state the key lemma of this section:

**Lemma 15.** *In the context of § 1.1, for  $\tau$  a stopping time,  $((Z_{\tau+v} - Z_\tau)_{v \geq 0} - \mathbf{D}((Z_{\tau+u} - Z_\tau)_{u \leq 0}))$  is independent of  $\mathcal{B}_\tau$ , and its law is the Lévy fBm.*

*Remark 16.* In other words, Lemma 15 states that, conditionally to  $\mathcal{B}_\tau$  (or, morally, knowing the past trajectory of  $Z$  until  $\tau$ ), the law of the future trajectory of  $Z$  is equal to a “deterministic” drift term  $\mathbf{D}((Z_{\tau+u} - Z_\tau)_{u \leq 0})$  plus a “random” noise term being a Lévy fBm.

*Proof of Lemma 15.* As the increments of  $W$  are adapted to  $(\mathcal{B}_t)_t$ , conditionally to  $\mathcal{B}_\tau$ , the past trajectory  $(W_{\tau+u} - W_\tau)_{u \leq 0}$  of (the increments of)  $W$  is deterministic, while its future trajectory  $(W_{\tau+v} - W_\tau)_{v \geq 0}$  still has the unconditioned law of a standard oBm. Therefore, for  $t \geq 0$ , we split

$$\begin{aligned} Z_{\tau+t} - Z_\tau &\stackrel{(1)}{=} C_1 \int_{s \in \mathbf{R}} ((\tau + t - s)_+^\eta - (\tau - s)_+^\eta) dW_s \\ &= C_1 \int_{s \leftarrow s - \tau}^{s \leftarrow s - \tau} ((t - s)_+^\eta - (-s)_+^\eta) dW_{\tau+s} \\ &= C_1 \int_{u=-\infty}^0 ((t - u)^\eta - (-u)^\eta) dW_{\tau+u} + C_1 \int_{v=0}^t (t - v)^\eta dW_{\tau+v}, \end{aligned} \quad (12)$$

in which the first term is deterministic and given by some function of  $(W_{\tau+u} - W_\tau)_{u \leq 0}$ , while the second term (seen as a trajectory indexed by  $t$ ) has the law of the Lévy fBm indeed.

To end the proof, it remains to show that the aforementioned first term (seen as a trajectory indexed by  $t$ ) is equal to  $\mathbf{D}((Z_{\tau+u} - Z_\tau)_{u \leq 0})$  indeed. Since this point is actually not needed to prove our main result, we will postpone it to Appendix A.  $\square$

## 2.3 Reformulation of the main theorem

Thanks to Lemma 15, we will be able to get a sufficient condition for Theorem 3 in which there are no stopping times any more. But first we observe that proving Theorem 3 is tantamount to proving that for any  $\Lambda < C_1 H^{-1/2}$ , for any stopping time  $\tau$ :

$$\mathbb{P} \left( \liminf_{v \searrow 0} \frac{Z_{\tau+v} - Z_\tau}{(\log |\log v|)^{1/2} v^H} \geq -\Lambda \right) = 0. \quad (13)$$

When the event of the probability in Equation (13) holds true, we will say that  $\tau(\omega)$  is a  $\Lambda$ -slow time for the trajectory  $(Z(\omega)_t)_{t \in \mathbf{R}}$ .

• From now on, we fix an arbitrary  $\Lambda > -C_1 H^{-1/2}$  all along.

Now we need an *ad hoc* definition:

**Definition 17** (Nasty path). We say that a deterministic path  $(X_u)_{u \leq 0} \in \mathcal{P}$  is *nasty*, and we denote “ $X \in \mathcal{A}$ ”, when, for  $Y$  a Lévy fBm:

$$\mathbb{P}(0 \text{ is a } \Lambda\text{-slow time for } (\mathbf{D}X + Y(\omega))) > 0. \quad (14)$$

*Remark 18.*  $\mathcal{A}$  is a measurable subset of  $\mathcal{P}^{[1]}$ , as can be checked along the following lines:

- The drift operator  $\mathbf{D}: \mathcal{P} \rightarrow \mathcal{F}$  is measurable. (This is obvious from its characterization (7) as an explicit kernel operator).
- Denoting  $\llbracket n \rrbracket := \{0, \dots, n-1\}$  and  $\vec{X}_I := (X_i)_{i \in I}$ , for all  $n \in \mathbf{N}$ , for all  $\vec{v}_{\llbracket n \rrbracket} \in (\mathbf{R}_+)^{\llbracket n \rrbracket}$ ,  $\mathbb{P}(\vec{Y}_{\vec{v}_{\llbracket n \rrbracket}} \in \prod_{i \in \llbracket n \rrbracket} [a_i, b_i])$  is a measurable function of  $(\vec{a}_{\llbracket n \rrbracket}, \vec{b}_{\llbracket n \rrbracket})$ . (This is because  $\vec{Y}_{\vec{v}_{\llbracket n \rrbracket}}$  is a Gaussian random vector with known parameters).

- Therefore, for all  $n \in \mathbf{N}$ , for all  $\vec{v}_{\llbracket n \rrbracket} \in (\mathbf{R}_+)^{\llbracket n \rrbracket}$ , for all  $\vec{a}_{\llbracket n \rrbracket}, \vec{b}_{\llbracket n \rrbracket} \in \mathbf{R}^{\llbracket n \rrbracket}$ , the probabilities

$$\mathbb{P}(\forall i \in \llbracket n \rrbracket \quad (\mathbf{D}X + Y(\omega))_{v_i} \in [a_i, b_i]) \quad (15)$$

are measurable functions of  $X$ ; by Dynkin's  $\pi$ - $\lambda$  theorem, it follows that for all measurable  $A \subset \mathcal{F}$ ,  $\mathbb{P}(\mathbf{D}X + Y(\omega) \in A)$  is a measurable function of  $X$ .

- But “0’s being a  $\Lambda$ -slow time” is a measurable subset of  $\mathcal{F}$  (because the trajectories of  $\mathcal{F}$  are continuous, so they can be described fully from a countable set of time indices), so the probability in the l-h.s. of (14) is a measurable function of  $X$ , which finally implies that  $\mathcal{A}$  is measurable.

That vocabulary being set, Equation (13), and hence Theorem 3, will be a consequence of the following

**Proposition 19.** *In the context of § 1.1:*

$$\mathbb{P}(\exists t \in \mathbf{R} \quad (Z(\omega)_{t+u} - Z(\omega)_t)_{u \leq 0} \in \mathcal{A}) = 0. \quad (16)$$

*Proof of Equation (13) from Proposition 19.* Let  $\tau$  be any stopping time. We introduce the notation  $\mathbb{P}'$  for the law of the Lévy fBm  $Y$ , and the probability  $\mathbb{P}'_{\tau, \omega}$  defined as the pushforward measure of  $\mathbb{P}'$  by the map  $(Y_v)_{v \geq 0} \mapsto (X_t)_{t \in \mathbf{R}}$  from  $\mathcal{F}$  to  $\mathcal{C}(\mathbf{R}, \mathbf{R})$  defined by:

$$X_t = \begin{cases} Z_t & t \leq \tau(\omega); \\ Z_{\tau(\omega)} + \mathbf{D}((Z(\omega)_{\tau(\omega)+u} - Z(\omega)_{\tau(\omega)})_{u \leq 0}) + Y_{t-\tau(\omega)} & t > \tau(\omega), \end{cases} \quad (17)$$

so that, by Lemma 15,  $(\mathbb{P}'_{\tau, \omega})_{\omega \in \Omega}$  is the conditional probability of  $\mathbb{P}$  given  $\mathcal{B}_\tau$ . (That conditional probability is regular, by the same arguments as in Remark 18). Then, writing the law of total probability w.r.t.  $\mathcal{B}_\tau$ :<sup>[2]</sup>

$$\begin{aligned} & \mathbb{P}(\tau \text{ is a } \Lambda\text{-slow time for } Z) \\ &= \mathbb{E}(\mathbb{P}'_{\tau, \omega}(\tau(\omega) \text{ is } \Lambda\text{-slow time for } Z(\omega'))) \\ &= \mathbb{E}(\mathbb{P}'(0 \text{ is a } \Lambda\text{-slow time for } (\mathbf{D}(Z(\omega)_{\tau(\omega)+u} - Z(\omega)_{\tau(\omega)})_{u \leq 0} + Y(\omega')))) \\ &= \mathbb{E}(0 \text{ whenever } (Z_{\tau+u} - Z_\tau)_{u \leq 0} \notin \mathcal{A}), \end{aligned} \quad (18)$$

which is zero by Proposition 19.  $\square$

So, in the sequel, our new goal will be to prove Proposition 19.

<sup>[1]</sup>The  $\sigma$ -algebra considered on  $\mathcal{P}$  (resp. on  $\mathcal{F}$ ) is obviously (the trace of) the product  $\sigma$ -algebra—a trajectory in  $\mathcal{P}$  (resp.  $\mathcal{F}$ ) being seen as a function from  $\mathbf{R}_-$  (resp.  $\mathbf{R}_+$ ) to  $\mathbf{R}$ .

<sup>[2]</sup>In the following computation, to avoid confusions, I denoted by ‘ $\omega$ ’ the variable bound to operator  $\mathbb{E}$ , and by ‘ $\omega'$ ’ the variable bound to probabilities  $\mathbb{P}'$  and  $\mathbb{P}'_{\tau, \omega}$ .

### 3 Local behaviour of fBm's trajectories

#### 3.1 A law of the iterated logarithm for the Lévy fBm

• In all this article, we denote  $\llbracket n \rrbracket := \mathbf{N} \cap [0, n) = \{0, 1, 2, \dots, n-1\}$ . A subset  $\mathcal{I} \subset \mathbf{N}$  will be said to be thick when it has positive upper asymptotic density:

$$\limsup_{n \rightarrow \infty} \frac{|\mathcal{I} \cap \llbracket n \rrbracket|}{n} > 0. \quad (19)$$

The first main result of this section is the following

**Lemma 20.** *Let  $r \in (0, 1)$  and let  $\mathcal{I}$  be a thick subset of  $\mathbf{N}$ ; then, for  $Y$  a Lévy fBm:*

$$\liminf_{\substack{i \in \mathcal{I} \\ i \rightarrow \infty}} \frac{Y_{r^i}}{(\log i)^{1/2} r^{Hi}} \leq -C_1 H^{-1/2} \quad a.s.. \quad (20)$$

*Remark 21.* From Lemma 20 one can deduce immediately that  $\liminf_{v \searrow 0} \{Y_v / (\log|\log v|)^{1/2} v^H\} \leq -C_1 H^{-1/2}$ , which is the classic formulation of (one sense of) a law of the iterated logarithm, hence the title of this subsection. Note by the way that one could also prove that there is actually equality in (20) as well as in the LIL derived from it; but that will not be needed here.

*Proof of Lemma 20.* It will be convenient in this proof to assume that  $Y$  is driven by some oBm  $W$  according to (11). Then, for  $v \geq 0$ , let us define

$$\tilde{Y}_v := C_1 \int_{rv}^v (v-s)^\eta dW_s, \quad (21)$$

resp.

$$Y'_v := Y_v - \tilde{Y}_v = C_1 \int_0^{rv} (v-s)^\eta dW_s. \quad (22)$$

First let us study the  $\tilde{Y}_{r^i}$ 's. Obviously these random variables are independent, with  $Y_{r^i} / r^{Hi} \sim \mathcal{N}(0, C_1^2(1-r)^{2H}/2H) \forall i$ . Now, using that  $\mathbb{P}(\mathcal{N}(0, 1) \leq -x) \geq e^{-x^2/2}/2\sqrt{2\pi}x$  for  $x \geq 1$ ,<sup>[3]</sup> we get that for  $i$  large enough: (having fixed some arbitrary small  $\varepsilon \in (0, 1)$ ),

$$\begin{aligned} \mathbb{P}(\tilde{Y}_{r^i} / (\log i)^{1/2} r^{Hi} \leq -(1-\varepsilon)C_1(1-r)^H H^{-1/2}) \\ = \mathbb{P}(\mathcal{N}(0, 1) \leq -(1-\varepsilon)\sqrt{2}(\log i)^{1/2}) \\ \geq i^{-(1-\varepsilon)^2} / (1-\varepsilon)4\sqrt{\pi}(\log i)^{1/2} \stackrel{i \rightarrow \infty}{=} \Omega(i^{-1}), \end{aligned} \quad (23)$$

where “ $f(i) = \Omega(g(i))$ ” means that  $g(i) = O(f(i))$ . As  $\mathcal{I}$  is thick, the series  $\sum_{i \in \mathcal{I}} i^{-1}$  is divergent, thus so is

$$\sum_{i \in \mathcal{I}} \mathbb{P}\left(\frac{\tilde{Y}_{r^i}}{(\log i)^{1/2} r^{Hi}} \leq -(1-\varepsilon)C_1(1-r)^H H^{-1/2}\right). \quad (24)$$

<sup>[3]</sup>This is because of convexity of the density  $y \mapsto \varphi(y) := e^{-y^2/2} / \sqrt{2\pi}$  on  $(-\infty, -1]$ : from this property you deduce that  $\int_{-\infty}^{-x} \varphi(y) dy \geq \varphi(-x)^2 / 2\varphi'(-x) = \varphi(-x)/2x$ .

Since the events concerning the different  $\tilde{Y}_{r^i}$ 's are independent, it follows by the (second) Borel–Cantelli lemma that almost-surely there are infinitely many  $i \in \mathcal{I}$  for which  $\tilde{Y}_{r^i} / (\log i)^{1/2} r^{Hi} \leq -(1 - \varepsilon) C_1 (1 - r)^H H^{-1/2}$ , so that:

$$\liminf_{\substack{i \in \mathcal{I} \\ i \rightarrow \infty}} \frac{\tilde{Y}_{r^i}}{(\log i)^{1/2} r^{Hi}} \leq -(1 - \varepsilon) C_1 (1 - r)^H H^{-1/2} \quad \text{a.s.}, \quad (25)$$

in which the factor  $(1 - \varepsilon)$  can be removed by letting  $\varepsilon \rightarrow 0$ .

Now let us handle the  $Y'_{r^i}$ 's. One has  $Y'_{r^i} / r^{Hi} \sim \mathcal{N}(0, C_1^2 (1 - (1 - r)^{2H}) / 2H)$ ; therefore, using that  $\mathbb{P}(\mathcal{N}(0, 1) \geq x) \leq e^{-x^2/2}$  for all  $x$ ,<sup>[4]</sup> we get that: (having fixed some arbitrary small  $\varepsilon > 0$ ),

$$\begin{aligned} \mathbb{P}(Y'_{r^i} / (\log i)^{1/2} r^{Hi} \geq (1 + \varepsilon) C_1 (1 - (1 - r)^{2H})^{1/2} H^{-1/2}) \\ = \mathbb{P}(\mathcal{N}(0, 1) \geq (1 + \varepsilon) \sqrt{2} (\log i)^{1/2}) \leq i^{-(1+\varepsilon)^2}. \end{aligned} \quad (26)$$

The series  $\sum_{i \in \mathcal{I}} i^{-(1+\varepsilon)^2}$  is convergent since  $\sum_{i \in \mathbf{N}} i^{-(1+\varepsilon)^2}$  is, thus so is

$$\sum_{i \in \mathcal{I}} \mathbb{P}\left(\frac{Y'_{r^i}}{(\log i)^{1/2} r^{Hi}} \geq (1 + \varepsilon) C_1 (1 - (1 - r)^{2H})^{1/2} H^{-1/2}\right). \quad (27)$$

It follows by the (first) Borel–Cantelli lemma that almost-surely there are only finitely many  $i$ 's for which  $\tilde{Y}_{r^i} / (\log i)^{1/2} r^{Hi} \geq (1 + \varepsilon) C_1 (1 - (1 - r)^{2H})^{1/2} H^{-1/2}$ , so that:

$$\limsup_{\substack{i \in \mathcal{I} \\ i \rightarrow \infty}} \frac{Y'_{r^i}}{(\log i)^{1/2} r^{Hi}} \leq (1 + \varepsilon) C_1 (1 - (1 - r)^{2H})^{1/2} H^{-1/2} \quad \text{a.s.}, \quad (28)$$

in which the factor  $(1 + \varepsilon)$  can be removed by letting  $\varepsilon \rightarrow 0$ .

Summing (25) and (28), we get an intermediate result:

**Proposition 22.** *Under the assumptions of Lemma 20, almost-surely:*

$$\liminf_{i \in \mathcal{I}} \frac{Y_{r^i}}{(\log i)^{1/2} r^{Hi}} \leq -\lambda(r), \quad (29)$$

where

$$\lambda(r) := C_1 \left( (1 - r)^H - (1 - (1 - r)^{2H})^{1/2} \right) H^{-1/2}. \quad (30)$$

So, now it remains to improve the constant  $\lambda(r)$  in (29) into  $C_1 H^{-1/2}$ . For this, we begin with observing that the Lévy fBm is scale-invariant with exponent  $H$  (by which I mean that for  $a \in \mathbf{R}_+^*$ ,  $(Y_{av} / a^H)_{v \geq 0}$  is also a Lévy fBm); therefore, Proposition 22 has the following

**Corollary 23.** *Under the assumptions of Lemma 20, for  $a \in \mathbf{R}_+^*$ , one has almost-surely:*

$$\liminf_{i \in \mathcal{I}} \frac{Y_{ar^i}}{a^H (\log i)^{1/2} r^{Hi}} \leq -\lambda(r). \quad (31)$$

---

<sup>[4]</sup>This is because  $\mathbb{E}(e^{x\mathcal{N}(0,1)}) = e^{x^2/2}$ , from which the claimed formula follows by Markov's inequality.

Now let  $k > 1$  be an arbitrary large integer; and take  $l \in \llbracket k \rrbracket$  such that  $\mathcal{J} := \{j \in \mathbf{N} \mid kj + l \in \mathcal{I}\}$  is thick—such an  $l$  exists since  $\mathcal{I}$  itself is thick. Then one has:

$$\begin{aligned} \liminf_{i \in \mathcal{I}} \frac{Y_{r^i}}{(\log i)^{1/2} r^{Hi}} &\leq \liminf_{j \in \mathcal{J}} \frac{Y_{r^{kj+l}}}{(\log(kj+l))^{1/2} r^{H(kj+l)}} \\ &= \liminf_{j \in \mathcal{J}} \frac{Y_{r^{l(r^k)j}}}{(r^l)^H (\log j)^{1/2} (r^k)^{Hj}}, \end{aligned} \quad (32)$$

where in the last equality we used that  $(\log(kj+l))^{1/2} \stackrel{j \rightarrow \infty}{\sim} (\log j)^{1/2}$ . But, applying Corollary 23 with ‘ $r$ ’ =  $r^k$ , ‘ $a$ ’ =  $r^l$  and ‘ $\mathcal{I}$ ’ =  $\mathcal{J}$ , the r.h.s. of (32) is bounded above by  $-\lambda(r^k)$ ; so,

$$\liminf_{i \in \mathcal{I}} \frac{Y_{r^i}}{(\log i)^{1/2} r^{Hi}} \leq -\lambda(r^k). \quad (33)$$

Letting  $k$  tend to infinity,  $\lambda(r^k)$  tends to  $C_1 H^{-1/2}$ , which finally proves (20).  $\square$

### 3.2 Nastiness condition as a limit

For all the sequel of this article, we fix some  $r \in (0, 1)$  small enough (in a sense to be made precise later); we also fix arbitrarily two parameters  $\alpha \in (0, C_1 H^{-1/2} - \Lambda)$  and  $p \in (0, 1)$ . Then we define, for all  $n > 0$ :

$$\mathcal{A}_n := \{(X_u)_{u \leq 0} \in \mathcal{P} \mid \text{card}\{i \in \llbracket n \rrbracket \mid (\mathbf{D}X)_{r^i} \geq \alpha (\log i)_+^{1/2} r^{Hi}\} \geq pn\}. \quad (34)$$

Then we have the following connection between  $\mathcal{A}$  and the  $\mathcal{A}_n$ ’s:

**Lemma 24.**

$$\mathcal{A} \subset \liminf_{n \in \mathbf{N}} \mathcal{A}_n. \quad (35)$$

*Proof.* We prove the contrapositive inclusion. Let  $(X_u)_{u \leq 0} \in \mathcal{P}$  be such that  $X \notin \liminf \{\mathcal{A}_n\}$ , that is, the set  $\{n \mid X \notin \mathcal{A}_n\}$  is unbounded; and set

$$\mathcal{I} := \{i \in \mathbf{N} \mid (\mathbf{D}X)_{r^i} < \alpha (\log i)_+^{1/2} r^{Hi}\}. \quad (36)$$

One has by definition that  $|\mathcal{I} \cap \llbracket n \rrbracket| / n \geq 1 - p$  for all  $n$  such that  $X \notin \mathcal{A}_n$ ; as these  $n$  are unbounded and  $1 - p > 0$ , it follows that  $\mathcal{I}$  is thick. Therefore, Lemma 20 gives that for almost-all Lévy fBm  $Y(\omega)$ , one has that

$$\liminf_{i \in \mathcal{I}} \frac{Y(\omega)_{r^i}}{(\log i)^{1/2} r^{Hi}} \leq -C_1 H^{-1/2}. \quad (37)$$

On the other hand, the definition of  $\mathcal{I}$  obviously implies that

$$\liminf_{i \in \mathcal{I}} \frac{(\mathbf{D}X)_{r^i}}{r^{Hi} (\log i)^{1/2}} \leq \alpha. \quad (38)$$

Summing (37) and (38), it follows that almost-surely:

$$\liminf_{i \in \mathcal{I}} \frac{(\mathbf{D}X + Y(\omega))_{r^i}}{(\log i)^{1/2} r^{Hi}} \leq \alpha - C_1 H^{-1/2} < -\Lambda, \quad (39)$$

so that  $X \notin \mathcal{A}$ .  $\square$

### 3.3 Second reformulation of the main theorem

Thanks to the work of this section, we are now able to show that the following result will be a sufficient condition for Proposition 19:

**Proposition 25.** *In the context of § 1.1:*

$$\mathbb{P}(\exists t \in [0, 1] \quad (Z(\omega)_{t+u} - Z(\omega)_t)_{u \leq 0} \in \mathcal{A}_n) \xrightarrow{n \rightarrow \infty} 0. \quad (40)$$

*Proof of Proposition 19 from Proposition 25.* Proposition 24 implies that

$$\begin{aligned} & \{\omega \in \Omega \mid \exists t \in [0, 1] \quad (Z(\omega)_{t+u} - Z(\omega)_t)_{u \leq 0} \in \mathcal{A}\} \\ & \subset \liminf_{n \rightarrow \infty} \{\omega \mid \exists t \in [0, 1] \quad (Z(\omega)_{t+u} - Z(\omega)_t)_{u \leq 0} \in \mathcal{A}_n\}; \end{aligned} \quad (41)$$

therefore, by the (first) Borel–Cantelli lemma, Proposition 25 yields that

$$\mathbb{P}(\exists t \in [0, 1] \quad (Z(\omega)_{t+u} - Z(\omega)_t)_{u \leq 0} \in \mathcal{A}) = 0. \quad (42)$$

But, since the increments of the fractional Brownian motion are stationary, in (42) we may replace  $[0, 1]$  by  $[n, n + 1]$  for all  $n \in \mathbf{Z}$ ; and then, by countable union, we get the wished result (16).  $\square$

So, in the sequel, our new goal will be to prove Proposition 25.

## 4 Pathwise control via pointwise control

### 4.1 A regularity result

One of our issues to prove Proposition 25 is that we have to bound the probability of an event defined as a union for uncountably infinitely many  $t$ 's. To overcome this issue, we will need a tool to “get rid of the trajectorial aspects” of the problem: this is the work of this section.

First, we need a little notation:

**Definition 26** (Processes  $\hat{\Gamma}_i$  and variables  $\Gamma_i$ ). Within the context of § 1.1, for  $i \in \mathbf{N}$ , we define the following random process (indexed by  $t \in \mathbf{R}$ ):

$$\hat{\Gamma}_i(\omega)_t := \frac{\mathbf{D}((Z(\omega)_{t+u} - Z(\omega)_t)_{u \leq 0})_{r^i}}{r^{Hi}}. \quad (43)$$

We also define the following random variable:

$$\Gamma_i(\omega) := \hat{\Gamma}_i(\omega)_0 = \frac{\mathbf{D}((Z(\omega)_u)_{u \leq 0})_{r^i}}{r^{Hi}}. \quad (44)$$

Then, the main result of this section is the following

**Lemma 27.** *In the context of this section, there exist absolute<sup>[5]</sup> constants  $C_a > 0$ ,  $C_b < \infty$  (whose exact expressions do not matter) such that for all  $i \in \mathbf{N}$ , for all  $T > 0$ :*

$$\mathbb{P}(\exists t \in [0, T] \quad |(\hat{\Gamma}_i)_t - \Gamma_i| \geq 1) \leq C_b \exp(-C_a(r^i / T)^{2H \wedge 1}). \quad (45)$$

<sup>[5]</sup>Remember that in this article, “absolute” constants may actually depend on  $H$ .

*Proof.* First, since  $Z$  is scale-invariant with exponent  $H$  (and operator  $\mathbf{D}$  preserves that scale invariance), it will be enough to prove Lemma 27 for  $i = 0$ ; so we will only handle that case. Then the subscript  $i$  becomes useless, so we remove it in our notation.

Because of the characterization (10) of  $\mathbf{D}$ ,  $\hat{\Gamma}_t$  may be written as a function of  $W$ :

$$\hat{\Gamma}_t = C_1 \int_{-\infty}^t ((t+1-s)^\eta - (t-s)^\eta) dW_s. \quad (46)$$

That shows that  $\hat{\Gamma}$  is a stationary centred Gaussian random process, with

$$\begin{aligned} \text{Var}(\hat{\Gamma}_t - \hat{\Gamma}_0) &= C_1^2 \int_{\mathbf{R}} (\mathbf{1}_{s \leq t} (t+1-s)^\eta - \mathbf{1}_{s \leq 0} (1-s)^\eta - (t-s)_+^\eta + (-s)_+^\eta)^2 ds \\ &\stackrel{t \rightarrow 0}{\asymp} O(t^{2H \wedge 1}). \end{aligned} \quad (47)$$

To go further, we need the following lemma, whose proof is postponed to Appendix B:

**Lemma 28.** *Let  $(X_t)_{t \in [0,1]}$  be a centred Gaussian process such that  $X_0 = 0$  a.s. and*

$$\forall t, s \in [0, 1] \quad \text{Var}(X_t - X_s) \leq |t - s|^{2\theta} \quad (48)$$

for some  $\theta \in (0, 1]$ . In such a case it is known that  $X$  has a continuous version (by the Kolmogorov theorem) and that, for this continuous version, the random variable  $\|X\|(\omega) := \sup_{t \in [0,1]} |X(\omega)_t|$  is sub-Gaussian (by the Fernique theorem), i.e. there exist constants  $C_c > 0, C_d < \infty$  such that

$$\forall x \geq 0 \quad \mathbb{P}(\|X\| \geq x) \leq C_d \exp(-C_c x^2). \quad (49)$$

The present lemma states that, moreover, the constants  $C_c$  and  $C_d$  can be made explicit, only depending on  $\theta$ .

We apply Lemma 28 in the following way. From (47), one has  $\text{Var}(\hat{\Gamma}_t - \hat{\Gamma}_s) \leq C_e |t - s|^{2H \wedge 1}$  for all  $t, s \in [0, 1]$ , for some  $C_e < \infty$ . Therefore, for  $T \leq 1$ , the random process

$$X(\omega)_t := (C_e T^{2H \wedge 1})^{-1/2} (\hat{\Gamma}(\omega)_{tT} - \Gamma(\omega)) \quad (50)$$

satisfies the assumptions of Lemma 28 with ' $\theta$ ' =  $H \wedge 1/2$ , so that (49) yields:

$$\mathbb{P}(\exists t \in [0, T] \quad |(\hat{\Gamma}_i)_t - \Gamma_i| \geq x) \leq C_d \exp(-C_c C_e^{-1} x^2 / T^{2H \wedge 1}). \quad (51)$$

This implies (45) for  $T \leq 1$ , with constants not depending on  $T$ . On the other hand, up to replacing  $C_b$  by  $(e^{C_a} \vee C_b)$ , (45) is automatically true for  $T > 1$ ; so the proof of Lemma 27 is completed.  $\square$

## 4.2 Third reformulation of the main result

Now, we will see how Proposition 27 allows one to find an easier sufficient condition for Proposition 25. First of all, we have to introduce a little notation: in this section, we fix some arbitrary  $\alpha' \in (0, \alpha)$ ,  $p' \in (0, p)$ , and we define  $\mathcal{A}'_n$  by the variant of Equation (34) in which  $\alpha$  and  $p$  are replaced by resp.  $\alpha'$  and  $p'$ ; also, we fix some arbitrary  $\tilde{r} \in (0, r)$ , and we set

$$T_n := \tilde{r}^n. \quad (52)$$

Now, we introduce the following events of  $\Omega$ :

**Definition 29** (Events  $A_n, A'_n, \bar{A}_n, \bar{A}_n^k$  and  $\bar{A}_n^*$ ).

$$A_n := \{\omega \mid (Z(\omega)_u)_{u \leq 0} \in \mathcal{A}_n\}; \quad (53)$$

$$A'_n := \{\omega \mid (Z(\omega)_u)_{u \leq 0} \in \mathcal{A}'_n\}; \quad (54)$$

$$\bar{A}_n := \{\omega \mid \exists t \in [0, T_n] \quad (Z(\omega)_{t+u} - Z(\omega)_t)_{u \leq 0} \in \mathcal{A}_n\}; \quad (55)$$

$$\bar{A}_n^k := \{\omega \mid \exists t \in [kT_n, (k+1)T_n] \quad (Z(\omega)_{t+u} - Z(\omega)_t)_{u \leq 0} \in \mathcal{A}_n\}; \quad (56)$$

$$\bar{A}_n^* := \{\omega \mid \exists t \in [0, 1] \quad (Z(\omega)_{t+u} - Z(\omega)_t)_{u \leq 0} \in \mathcal{A}_n\}. \quad (57)$$

Then we claim that

**Lemma 30.** *For  $n$  large enough,*

$$\omega \in \bar{A}_n \setminus A'_n \quad \Rightarrow \quad \exists i \in \llbracket n \llbracket \quad \exists t \in [0, T_n] \quad |\hat{\Gamma}_i(\omega)_t - \Gamma_i(\omega)| \geq 1. \quad (58)$$

*Proof.* Assume that  $\omega \in \bar{A}_n \setminus A'_n$ . Then the fact that  $\omega \in \bar{A}_n$  means that there exists some  $t \in [0, T_n]$  such that  $(Z_{t+u} - Z_t)_{u \leq 0} \in \mathcal{A}_n$ . For such a  $t$ , going back to the definitions (34) and (43) of  $\mathcal{A}_n$  and  $\hat{\Gamma}_i$ , this means that

$$\text{card}\{i \in \llbracket n \llbracket \mid (\hat{\Gamma}_i)_t \geq \alpha(\log i)_+^{1/2}\} \geq pn. \quad (59)$$

Similarly, the fact that  $\omega \notin A'_n$  means that

$$\text{card}\{i \in \llbracket n \llbracket \mid \Gamma_i \geq \alpha'(\log i)_+^{1/2}\} < p'n. \quad (60)$$

Therefore, there exist at least  $(p-p')n$  indices ' $i$ ' such that  $(\hat{\Gamma}_i)_t \geq \alpha(\log i)_+^{1/2}$  while (for the same  $i$ )  $\Gamma_i < \alpha'(\log i)_+^{1/2}$ . Necessarily one these indices is  $\geq (p-p')n - 1$ ; thus, for such an  $i$ , one has:

$$(\hat{\Gamma}_i)_t - \Gamma_i \geq (\alpha - \alpha')(\log((p-p')n - 1))^{1/2}. \quad (61)$$

But, provided  $n \geq (e^{(\alpha-\alpha')^{-2}} + 1) / (p-p')$ , the r.h.s. of (61) is  $\geq 1$ ; so in the end we have found  $i \in \llbracket n \llbracket, t \in [0, T_n]$  such that  $|\hat{\Gamma}_i(\omega)_t - \Gamma_i(\omega)| \geq 1$ , proving the lemma.  $\square$

Combining Lemma 30 with Lemma 27, we get that

$$\mathbb{P}(\bar{A}_n \setminus A'_n) \leq \sum_{i=0}^{n-1} C_b \exp(-C_a(r^i / T_n)^{2H \wedge 1}), \quad (62)$$

in which the right-hand side is obviously bounded by

$$nC_b \exp(-C_a(r / \tilde{r})^{(2H \wedge 1)n}), \quad (63)$$

which shows that  $\mathbb{P}(\bar{A}_n \setminus A'_n)$  decreases superexponentially in  $n$  (i.e. faster than any exponential).

Now  $\bar{A}_n^* \subset \bigcup_{k \in \llbracket [1/T_n] \llbracket} \bar{A}_n^k$ , where  $\mathbb{P}(\bar{A}_n^k) = \mathbb{P}(\bar{A}_n) \forall k$  by translation invariance, so it follows that

$$\mathbb{P}(\bar{A}_n^*) \leq [1/T_n] \mathbb{P}(\bar{A}_n) \leq [\tilde{r}^{-n}] (\mathbb{P}(A'_n) + \mathbb{P}(\bar{A}_n \setminus A'_n)) = [\tilde{r}^{-n}] \mathbb{P}(A'_n) + o(1). \quad (64)$$

Our goal being to prove that  $\mathbb{P}(\bar{A}_n^*) \rightarrow 0$  as  $n \rightarrow \infty$  (that is just re-writing Proposition 25 with the notation of this section), it will be sufficient for that to prove the following

**Proposition 31.**  $\mathbb{P}(A'_n)$  decreases superexponentially in  $n$ .

So, as  $A'_n$  corresponds to a condition on a finite-dimensional Gaussian vector, we have managed to get completely rid of the trajectorial aspects of the problem! Now our ultimate goal will be to prove Proposition 31.

*Remark 32.* As the “prime” symbols would be somehow cumbersome, we will drop them in the sequel, thus actually proving the superexponential decrease of  $\mathbb{P}(A_n)$ . Nevertheless this should not be confusing, as the constraints on  $\alpha$  and  $p$  (and therefore on  $A_n$ ) are the same as on  $\alpha'$  and  $p'$  (and therefore on  $A'_n$ ).

## 5 Final computations: controlling a Gaussian vector

### 5.1 Covariance structure

☛ In this section, for ‘ $X$ ’ a symbol and  $I$  a discrete set, “ $\vec{X}_I$ ” will be a shorthand for “ $(X_i)_{i \in I}$ ”.

So, our goal is to prove the superexponential decay of  $\mathbb{P}(A_n)$ , which can be re-written as

$$\mathbb{P}(A_n) = \mathbb{P}(\text{card}\{i \in \llbracket n \rrbracket \mid \Gamma_i \geq \alpha(\log i)_+^{1/2}\} \geq pn) \quad (65)$$

(where  $\Gamma_i$  was defined by (44)). (65) obviously implies that

$$\mathbb{P}(A_n) \leq \sum_{\substack{I \subset \llbracket n \rrbracket \\ |I| \geq pn}} \mathbb{P}(\forall i \in I \quad \Gamma_i \geq \alpha(\log i)_+^{1/2}). \quad (66)$$

As there are only  $2^n$  subsets of  $\llbracket n \rrbracket$ , to prove that  $\mathbb{P}(A_n)$  decreases superexponentially it is therefore sufficient to prove that

$$\sup_{\substack{I \subset \llbracket n \rrbracket \\ |I| \geq pn}} \mathbb{P}(\forall i \in I \quad \Gamma_i \geq \alpha(\log i)_+^{1/2}) \quad (67)$$

decreases superexponentially.

Now, by (10) one has

$$\Gamma_i(\omega) = C_1 r^{-Hi} \int_{-\infty}^0 ((r^i - s)^\eta - (-s)^\eta) dW(\omega)_s; \quad (68)$$

therefore  $\vec{\Gamma}_{\mathbf{N}}$  is a centred Gaussian vector, with:

$$\begin{aligned} \text{Cov}(\Gamma_i, \Gamma_j) &= C_1^2 r^{-(i+j)H} \int_{-\infty}^0 ((r^i - s)^\eta - (-s)^\eta)((r^j - s)^\eta - (-s)^\eta) ds \\ &= C_1^2 r^{-|i-j|H} \int_{-\infty}^0 ((r^{|i-j|} - s)^\eta - (-s)^\eta)((1 - s)^\eta - (-s)^\eta) ds \\ &\leq C_f r^{(1/2-|\eta||i-j|)}, \end{aligned} \quad (69)$$

for some absolute constant  $C_f < \infty$ . Therefore, provided  $r$  was chosen small enough, we have the following control on the covariance matrix of  $\vec{\Gamma}_{\mathbf{N}}$ :

$$\text{Cov}(\Gamma_i, \Gamma_j) = \sigma^2 \quad \text{for } i = j; \quad (70)$$

$$|\text{Cov}(\Gamma_i, \Gamma_j)| \leq \sigma^2 \varepsilon^{|i-j|} \quad \text{for } i \neq j, \quad (71)$$

where  $\varepsilon > 0$  is some small parameter which will be fixed later, and where  $\sigma := \text{Var}(\Gamma)^{1/2} > 0$  (since  $H \neq 1/2$ ).

## 5.2 Density estimates

To exploit (70)-(71), we need the following lemma (whose proof is postponed to Appendix C):

**Lemma 33.** *For  $n \in \mathbf{N}$ ,  $\varepsilon > 0$ , let  $\mathbf{A} =: ((a_{ij}))_{i,j \in [n]}$  be a square matrix such that  $a_{ii} = 1 \forall i$  and  $|a_{ij}| \leq \varepsilon^{|i-j|} \forall i \neq j$ . Then:*

$$\det \mathbf{A} \geq \exp(-n\Phi_g(\varepsilon)\varepsilon^2), \quad (72)$$

where  $\Phi_g: (0, \infty) \rightarrow [1, \infty]$  is some absolute function (in particular, not depending on  $n$ ) such that  $\Phi_g(\varepsilon) \xrightarrow{\varepsilon \rightarrow 0} 1$ —we will call such a function a quasi-one function. In particular, provided  $\varepsilon$  is small enough,  $\mathbf{A}$  is invertible. The present lemma asserts moreover that, then, denoting  $\mathbf{A}^{-1} =: ((b_{ij}))_{i,j}$ :

$$|b_{ij}| \leq 2^{|i-j|-1}(\Phi_h(\varepsilon)\varepsilon)^{|i-j|} \quad \forall i \neq j; \quad (73)$$

$$|b_{ii} - 1| \leq 2\Phi_i(\varepsilon)\varepsilon^2 \quad \forall i, \quad (74)$$

for  $\Phi_i$  and  $\Phi_h$  some other absolute “quasi-one functions”.

We apply Lemma 33 to the covariance matrix of  $\vec{\Gamma}_{[n]}$  (assuming  $\varepsilon$  was chosen small enough so that  $\Phi_g$  is finite); then, the formula for the density of Gaussian vectors gives that:

$$\begin{aligned} \frac{d\mathbb{P}(\vec{\Gamma}_{[n]} = \vec{\gamma}_{[n]})}{d\gamma} &\leq \left( \frac{\exp(\Phi_g(\varepsilon)\varepsilon^2)}{2\pi\sigma^2} \right)^{n/2} \times \\ &\exp\left( \frac{1}{2\sigma^2} \left( (-1 + 2\Phi_i(\varepsilon)\varepsilon^2) \sum_{i \in [n]} \gamma_i^2 + \frac{1}{2} \sum_{\substack{i,j \in [n] \\ i \neq j}} |\gamma_i||\gamma_j| \times (2\Phi_h(\varepsilon)\varepsilon)^{|i-j|} \right) \right). \end{aligned} \quad (75)$$

Bounding above  $|\gamma_i||\gamma_j|$  by  $\frac{1}{2}(\gamma_i^2 + \gamma_j^2)$ , that is bounded again by

$$(\Phi_j(\varepsilon)/2\pi\sigma^2)^{n/2} \exp\left( -\frac{1}{2\Phi_k(\varepsilon)\sigma^2} \sum_{i \in [n]} \gamma_i^2 \right), \quad (76)$$

where

$$\Phi_j(\varepsilon) := \exp(\Phi_g(\varepsilon)\varepsilon^2) \quad (77)$$

and

$$\Phi_k(\varepsilon) := \left( 1 - 2\Phi_i(\varepsilon)\varepsilon^2 - \frac{1}{2} \sum_{z \in \mathbf{Z}^*} (2\Phi_h(\varepsilon)\varepsilon)^{|z|} \right)_+^{-1} \quad (78)$$

are “quasi-one functions” again.

In the sequel we assume that  $\varepsilon$  was chosen small enough so that  $\Phi_j(\varepsilon), \Phi_k(\varepsilon) < \infty$ ; and we define the following vectorial random variable (which we are actually only interested in through its law):

**Definition 34** (Variable  $\vec{\Pi}_{\mathbf{N}}$ ).  $\vec{\Pi}_{\mathbf{N}}$  is a random vector on  $\mathbf{R}^{\mathbf{N}}$  whose entries are i.i.d.  $\mathcal{N}(0, \Phi_k(\varepsilon)\sigma^2)$ .

Then, Equation (76) can be rephrased into:

$$\frac{d\mathbb{P}(\vec{\Gamma}_{\llbracket n \rrbracket} = \vec{\gamma}_{\llbracket n \rrbracket})}{d\mathbb{P}(\vec{\Pi}_{\llbracket n \rrbracket} = \vec{\gamma}_{\llbracket n \rrbracket})} \leq (\Phi_j(\varepsilon)\Phi_k(\varepsilon))^{n/2} \quad \text{uniformly in } \vec{\gamma}_{\llbracket n \rrbracket}. \quad (79)$$

Therefore, for  $I \subset \llbracket n \rrbracket$  with  $|I| \geq pn$ :

$$\mathbb{P}(\forall i \in I \quad \Gamma_i \geq \alpha(\log i)_+^{1/2}) \leq (\Phi_j(\varepsilon)\Phi_k(\varepsilon))^{n/2} \mathbb{P}(\forall i \in I \quad \Pi_i \geq \alpha(\log i)_+^{1/2}). \quad (80)$$

But

$$\begin{aligned} \mathbb{P}(\forall i \in I \quad \Pi_i \geq \alpha(\log i)_+^{1/2}) &= \prod_{i \in I} \mathbb{P}(\Pi_i \geq \alpha(\log i)_+^{1/2}) \\ &= \prod_{i \in I} \mathbb{P}(\mathcal{N}(0, 1) \geq C_1(\log i)_+^{1/2}) \leq \prod_{i \in I} \exp(-C_1^2(\log i)_+/2) \\ &= \prod_{i \in I} (i \vee 1)^{-C_1^2/2} \leq (|I| - 1)_+!^{-C_1^2/2} \leq (\lceil pn \rceil - 1)_+!^{-C_1^2/2} \quad (81) \end{aligned}$$

(with  $C_1 := \alpha / \Phi_k(\varepsilon)^{1/2}\sigma$ ), where the penultimate inequality comes from ordering  $I =: \{i_0, i_1, \dots, i_{|I|-1}\}$  with  $i_0 < i_1 < \dots$ , and observing that then  $i_j \geq j$  for all  $j$ , so that  $(i_j \vee 1)^{-C_1^2/2} \leq (j \vee 1)^{-C_1^2/2}$ .

Combining (80) with (81) shows that  $\mathbb{P}(\Gamma_i \geq \alpha(\log i)_+^{1/2} \forall i \in I)$  decreases superexponentially in  $n$  uniformly in  $I$ , which finally proves Proposition 31 and hence Theorem 3.  $\square$

## 6 Application to financial mathematics

### 6.1 Two-way crossing property

Theorem 3 implies in particular that a stopping time  $\tau$  can almost-never be a local minimum at right for the trajectory of  $Z$ , i.e. that for almost-all  $\omega$ , there exist arbitrarily small  $\varepsilon > 0$  such that  $Z(\omega)_{\tau(\omega)+\varepsilon} < Z(\omega)_{\tau(\omega)}$ . Symmetrically (since  $-Z$  has the same law as  $Z$ ),  $\tau$  can almost-never be a local maximum for  $Z$ . The conjunction of these two facts implies that fBm has the “two-way crossing property” introduced by Bender in [Bender, 2012] (whose definition is recalled below for the sake of self-containment):

**Definition 35** (Two-way crossing property). A process  $X$  (whose increments<sup>[6]</sup> are) adapted to a filtration  $(\mathcal{B}_t)_t$  is said to satisfy the *two-way crossing (TWC) property* when, for any stopping time  $\tau$  adapted to  $(\mathcal{B}_t)_t$ , one has almost-surely:

$$\inf\{t > \tau \mid X_t > X_\tau\} = \inf\{t > \tau \mid X_t < X_\tau\}. \quad (82)$$

**Corollary 36.** *For any  $H \neq 1/2$ , fractional Brownian motion (hence also geometric fBm) satisfies the two-way crossing property.*

<sup>[6]</sup>Or logarithmic increments in the case of geometric fBm.

## 6.2 Financial context

The motivation for this work comes from the study of (geometric) fractional Brownian motion as a model for price processes in financial mathematics, which is a natural choice for modelling correlations and scale invariance, as was first suggested by [Mandelbrot, 1967]. fBm is however not a semimartingale, which in the classic setting leads to arbitrage opportunities, as was shown by [Delbaen and Schachermayer, 1994]: so, an efficient market should not allow for fBm price processes. . . [Rogers, 1997] showed however that arbitrage strategies for a fBm price process shall necessarily rely on making numerous transactions on extremely short scales of time. That suggested to look whether relaxing slightly some assumptions of the classic financial theory would nevertheless lead to a consistent theory allowing for fBm price processes: many work on this issue was carried out over the last two decades.

A first idea is to impose restrictions on the authorized trading strategies. In [Cheridito, 2003] it was suggested to restrict oneself to the so-called *Cheridito class* of strategies, in which a minimal waiting time is imposed between two transactions. In such a context it can be shown rather easily that fBm is arbitrage-free indeed. An intermediate class of trading strategies between the Cheridito class and the full class of predictable strategies, called *simple strategies*, was investigated by [Bender, 2012]: a simple strategy allows only for a fixed, finite number of transactions, but transaction times may be arbitrarily close to each other. In his paper Bender introduced the (TWC) property, and showed that this property (together with conditional full support, see below) was a necessary and sufficient condition for simple arbitrage to be ruled out. At that time it remained an open question however whether fBm did have this (TWC) property. . . My work answers positively to this question, and therefore proves that geometric fBm is proof against simple arbitrage. This was actually my initial motivation for this article.

Another approach for limiting arbitrage opportunities is to allow for any trading strategy, but to introduce transaction costs, generally proportional, of arbitrarily small rate. In this context [Guasoni, 2006] proved that fBm is arbitrage-free indeed. The proof of that property already relied on studying the properties of fBm after a stopping time: a sufficient condition for arbitrage-freeness is indeed the *stickiness* property, which says that the future trajectory of the price process after any stopping time may remain arbitrarily close to its initial value for an arbitrarily large amount of time, with positive probability. The stickiness property itself is a particular case of the *conditional full support* property, which says that future trajectories of the process after any stopping time have full support among the space of continuous trajectories (for the topology of locally uniform convergence). Conditional full support was shown to hold for fBm by [Guasoni et al., 2008].

Still in presence of transaction costs, more recently Czichowsky, Schachermayer & al. investigated on existence of a so-called *shadow price process* for an asset whose price is modelled by a non-semimartingale process (like geometric fBm): a shadow price process is a semimartingale process taking values in the bid-ask spread, whose optimal trading strategy in absence of transaction costs coincides with the optimal strategy for the actual price process in presence of transaction costs. The notion of optimality considered here is to maximise the expectation value of some (reasonable) utility function. Using the stickiness prop-

erty, [Czichowsky and Schachermayer, 2015] proved first that geometric fBm does admit a shadow price indeed in the case the utility function is bounded from above and defined on the whole  $\mathbf{R}$  (e.g.  $U(x) = 1 - e^{-x}$ ). Then they realised that a process satisfying Bender’s (TWC) condition (together with some boundedness estimate) would actually have a shadow price for a much larger class of utility functions, including in particular logarithmic and concave power utilities. Hence they used my results in [Czichowsky et al., 2016] to show that geometric fBm admits a shadow price for such utilities.

## A Conditional expectation of the fBm

This appendix is devoted to ending the proof of Lemma 15 initiated in § 2.2. At the point we have got to, what remains to do is showing that

$$C_1 \int_{s=-\infty}^0 ((v-s)^\eta - (-s)^\eta) dW_{\tau+s} \quad (83)$$

(seen as a trajectory indexed by  $v \in \mathbf{R}_+$ ) is actually equal to  $\mathbf{D}((Z_{\tau+u} - Z_\tau)_{u \leq 0})$  with  $\mathbf{D}$  defined by (7)-(8), where  $W$  is the ordinary Brownian motion driving the fBm  $Z$ . To alleviate notation, actually we will only prove this result for  $\tau \equiv 0$ , the original case being the same up to time translation of the increments (hence the informal definition (10) of  $\mathbf{D}$ ).

The starting point for our computation is the *Pipiras–Taqqu formula*, which says that Equation (1) defining the past increments of  $Z$  as a function of the past increments of  $W$  has an “inverse” giving back the past increments of  $W$  from the past increments of  $Z$ :

**Proposition 37** ([Pipiras and Taqqu, 2002, Cor. 1.1]). *In the context of § 1.1, one has almost-surely, for all  $t$ :*

$$W_t = \frac{C_1^{-1}}{\Pi(\eta)\Pi(-\eta)} \int_{\mathbf{R}} ((t-s)_+^{-\eta} - (-s)_+^{-\eta}) dZ_s. \quad (84)$$

(Recall that  $\Pi(\bullet)$  is Euler’s pi function extrapolating the factorial).

• From now on in this appendix, it will be convenient to shorthand “ $1 / \Pi(\eta)\Pi(-\eta)$ ” into “ $C_H$ ”.

So, let us use (84) to get (7)-(8). First, like (1), Equations (83) and (84) have to be interpreted by integrating by parts: for  $v \geq 0, s \leq 0$ , that means resp. that:

$$(83) = \eta C_1 \int_{-\infty}^0 \xi_{\eta-1}(-s, v) W_s ds; \quad (85)$$

$$\frac{W_t}{C_1^{-1} C_H} = \eta \int_{-\infty}^t \xi_{-\eta-1}(t-s, -t) (Z_s - Z_t) ds + \eta \int_t^0 (-s)^{-\eta-1} Z_s ds + (-t)^{-\eta} Z_t, \quad (86)$$

where we recall that  $\xi_r(a, b) := (a + b)^r - a^r$ . Hence, (83) is equal to:

$$\eta^2 C_H \int_{s=-\infty}^0 \int_{u=-\infty}^s \xi_{\eta-1}(-s, v) \xi_{-\eta-1}(s-u, -s) (Z_u - Z_s) ds du \quad (87)$$

$$+ \eta^2 C_H \int_{s=-\infty}^0 \int_{u=s}^0 \xi_{\eta-1}(-s, v) (-u)^{-\eta-1} Z_u ds du \quad (88)$$

$$+ \eta C_H \int_{-\infty}^0 \xi_{\eta-1}(-s, v) (-s)^{-\eta} Z_s ds. \quad (89)$$

Now we are going to re-write each of the terms (87)–(89) as an integral against  $Z_u du$ , in order to get (7)–(8). First, Term (89) is already of the wanted form, up to renaming ‘ $s$ ’ into ‘ $u$ ’. Next, Term (88) simplifies into:

$$\begin{aligned} (88) &= \eta^2 C_H \int_{u=-\infty}^0 \left( \int_{s=-\infty}^u \xi_{\eta-1}(-s, v) ds \right) (-u)^{-\eta-1} Z_u du \\ &= -\eta C_H \int_{-\infty}^0 \left[ \xi_{\eta}(-s, v) \right]_{s=-\infty}^u (-u)^{-\eta-1} Z_u du \\ &= -\eta C_H \int_{-\infty}^0 \xi_{\eta}(-u, v) (-u)^{-\eta-1} Z_u du. \end{aligned} \quad (90)$$

Term (87) is the hardest to get into the wanted form, because splitting naively the factor  $(Z_u - Z_s)$  would yield divergent integrals. To bypass that problem, we first make a truncation: for  $\varepsilon$  a small positive number,

$$\begin{aligned} (87) &\approx \eta^2 C_H \int_{s=-\infty}^0 \int_{u=-\infty}^{(1+\varepsilon)s} \xi_{\eta-1}(-s, v) \xi_{-\eta-1}(s-u, -s) (Z_u - Z_s) ds du \\ &= \eta^2 C_H \iint_{\substack{s < 0 \\ u < (1+\varepsilon)s}} \xi_{\eta-1}(-s, v) \xi_{-\eta-1}(s-u, -s) Z_u ds du \end{aligned} \quad (91)$$

$$- \eta^2 C_H \iint_{\substack{s < 0 \\ u < (1+\varepsilon)s}} \xi_{\eta-1}(-s, v) \xi_{-\eta-1}(s-u, -s) Z_s ds du. \quad (92)$$

By the change of variables  $(s, u) \leftarrow (u - s, u)$ ,

$$\begin{aligned} (91) &= \eta^2 C_H \int_{u=-\infty}^0 \left( \int_{s=-\infty}^{\varepsilon u / (1+\varepsilon)} \mathbf{1}_{s > u} \xi_{\eta-1}(s-u, v) \xi_{-\eta-1}(-s, s-u) ds \right) Z_u du \\ &\approx \eta^2 C_H \int_{u=-\infty}^0 \left( \int_{s=-\infty}^{\varepsilon u} \mathbf{1}_{s > u} \xi_{\eta-1}(s-u, v) \xi_{-\eta-1}(-s, s-u) ds \right) Z_u du \end{aligned} \quad (93)$$

(where by “ $\approx$ ” we mean that, for all  $v$ , the difference between the two members from either side of the ‘ $\approx$ ’ sign tends to 0 as  $\varepsilon \rightarrow 0$ , as one can check by simple estimates); and by the change of variables  $(s, u) \leftarrow (u - s, s)$ ,

$$(92) = -\eta^2 C_H \int_{u=-\infty}^0 \left( \int_{s=-\infty}^{\varepsilon u} \xi_{\eta-1}(-u, v) \xi_{-\eta-1}(-s, -u) ds \right) Z_u du. \quad (94)$$

So,

$$(87) \approx \eta^2 C_H \int_{-\infty}^0 \left( \int_{-\infty}^{\varepsilon u} J(v, u, s) ds \right) Z_u du, \quad (95)$$

with

$$J(v, u, s) := \mathbf{1}_{s>u} \xi_{\eta-1}(s-u, v) \xi_{-\eta-1}(-s, s-u) - \xi_{\eta-1}(-u, v) \xi_{-\eta-1}(-s, -u). \quad (96)$$

But  $\int^0 J(v, u, s) ds$  does converge, so, letting  $\varepsilon$  tend to 0, we get in the end:

$$(87) = \eta^2 C_H \int_{-\infty}^0 \left( \int_{-\infty}^0 J(v, u, s) ds \right) Z_u du. \quad (97)$$

Summing (89), (90) and (97), and observing that  $\xi_{\eta-1}(-u, v)(-u)^{-\eta} - \xi_{\eta}(-u, v) \times (-u)^{-\eta-1} = -v(v-u)^{\eta-1}(-u)^{-\eta-1}$ , finally yields Equations (7)-(8).  $\square$

## B Explicit estimate for the supremum of Gaussian processes

*Proof of Lemma 28.* Let  $\theta \in (0, 1]$  and let  $X$  satisfying the assumptions of the lemma. Then obviously, for the continuous version of  $X$ :

$$\|X\|(\omega) \leq \sum_{i=0}^{\infty} \sup_{a \in [2^i \llbracket} |X(\omega)_{a2^{-i}} - X(\omega)_{(a+1)2^{-i}}|. \quad (98)$$

Therefore, for  $(\gamma_i)_i$  a sequence of positive real numbers such that  $\sum_i \gamma_i = 1$ , one has that, for all  $x \geq 0$ :

$$\begin{aligned} \mathbb{P}(\|X\| \geq x) &\leq \sum_{i=0}^{\infty} \mathbb{P} \left( \sup_{a \in [2^i \llbracket} |X_{a2^{-i}} - X_{(a+1)2^{-i}}| \geq \gamma_i x \right) \\ &\leq \sum_i 2^i \sup_a \mathbb{P}(|X_{a2^{-i}} - X_{(a+1)2^{-i}}| \geq \gamma_i x). \end{aligned} \quad (99)$$

But, uniformly in  $a$ ,

$$\begin{aligned} \mathbb{P}(|X_{a2^{-i}} - X_{(a+1)2^{-i}}| \geq \gamma_i x) &= \mathbb{P}(\mathcal{N}(0, 1) \geq \gamma_i x / \text{Var}^{1/2}(X_{a2^{-i}} - X_{(a+1)2^{-i}})) \\ &\leq \mathbb{P}(\mathcal{N}(0, 1) \geq 2^{i\theta} \gamma_i x) \leq 2 \exp(-2^{2i\theta-1} (\gamma_i x)^2); \end{aligned} \quad (100)$$

so, taking  $\gamma_i := (1 - 2^{-\theta/2})2^{-i\theta/2}$ :

$$\mathbb{P}(\|X\| \geq x) \leq \sum_{i=0}^{\infty} 2^{i+1} \exp(-(1 - 2^{-\theta/2})^2 \times 2^{i\theta-1} x^2). \quad (101)$$

Provided  $x \geq 2 / (1 - 2^{-\theta/2})\theta^{1/2} =: C_m(\theta)$ , one has (bounding  $2^{i\theta}$  below by  $(1 + i\theta \log 2)$ )

$$\exp(-2^{i\theta} x^2 / 2(1 - 2^{-\theta/2})^2) \leq \exp(-x^2 / 2(1 - 2^{-\theta/2})^2 - \log(4)i), \quad (102)$$

so that, for  $x \geq C_m$ :

$$\mathbb{P}(\|X\| \geq x) \leq \left( \sum_{i=0}^{\infty} 2^{i+1} 4^{-i} \right) \exp(-x^2 / 2(1 - 2^{-\theta/2})^2) =: 4e^{-C_c(\theta)x^2}. \quad (103)$$

On the other hand, for  $x < C_m$  one has obviously  $\mathbb{P}(\|X\| \geq x) \leq 1$ ; so Equation (49) follows with  $C_d := 4 \vee e^{C_c C_m^2}$ .  $\square$

## C Almost diagonal matrices

*Proof of Lemma 33.* Consider  $\mathbf{A}$  satisfying the assumptions of the lemma, and denote  $\mathbf{I}_n - \mathbf{A} =: \mathbf{H}$ . The first part of this proof will consist in deriving estimates on the entries of  $\mathbf{H}$  and its powers. Denote respectively

$$\mathbf{H} =: ((h_{ij}))_{i,j \in \llbracket n \rrbracket}; \quad (104)$$

$$\mathbf{H}^k =: ((h_{ij}^{(k)}))_{i,j \in \llbracket n \rrbracket} \quad \forall k \geq 0. \quad (105)$$

Then the assumptions of the lemma ensure that one has  $|h_{ii}| = 0 \forall i$ , resp.  $|h_{ij}| \leq \varepsilon^{|i-j|} \forall i \neq j$ , and hence

$$|h_{ij}^{(k+1)}| \leq \sum_{i' \in \llbracket n \rrbracket} |h_{ii'}| |h_{i'j}^{(k)}| = \sum_{i' \neq i} \varepsilon^{|i-i'|} |h_{i'j}^{(k)}| \quad \forall i, j \quad \forall k. \quad (106)$$

That suggests to define by induction:

$$\begin{cases} \mathfrak{h}_z^{(0)} := \mathbf{1}_{z=0} & \forall z \in \mathbf{Z} \\ \mathfrak{h}_z^{(k+1)} := \sum_{z' \neq z} \varepsilon^{|z-z'|} \mathfrak{h}_{z'}^{(k)} & \forall z \in \mathbf{Z} \quad \forall k \geq 0, \end{cases} \quad (107)$$

so that one has

$$|h_{ij}^{(k)}| \leq \mathfrak{h}_{i-j}^{(k)} \quad \forall i, j \quad \forall k. \quad (108)$$

The interest of having introduced the  $\mathfrak{h}_z^{(k)}$ 's is that these are easier to bound than the  $h_{ij}^{(k)}$ 's themselves. To bound the  $\mathfrak{h}_z^{(k)}$ 's, we begin with observing that one has obviously by induction:

$$\begin{aligned} \mathfrak{h}_z^{(k)} &= \sum_{\substack{(0=:s_0, s_1, s_2, \dots, s_{k-1}, z=:s_k) \\ 0 \neq s_1, s_1 \neq s_2, \dots, s_{k-1} \neq z}} \varepsilon^{\sum_{i \in \llbracket k \rrbracket} |s_i - s_{i+1}|} \\ &= \sum_{n \geq 0} \text{card} \left\{ (0 =: s_0, s_1, \dots, s_{k-1}, z =: s_k) \mid s_{i+1} \neq s_i \forall i, \sum_i |s_i - s_{i+1}| = n \right\} \varepsilon^n. \end{aligned} \quad (109)$$

To bound the cardinality appearing in (109), we observe that a  $(k+1)$ -tuple  $(0, s_1, s_2, \dots, s_{k-1}, z)$  such that  $s_{i+1} \neq s_i \forall i$  and  $\sum_i |s_i - s_{i+1}| = n$  (we will call such a  $(k+1)$ -tuple as *valid*) can be *coded* by a word of  $n$  symbols from  $\{+, +_1, -, -_1\}$ , in the following way: successively, for each  $i$  we write  $(|s_{i+1} - s_i| - 1)$  symbols “ $\text{sgn}(s_{i+1} - s_i)$ ” followed by a symbol “ $\text{sgn}(s_{i+1} - s_i)$ ”—for instance, for  $k = 5$ ,  $z = 2$ ,  $n = 8$ , one would have

$$(0, 1, 4, 2, 1, 2) \mapsto “+|++++|---|+|”. \quad (110)$$

Obviously such a coding is injective. Moreover, for given  $k, z, n$ , an  $n$ -character word may be the image of a valid  $(k+1)$ -tuple only if  $z$  and  $n$  have the same parity, that the word contains  $(n+z)/2$  symbols from  $\{+, +_1\}$  vs.  $(n-z)/2$  symbols from  $\{-, -_1\}$ , and that  $k$  exactly of the  $n$  symbols, necessarily including the last one, are from  $\{+|, -|\}$ . Henceforth:

$$\begin{aligned} \text{card} \left\{ (0, s_1, s_2, \dots, s_{k-1}, z) \mid s_{i+1} \neq s_i \forall i \text{ and } \sum_i |s_i - s_{i+1}| = n \right\} \\ \leq \mathbf{1}_{2|n-z} \binom{n}{(n-|z|)/2} \binom{n-1}{k-1}. \end{aligned} \quad (111)$$

In the end, combining (108), (109) and (111):

$$|h_{ij}^{(k)}| \leq \sum_{m \geq 0} \binom{|i-j|+2m}{m} \binom{|i-j|+2m-1}{k-1} \varepsilon^{|i-j|+2m} \quad \forall i, j \quad \forall k. \quad (112)$$

After these preliminary estimates, let us turn to proving the lemma itself. We begin with the first part, namely, bounding  $\det \mathbf{A}$  from below. For  $\mathbf{X} =: ((x_{ij}))_{i,j}$  an  $n \times n$  matrix, denote

$$\|\mathbf{X}\| := \sup_{j \in \llbracket n \rrbracket} \sum_{i \in \llbracket n \rrbracket} |x_{ij}| : \quad (113)$$

$\|\cdot\|$  is the operator norm of  $\mathbf{X}$  when seen as an operator from  $\ell^1(\llbracket n \rrbracket)$  into itself, so it is sub-multiplicative. Then, the formula

$$\log(\mathbf{I}_n - \mathbf{H}) = \sum_{k=1}^{\infty} k^{-1} \mathbf{H}^k \quad (114)$$

converges as soon as  $\|\mathbf{H}\| < 1$ , then yielding:

$$\begin{aligned} |\operatorname{tr} \log(\mathbf{I}_n - \mathbf{H})| &\leq \sum_{k=1}^{\infty} k^{-1} |\operatorname{tr} \mathbf{H}^k| \\ &\leq 0 + \frac{1}{2} |\operatorname{tr} \mathbf{H}^2| + n \sum_{k \geq 3} k^{-1} \|\mathbf{H}^k\| \\ &\leq \frac{1}{2} |\operatorname{tr} \mathbf{H}^2| + n \sum_{k \geq 3} k^{-1} \|\mathbf{H}\|^k. \end{aligned} \quad (115)$$

But the assumptions on the entries of  $\mathbf{H}$  imply that

$$\|\mathbf{H}\| \leq \sum_{z \in \mathbf{Z}^*} \varepsilon^{|z|} = \frac{2\varepsilon}{1-\varepsilon}, \quad (116)$$

which is  $< 1$  as soon as  $\varepsilon < 1/3$ ; and on the other hand, we get from (112) that, for all  $i$ ,

$$|h_{ii}^{(2)}| \leq \sum_{m \geq 1} \binom{2m}{m} (2m-1) \varepsilon^{2m} \leq 2\varepsilon^2 + \sum_{m \geq 2} 2^{2m} (2m-1) \varepsilon^{2m} =: 2\Phi_n(\varepsilon) \varepsilon^2, \quad (117)$$

so that  $|\operatorname{tr} \mathbf{H}^2| \leq 2n\Phi_n(\varepsilon) \varepsilon^2$ . In the end:

$$\begin{aligned} \det \mathbf{A} &= \det \exp \log(\mathbf{I}_n - \mathbf{H}) = \exp \operatorname{tr} \log(\mathbf{I}_n - \mathbf{H}) \\ &\geq \exp \left( -n\Phi_n(\varepsilon) \varepsilon^2 - n \sum_{k \geq 3} k^{-1} \left( \frac{2\varepsilon}{1-\varepsilon} \right)^k \right) =: \exp(-n\Phi_g(\varepsilon) \varepsilon^2), \end{aligned} \quad (118)$$

which is Equation (72).

Now let us handle the second part of the lemma, namely, bounding the entries of  $(\mathbf{A}^{-1} - \mathbf{I}_n)$ . Provided  $\|\mathbf{H}\| < 1$ , one has

$$\mathbf{A}^{-1} - \mathbf{I}_n = \sum_{k=1}^{\infty} \mathbf{H}^k, \quad (119)$$

so that

$$|b_{ij} - \mathbf{1}_{i=j}| \leq \sum_{k=1}^{\infty} |h_{ij}^{(k)}| \quad \forall i, j. \quad (120)$$

To bound the r-h.s. of (120), we write that, starting from (112):

$$\begin{aligned} \sum_{k \geq 1} |h_{ij}^{(k)}| &\leq \sum_{m \geq 0} \binom{|i-j|+2m}{m} \left( \sum_{k=1}^{|i-j|+2m} \binom{|i-j|+2m-1}{k-1} \right) \varepsilon^{|i-j|+2m} \\ &= \sum_{m \geq 0} \binom{|i-j|+2m}{m} \mathbf{1}_{|i-j|+2m \geq 1} 2^{|i-j|+2m-1} \varepsilon^{|i-j|+2m} \\ &= \left( \sum_{m \geq 0} \mathbf{1}_{|i-j|+2m \geq 1} \binom{|i-j|+2m}{m} (4\varepsilon^2)^m \right) 2^{|i-j|-1} \varepsilon^{|i-j|}. \end{aligned} \quad (121)$$

But we observe that, for  $z \geq 1$ ,  $x > 0$ ,

$$\begin{aligned} \sum_{m \geq 0} \binom{z+2m}{m} x^m &= 1 + \sum_{m \geq 1} \binom{z+2m}{m} x^m \leq 1 + \sum_{m \geq 1} \frac{(z+2m)^m}{m!} x^m \\ &\leq 1 + \sum_{m \geq 1} \frac{(2z)^m + (4m)^m}{m!} x^m \\ &\leq 1 + \sum_{m \geq 1} \frac{(2zx)^m}{m!} + \sum_{m \geq 1} (4ex)^m = e^{2zx} + \frac{4ex}{1-4ex} \\ &\leq \left( e^{2x} + \frac{4ex}{1-4ex} \right)^z =: \Phi_o(x)^z, \end{aligned} \quad (122)$$

so that here, for  $i \neq j$ :

$$\sum_{k=1}^{\infty} |h_{i,j}^{(k)}| \leq \Phi_o(4\varepsilon^2)^{|i-j|} 2^{|i-j|-1} \varepsilon^{|i-j|} =: 2^{|i-j|-1} (\Phi_h(\varepsilon)\varepsilon)^{|i-j|}, \quad (123)$$

which is Equation (73).

Equation (74) for the case  $i = j$  is derived in the same way as (73), with just a few minor differences at the beginning of the computation: namely, in the l-h.s. of (121), we treat apart the cases “ $k = 1$ ” (which yields zero here since  $h_{ii} = 0$  by assumption) and “ $k = 2$ ” (which has already been handled by (117)); then all the sequel is the same.  $\square$

*Remark 38.* The bounds (72)–(74) of Lemma 33 are optimal at first order. Actually this is much more than needed to prove Theorem 3, and we could have got a sufficient result with a shorter proof; but it seemed interesting to me to state the sharp version of the lemma.

## Acknowledgements

This work owes much to Christian Bender, the author of [Bender, 2012]. First, he was the person who drew my attention onto the question that Corollary 36 solves—that was at a seminar in Nancy in 2012. Moreover, Christian invited

me several times in Saarbrücken to discuss about my proof sketch, and gave me several worthy pieces of technical help to write it down in a proper way.

I also want to thank Walter Schachermayer for his enthusiastic support after he read my preprint on arXiv, for his enlightening explanations on the financial context around my work, and for the quite pleasant discussions that we had in Zurich together with Christoph Czichowsky (who helped me very kindly too) about possible applications of my results.

Last but not least, the careful work of my two referees has led to significant improvements in the final version of this paper: may they be thanked too!

## References

- Miguel Arcones. On the law of the iterated logarithm for Gaussian processes. *J. Theoret. Probab.*, 8(4):877–903, 1995.
- Christian Bender. Simple arbitrage. *Ann. Appl. Probab.*, 22(5):2067–2085, 2012.
- Patrick Cheridito. Arbitrage in fractional Brownian motion models. *Finance Stoch.*, 7(4):533–553, 2003.
- Christoph Czichowsky and Walter Schachermayer. Portfolio optimisation beyond semimartingales: Shadow princes and fractional Brownian motion. Preprint, [arXiv:1505.02416](https://arxiv.org/abs/1505.02416), 2015.
- Christoph Czichowsky, Rémi Peyre, Walter Schachermayer, and Junjian Yang. Shadow prices, fractional Brownian motion, and portfolio optimisation. Work in progress, 2016.
- Freddy Delbaen and Walter Schachermayer. A general version of the fundamental theorem of asset pricing. *Math. Ann.*, 300(3):463–520, 1994.
- Paolo Guasoni. No arbitrage under transaction costs, with fractional Brownian motion and beyond. *Math. Finance*, 16(3):569–582, 2006.
- Paolo Guasoni, Miklós Rásonyi, and Walter Schachermayer. Consistent price systems and face-lifting pricing under transaction costs. *Ann. Appl. Probab.*, 18(2):491–520, 2008.
- Benoît Mandelbrot. The variation of some other speculative prices. *The Journal of Business*, 40(4):393–413, 1967.
- Ivan Nourdin. *Selected aspects of fractional Brownian motion*, volume 4 of *Bocconi & Springer Series*. Springer; Bocconi University Press, 2012.
- Vladas Pipiras and Murad S. Taqqu. Deconvolution of fractional Brownian motion. *J. Time Ser. Anal.*, 23(4):487–501, 2002.
- Chris Rogers. Arbitrage with fractional Brownian motion. *Math. Finance*, 7(1): 95–105, 1997.