

Regulation of bacterial motility in response to low pH in *Escherichia coli*: the role of H-NS protein

Olga A. Soutourina,^{1†} Evelyne Krin,¹ Christine Laurent-Winter,² Florence Hommais,¹ Antoine Danchin¹ and Philippe N. Bertin^{1‡}

Author for correspondence: Philippe N. Bertin. Tel: +32 14 33 34 36. Fax: +32 14 32 03 13.
e-mail: pbertin@sckcen.be

Unité de Génétique des
Génomés Bactériens¹ and
Génopole-Plateau
Protéomique², Institut
Pasteur, Paris, France

The effect of detrimental conditions on bacterial motility in *Escherichia coli* was investigated. Expression profiling of mutant *E. coli* strains by DNA arrays and analysis of phenotypic traits demonstrated that motility and low-pH resistance are coordinately regulated. Analysis of transcriptional fusions suggests that bacterial motility in response to an acidic environment is mediated via the control by H-NS of *flhDC* expression. Moreover, the results suggested that the presence of an extended mRNA 5' end and DNA topology are required in this process. Finally, the presence of a similar regulatory region in several Gram-negative bacteria implies that this mechanism is largely conserved.

Keywords: acidic pH, osmolarity, DNA supercoiling, DNA array

INTRODUCTION

Complex cellular responses are often controlled by regulatory networks in which transcription factors regulate the expression of a diverse set of target genes. In eukaryotes such complex systems are implicated in cell differentiation into specialized tissues and in the maintenance of tissue homeostasis (Arnold & Winter, 1998; Kawakami *et al.*, 2000; Relaix & Buckingham, 1999). In prokaryotes, multiple regulatory networks are usually organized in a similar way to those in higher organisms, with master regulatory genes at the top of the hierarchy, ensuring adequate responses and/or transformation into another stable cellular state to fine tune cellular metabolism. As an example, RpoS is a central element that governs the expression of bacterial stress- or stationary-phase-induced genes (Hengge-Aronis, 1999). Studies of genetic circuits such as those involved in bacteriophage infection, chemotaxis or the cell division cycle in prokaryotes have also provided important insight into the understanding of regulatory networks (Huang, 1999).

Bacterial flagellum biosynthesis is under the control of the *flhDC* master operon, which governs motility and chemotaxis, as well as differentiation into swarming cells in enterobacteria. Moreover, this operon ensures global communication between flagellar genes and external factors, as well as cell division (Aizawa & Kubori, 1998). The complex motility and chemotaxis system in *Escherichia coli* includes nearly 50 genes organized in an ordered cascade in which the expression of a gene located at a given level requires the transcription of another one at a higher level (Macnab, 1996). This system is subject to a complex regulation by multiple environmental factors and regulatory proteins. For example, flagellum biosynthesis is sensitive to catabolite repression (Adler & Templeton, 1967; Silverman & Simon, 1974; Yokota & Gots, 1970) and is inhibited by stressful environmental conditions, such as increased temperature and high osmolarity (Adler & Templeton, 1967; Li *et al.*, 1993). Moreover, numerous mutations in stress-related genes, such as those encoding heat-shock proteins, membrane components or DNA replication initiation factors, are known to affect motility (Farr *et al.*, 1989; Kitamura *et al.*, 1994; Mizushima *et al.*, 1995, 1997; Shi *et al.*, 1992, 1993a) by repressing transcription of the *flhDC* master operon (Mizushima *et al.*, 1995, 1997; Shi *et al.*, 1993b). Unlike *flhDC* transcriptional control by cAMP–CAP complex (Soutourina *et al.*, 1999), the mechanism by which stress-related conditions affect master operon expression remains still largely unknown.

[†] Present address: Laboratoire de Biochimie, UMR 7654, CNRS-Ecole Polytechnique, 91128 Palaiseau Cedex, France.

[‡] Present address: Laboratory of Microbiology, Boeretang 200, B-2400 Mol, Belgium.

Abbreviation: CAT, chloramphenicol acetyltransferase.

H-NS is a nucleoid-associated protein known to be involved in the control of motility in *E. coli* (Bertin *et al.*, 1994) and in *Salmonella typhimurium* (Hinton *et al.*, 1992). This protein positively controls the master flagellar operon but the mechanism of this regulation remains unclear (Kutsukake, 1997; Soutourina *et al.*, 1999). In the present study we demonstrated that flagellar gene expression is inhibited under low-pH conditions and that this regulation of the *flhDC* master operon may be dependent on the H-NS protein. For the first time, we provide evidence that the 5' end of mRNA plays a crucial role in *flhDC* expression in response to environmental factors.

METHODS

Bacterial strains and growth conditions. Bacterial strains and plasmids used in this study are listed in Table 1. Strains were grown in Luria-Bertani (LB), tryptone, M9 or M63 media (all media as in Miller, 1992), supplemented as indicated with 0.1% (w/v) Casamino acids, 0.4% (w/v) sodium succinate, 0.4% (w/v) glycerol or 0.4% (w/v) glucose as a carbon source. Tryptone swarm plates containing 1% Bacto-tryptone, 0.5% NaCl and 0.3% Bacto-agar were used to test bacterial motility as previously described (Bertin *et al.*, 1999), except for plates at pH 4.6, which contained 0.5% Bacto-agar. Metabolism of β -glucosides was tested on MacConkey indicator agar plates with 1% salicin as a carbon source. When required, antibiotics were added at the following concentration: ampicillin, 50 $\mu\text{g ml}^{-1}$, and kanamycin, 20 $\mu\text{g ml}^{-1}$. All experiments were performed in accordance with the European regulation requirements concerning the contained use of Genetically Modified Organisms of Group I (agreement no. 2735).

Chloramphenicol acetyltransferase (CAT) assay. Strains were grown in tryptone medium supplemented with sodium succinate or M9 medium supplemented with Casamino acids and glycerol at pH 7.0 or 4.6 to an OD₆₀₀ of 0.15 to 0.3. CAT

activity was measured with 5,5'-dithiobis-2-nitrobenzoic acid (DTNB) on cell extracts as previously described (Soutourina *et al.*, 1999).

Resistance to low pH. Strains were grown to stationary phase overnight in M9 medium, pH 5.5, supplemented with glucose and Casamino acids. Acidic stress was analysed in M9 medium at pH 2.5 supplemented with 0.012% glutamate as previously described (Hommais *et al.*, 2001).

Two-dimensional gel electrophoresis. Strains were grown in M9 medium supplemented with Casamino acids and glycerol at pH 7.0 or 4.6 to an OD₆₀₀ of 0.7. Total protein extracts and two-dimensional gel electrophoresis were carried out as previously described (Hommais *et al.*, 2001; Laurent-Winter *et al.*, 1997).

Expression profiling. Bacterial cells were grown in M63 minimal medium supplemented with glucose (Miller, 1992) to an OD₆₀₀ of 0.6. Handling of RNA, cDNA synthesis from 10 μg RNA, hybridization on DNA arrays (Panorama *E. coli* gene arrays from Sigma-GenoSys Biotechnologies) and data analysis were performed as previously described (Hommais *et al.*, 2001). Briefly, hybridization probes were generated from 10 μg RNA following standard cDNA synthesis using [α -³²P]dCTP (7.4 \times 10¹³–1.1 \times 10¹⁴ Bq mmol⁻¹, New England Nuclear), AMV reverse transcriptase (Roche) and *E. coli* labelling primers (Sigma-Genosys). The prehybridization and hybridization were carried out according to the manufacturer's recommendations with some modifications (Hommais *et al.*, 2001). Blots were exposed to PhosphorImager screen (Molecular Dynamics) and were then scanned on a 445SI PhosphorImager. The intensity of each dot was measured with the XDOTSREADER software (Cose) and analysed using an Excel spreadsheet.

In vitro transcription assays. *In vitro* transcription experiments were performed with pDIA546 containing the entire *flhDC* regulatory region as previously described (Soutourina *et al.*, 1999). Plasmid pDIA546 was restricted by *EcoRI* for 2 h at 37 °C and used as linearized template for *in vitro* transcription.

Table 1. Bacterial strains and plasmids

Strain or plasmid	Relevant genotype	Reference or source
Strain		
MG1655	Wild-type	CGSC 6300
PS2209	Wild-type	Bertin <i>et al.</i> (1994)
PS2652	PS2209 <i>hms-1001</i>	Bertin <i>et al.</i> (1994)
FB8	Wild-type	Bruni <i>et al.</i> (1977)
BE1410	FB8 <i>hms-1001</i>	Laurent-Winter <i>et al.</i> (1997)
BE2120	<i>hms-1001</i> carrying a <i>mot</i> ⁺ suppressor mutation	This study
BE2121	<i>hms-1001</i> carrying a <i>mot</i> ⁺ suppressor mutation	This study
Plasmid		
pDIA528	pKK232-8-derivative carrying the <i>flhDC</i> promoter (nt -213 to +78)	Soutourina <i>et al.</i> (1999)
pDIA545	pKK232-8 derivative carrying the <i>flhDC</i> regulatory region (nt -213 to +205)	Soutourina <i>et al.</i> (1999)
pDIA546	pJCD01 derivative carrying the <i>flhDC</i> regulatory region (nt -213 to +202)	Soutourina <i>et al.</i> (1999)
pDIA559	pKK232-8-derivative carrying the <i>flhDC</i> promoter region	Soutourina <i>et al.</i> (1999)
pPM61	ColE1 derivative multicopy expression vector carrying the <i>flhDC</i> operon under its native promoter	Bartlett <i>et al.</i> (1988)

RESULTS**Isolation and characterization of suppressor mutations**

The comparative analysis of expression profiles in *E. coli* wild-type and *hns* strains (Hommais *et al.*, 2001) revealed two major alterations in the *hns* background: on the one hand, a decrease in flagellar gene expression, and on the other hand, an increase in the acidic pH resistance gene expression, suggesting that these two H-

NS-controlled phenotypes may be related to each other. To test this possible link, we isolated and characterized in an *hns* background suppressor mutations with regard to motility. Several spontaneous mutants showing H-NS-independent swarming were isolated at a frequency of about 10^{-9} by incubating *hns* strains on semi-solid agar plates at 30 °C for 24–48 h, and two of them were purified. Attempts to locate these suppressor mutations by genetic techniques were unsuccessful, which suggests that they may involve more than one gene, e.g. an

Table 2. Effect of suppressor mutations on various H-NS-related phenotypes

Strain (phenotype)	Mucoidy*	β -Glucoside utilization†	Motility‡	Survival (%)§
FB8 (wild-type)	–	–	30+3	<0.001
BE1410 (<i>hns</i>)	+	+	3+1	24+4
BE2120 (<i>mot</i> ⁺)	+	+	18+2	<0.005
BE2121 (<i>mot</i> ⁺)	+	+	21+2	<0.001

* +, Mucoid phenotype observed on LB medium; –, no mucoid phenotype.

† +, Ability to metabolize β -glucoside revealed by the appearance of red colonies on MacConkey-salicylic agar plates; –, lack of ability to metabolize β -glucoside, indicated by white colonies.

‡ Diameter of the swarming ring (in mm) after 15 h at 30 °C. Data are the mean values \pm standard deviations of three independent experiments.

§ Percentage survival is calculated as $100 \times$ number of c.f.u. per ml remaining after low-pH treatment divided by the initial c.f.u. per ml at time zero. Data are the mean values of three independent experiments.

Table 3. Expression profiling by DNA arrays of various *E. coli* strains

Gene	Expression level* in strain (phenotype):				Function
	FB8 (wild-type)	BE1410 (<i>hns</i>)	BE2120 (<i>mot</i> ⁺)	BE2121 (<i>mot</i> ⁺)	
					Flagellum biosynthesis
<i>flgB</i>	0.80	–	0.38	1.36	Proximal rod protein
<i>flgD</i>	0.67	–	0.61	0.86	Basal body rod modification protein
<i>flgG</i>	0.75	0.29	0.61	1.32	Distal rod protein
<i>flgK</i>	1.02	–	0.80	1.04	Hook-associated protein 1
<i>flgL</i>	1.02	–	0.44	1.34	Hook-associated protein 3
<i>fliA</i>	0.72	–	0.30	0.75	Flagella-specific sigma factor
<i>fliC</i>	7.04	–	0.26	0.41	Flagellin
<i>fliD</i>	1.57	–	0.44	1.66	Hook-associated protein 2
<i>fliJ</i>	1.10	0.12	1.01	0.95	Export for some substrates
<i>fliK</i>	0.83	–	0.66	1.39	Hook length control protein
					Acid pH resistance
<i>gadA</i>	0.57	16.98	2.76	2.03	Glutamate decarboxylase α
<i>gadB</i>	0.43	15.16	2.33	1.83	Glutamate decarboxylase β
<i>gadC</i>	0.26	8.91	1.21	0.39	Putative amino acid antiporter
<i>gadX</i>	–	3.48	2.83	0.83	Regulatory protein
<i>hdeA</i>	0.36	10.67	2.75	0.66	Putative chaperone-like protein involved in low pH resistance
<i>hdeB</i>	–	9.42	2.08	0.37	Involved in low pH resistance
<i>hdeD</i>	–	3.17	0.40	0.57	Unknown function

* Expression level (arbitrary units) was measured as previously described (Hommais *et al.*, 2001). –, Below the background level.

essential locus with an additional compensatory mutation. The impossibility of transferring these mutations to other strains by P1 transduction or of obtaining strains with similar phenotype by transposon insertion further supports the possible existence of distant mutations. The presence of such multiple genetic alterations could be explained by the high frequency of spontaneous mutation observed in *hns* strains (Lejeune & Danchin, 1990). A similar failure to characterize suppressor mutations involved in the thermoregulation of motility has been reported in *Yersinia enterocolitica* (Rohde *et al.*, 1994).

To investigate the effect of the suppressor mutations on the physiology of *E. coli*, we tested various H-NS-related phenotypes in strains BE2120 and BE2121. Like the original *hns* mutant, both strains remained mucoid and able to use salicin as a carbon source (Table 2). In contrast, the mutations reversed, as expected, the loss of motility on semi-solid medium. Surprisingly, the suppressor mutants also showed a strong susceptibility to low pH, similar to that of the wild-type strain (Table 2).

To address the mechanism underlying the possible link between loss of motility and low pH resistance, expression profiling was performed using DNA arrays. As seen in Table 3, the major differences concerned the expression of genes involved in flagellum biosynthesis and in low pH resistance. Indeed, the transcript level of many flagellar genes measured in suppressor strains BE2120 and BE2121 was close to that in the wild-type strain. More importantly, a strong reduction in the expression of the *gad* and *hde* genes involved in resistance to low pH (Hommais *et al.*, 2001) was observed in these mutants in comparison with the *hns* strain (Table 3). These results are consistent with the alteration of motility and acid resistance we observed (Table 2) and support a coordinate regulation by H-NS of both processes in *E. coli*.

The analysis of expression profiles of the suppressor mutants did not allow us to identify an altered expression of genes known to be involved in the control of motility, i.e. CsrA- or HdfR-encoding genes (Ko & Park, 2000; Romeo, 1998; Wei *et al.*, 2001). Furthermore, both genes were not mutated in the suppressor strains (data not shown). Finally, spontaneous mutants were also obtained in an *hns stpA* double mutant context (data not shown), suggesting that the motility reversion process we observed is independent of the presence of StpA, in agreement with our recent data (Bertin *et al.*, 2001). Taken together, these observations suggest that these proteins do not play any role in the H-NS-regulated control of motility and low pH resistance.

Regulation of motility under acidic pH conditions

In *E. coli*, the optimum pH for motility and chemotaxis is close to that for growth (Adler, 1973; Adler & Templeton, 1967). However, the control of bacterial motility by acidic pH, which can reflect the growth conditions frequently encountered by enterobacteria




Fig. 1. Effect of acidic pH on motility and flagellin synthesis. (A) Motility of the *E. coli* wild-type strain was assayed on semi-solid medium plates at pH 7.0 and at pH 4.6. Plates were incubated for 13–15 h at 30 °C. (B) Protein extracts of wild-type strain MG1655 grown at pH 7.0 and at pH 4.6 were resolved by two-dimensional electrophoresis and silver stained. Only the region in the vicinity of FliC is shown. FliC is indicated by an arrow. To facilitate the comparison, NusA and PtsI are indicated as landmarks.

inside their host (Mahan *et al.*, 1996), has not yet been well documented. To investigate the direct effect of low pH on motility, we tested the swarming behaviour of wild-type *E. coli* on semi-solid plates at neutral and acidic pH. As seen in Fig. 1(A), a loss of motility was observed under low pH conditions. To determine whether this alteration in swarming behaviour resulted from a lack of flagella, we analysed the flagellin content by two-dimensional gel electrophoresis. As seen in Fig. 1(B), the spot corresponding to the FliC protein on our *E. coli* two-dimensional protein map (Hommais *et al.*, 2001) was undetectable in the protein extract of bacteria grown at low pH by comparison with those grown at




Fig. 2. Effect of low pH and novobiocin on *flhDC* transcription. The *flhDC*-*cat* transcriptional fusion activity was measured in exponential growth phase in M9 medium supplemented with 0.1% (w/v) Casamino acids and 0.4% (w/v) glycerol at pH 7 and pH 4.6 or in the presence of 200 μ M novobiocin. The data are the mean values from three independent assays. WT, wild-type; pDIA528, plasmid carrying the *flhDC* promoter; pDIA545, plasmid carrying the entire *flhDC* regulatory region.

neutral pH. It has been proposed that disintegration of flagella into subunits occurs at acidic pH (Stocker & Campbell, 1959; Weibull, 1948). However, to prevent unnecessary energy consumption – the cost to cell of flagellar synthesis is about 2% of total biosynthetic energy expenditure (Macnab, 1996) – it can be assumed that detrimental conditions such as acidic pH repress expression of flagellar genes. To test this hypothesis, we measured the expression level of the flagellin-encoding gene using a *flhC*-*cat* transcriptional fusion under neutral and acidic pH conditions. A sevenfold decrease in *flhC* transcription was observed at acidic as compared to neutral pH: CAT activities of 530 ± 30 and 68 ± 8 units were measured at pH 7.0 and pH 4.6, respectively (1 unit corresponds to 1 μ mol chloramphenicol acetylated per min per μ g protein).

Regulation of the *flhDC* expression by low pH

The *flhDC* master operon, located at the top of the flagellum biosynthesis cascade, constitutes the major target for regulatory proteins, such as H-NS and cAMP-CAP (Soutourina *et al.*, 1999). To determine whether the control by acidic pH affected bacterial motility by downregulating the expression of the master operon, the motility of an *E. coli* wild-type strain overexpressing the *flhDC* operon (Table 1) was assayed under low-pH conditions. FlhDC overproduction from plasmid pPM61 resulted in a partial restoration of the motility defect (data not shown), suggesting that low pH affects the motility via the master regulator, as observed with some other environmental factors (Shi *et al.*, 1993b). This hypothesis was tested with CAT transcriptional fusions carrying either the *flhDC* promoter region alone or the extended regulatory region that includes the mRNA untranslated 5' end (Fig. 2). Our previous studies (Soutourina *et al.*, 1999) have demonstrated a crucial

role for the *flhDC* 5' end in the positive control of the flagellar master operon by H-NS. In the wild-type strain, similar values were obtained under neutral and low-pH conditions from the transcriptional fusion carrying the promoter region alone on plasmid pDIA528. In contrast, a more than threefold decrease in CAT activity was measured under acidic pH in comparison with neutral pH from the transcriptional fusion carrying the entire *flhDC* regulatory region on plasmid pDIA545. A similar reduction in *flhDC* expression has been previously measured in *E. coli* strains subjected to various stresses, such as high osmolarity or increased temperature (Shi *et al.*, 1993b), supporting a role for the *flhDC* 5' end in the control of flagellar genes in response to acidic pH.

Both low pH and *hns* mutation resulted in an up to threefold decrease in *flhDC*-*cat* activity in the presence of the 5' end region. Such a reduction in *flhDC* expression has been previously observed in *hns* strains carrying a chromosomal *flhDC*-*lacZ* transcriptional fusion (Soutourina *et al.*, 1999). Moreover, a comparative analysis of the plasmid content revealed no difference between wild-type and *hns* strains (data not shown). This suggests that the reduced level of *cat* activity measured in the *hns* strain did not result from any effect of the mutation on the plasmid copy number of pKK232-8 derivatives, in accordance with the results obtained with plasmid pGR71 carrying the same origin of replication (Bertin *et al.*, 1992). More importantly, no further reduction in transcription was observed in the *hns* mutant under low pH (Fig. 2), indicating that the effects of low pH and H-NS deficiency are not additive. This suggests that the control of bacterial motility in response to acidic pH is mediated by the H-NS protein. These results are in accordance with the coordinate regulation of motility and low pH response by H-NS protein observed in suppressor mutants (Tables 2 and 3).

One possibility that could explain the role of H-NS in the control of gene expression is its implication in DNA topology (Higgins *et al.*, 1988; Dorman *et al.*, 2001). Moreover, some environmental factors or drugs that are known to inhibit bacterial motility also affect DNA supercoiling, e.g. high salt concentration or novobiocin (Anderson & Bauer, 1978; Goldstein & Drlica, 1984; Higgins *et al.*, 1988; Shi *et al.*, 1993b). The overexpression of DNA gyrase subunit GyrB resulted in a partial restoration of motility in an *hns* mutant (data not shown). Furthermore, in the presence of novobiocin, a DNA gyrase inhibitor decreasing DNA supercoiling, we observed a more than twofold decrease in *flhDC* activity from the *flhDC* transcriptional fusion containing the entire regulatory region, similar to that obtained in the presence of an *hns* mutation or at low pH (Fig. 2). Finally, as compared to the wild-type strain, which was non-motile in the presence of novobiocin (28 and 4 mm swarming ring diameter in the absence and presence of 200 μ M novobiocin, respectively), a significant restoration of motility was observed in the presence of plasmid pPM61 overexpressing the *flhDC* operon (10 mm swarming ring diameter).




Fig. 3. Effect of plasmid DNA supercoiling on *in vitro* *flhDC* transcription assays. Supercoiled (1) and linearized plasmid pDIA546 (2) were incubated with RNA polymerase. Samples were subjected to electrophoresis on 7% polyacrylamide sequencing gel. The *flhDC* transcript originating from the *flhDC* promoter (271 nt) and that from the RNA-I promoter located on the same plasmid (108 nt) used as a control are indicated by arrows.

To further investigate the effect of DNA topology on *flhDC* expression, we performed *in vitro* transcription experiments with either supercoiled or linearized plasmid pDIA546 carrying the entire *flhDC* regulatory region. A severe reduction in the *flhDC* transcription level was observed when linearized plasmid was used as a template (Fig. 3). This suggests that variations in DNA topology involving the 5' end of the *flhDC* operon may play an important role in the transcriptional control of the flagellar master operon by H-NS in response to low pH.

DISCUSSION

To date, only limited data are available on the motility control by acidic pH (Adler, 1973; Adler & Templeton, 1967; Bowra & Dilworth, 1981). The fact that bacterial flagella, with other extracellular organelles, are directly exposed to damage caused by acidic pH (Stocker & Campbell, 1959; Weibull, 1948) suggests that these structures might be disintegrated under these conditions. Nevertheless, our results provide evidence that acidic pH regulates flagellar gene expression in *E. coli* by a transcriptional control on the *flhDC* master operon

(Figs 1 and 2). These results are consistent with general energy consumption considerations under detrimental conditions, flagellum biosynthesis being extremely expensive for the cell (Macnab, 1996), and also with the necessity to close the proton entrance during flagellum motor functioning by reducing the biosynthesis of this structure. Moreover, we demonstrated here a remarkable parallelism in the regulation of motility by the H-NS protein and low pH (Tables 2 and 3, Fig. 2). In particular, we showed that the H-NS protein might play a central role in the pH-mediated control of bacterial flagellum biosynthesis (Fig. 2). The control of flagellar master operon expression in response to low pH (Fig. 2) or high osmolarity (O.S., unpublished) requires an extended *flhDC* mRNA 5' untranslated region. The comparative analysis of the upstream sequence of *flhDC* homologous operons in enterobacteria revealed the presence of a similar domain in many Gram-negative bacteria. Furthermore, a long 5' untranslated region has also been identified in the master flagellar regulatory gene of polarly flagellated bacteria of the genera *Vibrio* and *Pseudomonas* (Soutourina *et al.*, 2001). Taken together, these results suggest that the mechanism that controls expression of flagellar master regulatory genes in response to environmental factors may be largely conserved among Gram-negative bacteria.

Despite the similarities between *E. coli* and *S. typhimurium*, some differences have been reported in the regulation of motility in these two organisms. These include the autogenous and global control by H-NS on *flhDC* master operon expression (Kutsukake, 1997) and the initiation of *flhDC* transcription (Soutourina *et al.*, 1999; Yanagihara *et al.*, 1999). On the other hand, some specificities may also exist in the response of these bacteria to acidic pH (Lin *et al.*, 1995), including the PhoPQ-mediated acid tolerance response (ATR) (Bearson *et al.*, 1998). As the synthesis of H-NS protein has been shown to be unaffected by low pH (Adams *et al.*, 2001), it has been recently suggested that H-NS is not involved in the control of flagellar biosynthesis in *S. typhimurium* under the acidic conditions tested. Similarly, in *E. coli*, we observed no alteration in *hns* expression at acidic pH. However, *flhDC* transcription measurement (Fig. 2) and characterization of suppressive mutants (Table 3) demonstrated the role of H-NS protein in motility control under low-pH conditions in *E. coli*. Nevertheless, these observations do not exclude the possibility that other regulators may participate in this process.

The presence of a 5' untranslated mRNA region has been usually associated with post-transcriptional regulation mechanisms. For example, such regions are responsible for either transcriptional anti-termination and RNA processing or translational control of threonyl-tRNA synthetase genes in *E. coli* and *Bacillus subtilis* (Condon *et al.*, 1997; Putzer *et al.*, 1995; Sacerdot *et al.*, 1998). As compared to the wild-type strain, an up to threefold decrease in *flhDC* activity was observed in an *hns* strain from both transcriptional and translational fusions (Soutourina *et al.*, 1999). More-

over, a similar reduction in the level of *flhDC* mRNA was measured in the *hns* mutant as compared to the wild-type in RT-PCR experiments and this reduction did not result from an effect of the H-NS protein on the *flhDC* mRNA stability (Soutourina, 2001). Finally, *in vivo* activity measurements from a transcriptional fusion containing the extended *flhDC* regulatory region (Fig. 2) demonstrated that H-NS, acidic pH and DNA supercoiling affect the *flhDC* expression at the level of transcription initiation.

Despite numerous studies on the control of bacterial motility by environmental factors, the molecular basis of this process remains largely unknown. In *E. coli*, the H-NS protein affects the expression of many genes involved in the cellular response to environmental changes, including those required for acidic pH resistance (Hommais *et al.*, 2001). Although the mechanism by which H-NS controls gene expression remains the subject of debate (Williams & Rimsky, 1997), an alteration of plasmid and chromosomal DNA supercoiling has been demonstrated *in vivo* in an *hns* mutant (Mojica & Higgins, 1997). Moreover, the involvement of DNA supercoiling has been proposed, for example, to explain the regulation by H-NS of osmotically regulated genes (Higgins *et al.*, 1988), stringently controlled bacterial promoters (Johansson *et al.*, 2000) or virulence gene expression in *Shigella flexneri* (Dorman *et al.*, 2001). On the other hand, various environmental conditions are also known to affect the level of DNA supercoiling, even though a cause-and-effect relationship has not yet been established (Higgins *et al.*, 1988; Tse-Dinh *et al.*, 1997). It has been proposed that a direct effect of environmental signals on promoter architecture, and then transcription, through influencing the interaction of architectural proteins with DNA, might be an important concept in understanding the environmental regulation of gene expression in bacteria (Jordi *et al.*, 1997). Similarly, different environmental cues might influence the action of H-NS by changing the structure of regulatory regions, the ability of H-NS to bind to DNA target and/or the conformation or the oligomerization state of H-NS. We did not observe any alteration in the level of *hns* gene expression or in the isoform composition of the H-NS protein under low pH or in the presence of DNA gyrase inhibitor (data not shown). In contrast, *flhDC* expression could be modulated by local alteration of DNA topology, resulting from interactions between H-NS and the regulatory region. Several observations argue in favour of this hypothesis. First, the alteration of swarming properties in presence of novobiocin, a DNA gyrase inhibitor, or in strains overproducing DNA gyrase or mutated in its structural gene suggests the existence of a critical DNA supercoiling level for normal motility in *E. coli* (see Results) (Shi *et al.*, 1993b; O.S., unpublished). Second, we observed a severe reduction in *flhDC* expression when linearized rather than supercoiled plasmid was used in *in vitro* transcription assay (Fig. 3). Third, the involvement of the extended *flhDC* regulatory region in the control of the master operon by H-NS, acidic pH

and novobiocin suggests a strong correlation between these regulatory processes (Fig. 2). Finally, further support is provided by the partial restoration of motility in the *hns* mutant by overexpression of DNA gyrase subunit gene *gyrB* (data not shown), and the alteration of topoisomer distribution of plasmids carrying the entire *flhDC* regulatory region in the presence of chloroquine phosphate as an intercalating agent (Soutourina, 2001). Taken together, these data suggest that the H-NS-mediated effect on motility may be at least in part explained by an alteration in the level of DNA topology of the *flhDC* regulatory region. They are consistent with the recent demonstration that a 339 bp DNA fragment having a bent structure can strongly affect the level of plasmid DNA supercoiling (Rohde *et al.*, 1999) and suggest that the regulatory region of the flagellar master operon may play a crucial role for an adequate control by the H-NS protein and environmental factors.

The control of bacterial motility via the *flhDC* operon includes several participants at multiple regulatory levels, e.g. transcription initiation control by cAMP-CAP complex and H-NS (Soutourina *et al.*, 1999), mRNA stability control by CsrA (Wei *et al.*, 2001) in *E. coli*, or FlhDC protein degradation by Lon protease in *Proteus mirabilis* (Claret & Hughes, 2000). Our results extend the knowledge of the regulation of the flagellar system and represent an important step toward the understanding of complex mechanisms governing bacterial motility in response to environmental challenges.

ACKNOWLEDGEMENTS

We are grateful to A. Pugsley for helpful advice and discussions and to G. Karimova for critical reading of the manuscript. We thank A. Kolb for technical assistance in *in vitro* transcription experiments.

Financial support came from the Institut Pasteur and the Centre National de la Recherche Scientifique (URA 2171). O.S. was supported by a French Government fellowship.

REFERENCES

- Adams, P., Fowler, R., Kinsella, N., Howell, G., Farris, M., Coote, P. & O'Connor, C. D. (2001). Proteomic detection of PhoPQ- and acid-mediated repression of *Salmonella* motility. *Proteomics* **1**, 597–607.
- Adler, J. (1973). A method for measuring chemotaxis and use of the method to determine optimum conditions for chemotaxis by *Escherichia coli*. *J Gen Microbiol* **74**, 77–91.
- Adler, J. & Templeton, B. (1967). The effect of environmental conditions on the motility of *Escherichia coli*. *J Gen Microbiol* **46**, 175–184.
- Aizawa, S.-I. & Kubori, T. (1998). Bacterial flagellation and cell division. *Genes Cells* **3**, 625–634.
- Anderson, P. & Bauer, W. (1978). Supercoiling in closed circular DNA: dependence upon ion type and concentration. *Biochemistry* **17**, 594–601.
- Arnold, H. H. & Winter, B. (1998). Muscle differentiation: more complexity to the network of myogenic regulators. *Curr Opin Genet Dev* **8**, 539–544.

- Bartlett, D. H., Frantz, B. B. & Matsumura, P. (1988). Flagellar transcriptional activators FlbB and FlaI: gene sequences and 5' consensus sequences of operons under FlbB and FlaI control. *J Bacteriol* **170**, 1575–1581.
- Bearson, B., Wilson, L. & Foster, J. (1998). A low pH-inducible, PhoQP-dependent acid tolerance response protects *Salmonella typhimurium* against inorganic acid stress. *J Bacteriol* **180**, 2409–2417.
- Bertin, P., Lejeune, P., Colson, C. & Danchin, A. (1992). Mutations in *bglY*, the structural gene for the DNA-binding protein H1 of *Escherichia coli*, increase the expression of the kanamycin resistance gene carried by plasmid pGR71. *Mol Gen Genet* **233**, 184–192.
- Bertin, P., Terao, E., Lee, E. H., Lejeune, P., Colson, C., Danchin, A. & Collatz, E. (1994). The H-NS protein is involved in the biogenesis of flagella in *Escherichia coli*. *J Bacteriol* **176**, 5537–5540.
- Bertin, P., Benhabiles, N., Krin, E., Laurent-Winter, C., Tendeng, C., Turlin, E., Thomas, A., Danchin, A. & Brasseur, R. (1999). The structural and functional organization of H-NS-like proteins is evolutionarily conserved in Gram-negative bacteria. *Mol Microbiol* **31**, 319–329.
- Bertin, P., Hommais, F., Krin, E., Soutourina, O., Tendeng, C., Derzelle, S. & Danchin, A. (2001). H-NS and H-NS-like proteins in Gram-negative bacteria and their multiple role in the regulation of bacterial metabolism. *Biochimie* **83**, 235–241.
- Bowra, B. J. & Dilworth, M. J. (1981). Motility and chemotaxis towards sugars in *Rhizobium leguminosarum*. *J Gen Microbiol* **126**, 231–235.
- Bruni, C. B., Colantuoni, V., Sbordone, L., Cortese, R. & Blasi, F. (1977). Biochemical and regulatory properties of *Escherichia coli* K-12 *his* mutants. *J Bacteriol* **130**, 4–10.
- Claret, L. & Hughes, C. (2000). Rapid turnover of FlhD and FlhC, the flagellar regulon transcriptional activator proteins, during *Proteus* swarming. *J Bacteriol* **182**, 833–836.
- Condon, C., Putzer, H., Luo, D. & Grunberg-Manago, M. (1997). Processing of the *Bacillus subtilis* *thrS* leader mRNA is RNase E-dependent in *Escherichia coli*. *J Mol Biol* **268**, 235–242.
- Dorman, C. J., McKenna, S. & Beloin, C. (2001). Regulation of intralocus gene expression in *Shigella flexneri*, a facultative intracellular pathogen. *Int J Med Microbiol* **291**, 89–96.
- Farr, S. B., Arnosti, D. N., Chamberlin, M. J. & Ames, B. (1989). An *apaH* mutation causes AppppA to accumulate and affects motility and catabolite repression in *Escherichia coli*. *Proc Natl Acad Sci USA* **86**, 5010–5014.
- Goldstein, E. & Drlica, K. (1984). Regulation of bacterial DNA supercoiling: plasmid linking numbers vary with growth temperature. *Proc Natl Acad Sci USA* **81**, 4046–4050.
- Hengge-Aronis, R. (1999). Interplay of global regulators and cell physiology in the general stress response of *Escherichia coli*. *Curr Opin Microbiol* **2**, 148–152.
- Higgins, C. F., Dorman, C. J., Stirling, D. A., Waddell, L., Booth, I. R., May, G. & Bremer, E. (1988). A physiological role for DNA supercoiling in the osmotic regulation of gene expression in *S. typhimurium* and *E. coli*. *Cell* **52**, 569–584.
- Hinton, J. C. D., Santos, D. S., Seirafi, A., Hulton, C. J., Pavitt, G. D. & Higgins, C. F. (1992). Expression and mutational analysis of the nucleoid-associated protein H-NS of *Salmonella typhimurium*. *Mol Microbiol* **6**, 2327–2337.
- Hommais, F., Krin, E., Laurent-Winter, C., Soutourina, O., Malpertuy, A., Le Caer, J. P., Danchin, A. & Bertin, P. (2001). Large-scale monitoring of pleiotropic regulation of gene expression by the prokaryotic nucleoid-associated protein, H-NS. *Mol Microbiol* **40**, 20–36.
- Huang, S. (1999). Gene expression profiling, genetic networks, and cellular states: an integrating concept for tumorigenesis and drug discovery. *J Mol Med* **77**, 469–480.
- Johansson, J., Balsalobre, C., Wang, S. Y., Urbonaviciene, J., Jin, D. J., Sonden, B. & Uhlin, B. E. (2000). Nucleoid proteins stimulate stringently controlled bacterial promoters: a link between the cAMP-CRP and the (p)ppGpp regulons in *Escherichia coli*. *Cell* **102**, 475–485.
- Jordi, B. J. A. M., Fielder, A., Burns, C. M., Hinton, J. C. D., Dover, N., Ussery, D. W. & Higgins, C. F. (1997). DNA binding is not sufficient for H-NS-mediated repression of *proU* expression. *J Biol Chem* **272**, 12083–12090.
- Kawakami, K., Sato, S., Ozaki, H. & Ikeda, K. (2000). Six family genes: structure and function as transcription factors and their roles in development. *Bioessays* **22**, 616–626.
- Kitamura, E., Nakayama, Y., Matsuzaki, H., Matsumoto, K. & Shibuya, I. (1994). Acidic-phospholipid deficiency represses the flagellar master operon through a novel regulatory region in *Escherichia coli*. *Biosci Biotechnol Biochem* **58**, 2305–2307.
- Ko, M. & Park, C. (2000). H-NS-dependent regulation of flagellar synthesis is mediated by a LysR family protein. *J Bacteriol* **182**, 4670–4672.
- Kutsukake, K. (1997). Autogenous and global control of the flagellar master operon, *flbD*, in *Salmonella typhimurium*. *Mol Gen Genet* **254**, 440–448.
- Laurent-Winter, C., Ngo, S., Danchin, A. & Bertin, P. (1997). Role of *Escherichia coli* histone-like nucleoid-structuring protein in bacterial metabolism and stress response. *Eur J Biochem* **244**, 767–773.
- Lejeune, P. & Danchin, A. (1990). Mutations in the *bglY* gene increase the frequency of spontaneous deletion in *Escherichia coli* K-12. *Proc Natl Acad Sci USA* **87**, 360–363.
- Li, C., Louise, C. J., Shi, W. & Adler, J. (1993). Adverse conditions which cause lack of flagella in *Escherichia coli*. *J Bacteriol* **175**, 2229–2235.
- Lin, J., Lee, I. S., Frey, J., Slonczewski, J. L. & Foster, J. W. (1995). Comparative analysis of extreme acid survival in *Salmonella typhimurium*, *Shigella flexneri*, and *Escherichia coli*. *J Bacteriol* **177**, 4097–4104.
- Macnab, R. M. (1996). Flagella and motility. In *Escherichia coli and Salmonella typhimurium: Cellular and Molecular Biology*, pp. 123–145. Edited by F. C. Neidhardt and others. Washington, DC: American Society for Microbiology.
- Mahan, M. J., Slauch, J. M. & Mekalanos, J. J. (1996). Environmental regulation of virulence gene expression in *Escherichia*, *Salmonella*, and *Shigella* spp. In *Escherichia coli and Salmonella typhimurium: Cellular and Molecular Biology*, pp. 2803–2815. Edited by F. C. Neidhardt and others. Washington, DC: American Society for Microbiology.
- Miller, J. H. (1992). *A Short Course in Bacterial Genetics*. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory.
- Mizushima, T., Koyanagi, R., Suzuki, E., Tomura, A., Kutsukake, K., Miki, T. & Sekimizu, K. (1995). Control by phosphatidylglycerol of expression of the *flbD* gene in *Escherichia coli*. *Biochim Biophys Acta* **1245**, 397–401.
- Mizushima, T., Koyanagi, R., Katayama, T., Miki, T. & Sekimizu, K. (1997). Decrease in expression of the master operon of flagellin synthesis in a *dnaA46* mutant of *Escherichia coli*. *Biol Pharm Bull* **20**, 327–331.

- Mojica, F. J. M. & Higgins, C. F. (1997).** *In vivo* supercoiling of plasmid and chromosomal DNA in an *Escherichia coli* *hns* mutant. *J Bacteriol* **179**, 3528–3533.
- Putzer, H., Grunberg-Manago, M. & Springer, M. (1995).** Bacterial aminoacyl-tRNA synthetases: genes and regulation of expression. In *tRNA: Structure, Biosynthesis and Function*, pp. 293–333. Edited by D. Soll & U. L. RajBhandary. Washington, DC: American Society for Microbiology.
- Relaix, F. & Buckingham, M. (1999).** From insect eye to vertebrate muscle: redeployment of a regulatory network. *Genes Dev* **13**, 3171–3178.
- Rohde, J. R., Fox, J. M. & Minnich, S. A. (1994).** Thermoregulation in *Yersinia enterocolitica* is coincident with changes in DNA supercoiling. *Mol Microbiol* **12**, 187–199.
- Rohde, J. R., Luan, X. S., Rohde, H., Fox, J. M. & Minnich, S. A. (1999).** The *Yersinia enterocolitica* pYV virulence plasmid contains multiple intrinsic DNA bends which melt at 37 °C. *J Bacteriol* **181**, 4198–4204.
- Romeo, T. (1998).** Global regulation by the small RNA-binding protein CsrA and the non-coding RNA molecule CsrB. *Mol Microbiol* **29**, 1321–1330.
- Sacerdot, C., Caillet, J., Graffe, M., Eyermann, F., Ehresmann, B., Ehresmann, C., Springer, M. & Romby, P. (1998).** The *Escherichia coli* threonyl-tRNA synthetase gene contains a split ribosomal binding site interrupted by a hairpin structure that is essential for autoregulation. *Mol Microbiol* **29**, 1077–1090.
- Shi, W., Zhou, Y., Wild, J., Adler, J. & Gross, C. A. (1992).** DnaK, DnaJ, and GrpE are required for flagellum synthesis in *Escherichia coli*. *J Bacteriol* **174**, 6256–6263.
- Shi, W., Bogdanov, M., Dowhan, W. & Zusman, D. R. (1993a).** The *pss* and *psd* genes are required for motility and chemotaxis in *Escherichia coli*. *J Bacteriol* **175**, 7711–7714.
- Shi, W., Li, C., Louise, C. & Adler, J. (1993b).** Mechanism of adverse conditions causing lack of flagella in *Escherichia coli*. *J Bacteriol* **175**, 2236–2240.
- Silverman, M. & Simon, M. (1974).** Characterization of *Escherichia coli* flagellar mutants that are insensitive to catabolite repression. *J Bacteriol* **120**, 1196–1203.
- Soutourina, O. (2001).** *Control of gene expression in the motility process in Gram-negative bacteria*. PhD thesis, University of Versailles-Saint-Quentin.
- Soutourina, O., Kolb, A., Krin, E., Laurent-Winter, C., Rimsky, S., Danchin, A. & Bertin, P. (1999).** Multiple control of flagellum biosynthesis in *Escherichia coli*: role of H-NS protein and the cyclic AMP-catabolite activator protein complex in transcription of the *flhDC* master operon. *J Bacteriol* **181**, 7500–7508.
- Soutourina, O. A., Semenova, E. A., Parfenova, V. V., Danchin, A. & Bertin, P. (2001).** Control of bacterial motility by environmental factors in polarly flagellated and peritrichous bacteria isolated from Lake Baikal. *Appl Environ Microbiol* **67**, 3852–3859.
- Stocker, B. A. D. & Campbell, J. C. (1959).** The effect of non-lethal deflagellation on bacterial motility and observations on flagellar regeneration. *J Gen Microbiol* **20**, 670–685.
- Tse-Dinh, Y. C., Qi, H. & Menzel, R. (1997).** DNA supercoiling and bacterial adaptation: thermotolerance and thermoresistance. *Trends Microbiol* **5**, 323–326.
- Wei, B. L., Brun-Zinkernagel, A.-M., Simecka, J. W., Pruss, B. M., Babitzke, P. & Romeo, T. (2001).** Positive regulation of motility and *flhDC* expression by the RNA-binding protein CsrA of *Escherichia coli*. *Mol Microbiol* **40**, 245–256.
- Weibull, C. (1948).** Some chemical and physico-chemical properties of the flagella of *Proteus vulgaris*. *Biochim Biophys Acta* **2**, 351–355.
- Williams, R. M. & Rimsky, S. (1997).** Molecular aspects of the *Escherichia coli* nucleoid protein, H-NS: a central controller of gene regulatory networks. *FEMS Microbiol Lett* **156**, 175–185.
- Yanagihara, S., Iyoda, S., Ohnishi, K., Iino, T. & Kutsukake, K. (1999).** Structure and transcriptional control of the flagellar master operon of *Salmonella typhimurium*. *Genes Genet Syst* **74**, 105–111.
- Yokota, T. & Gots, J. (1970).** Requirement of adenosine 3',5'-cyclic monophosphate for flagellation in *Escherichia coli* and *Salmonella typhimurium*. *J Bacteriol* **103**, 513–516.

Received 1 November 2001; revised 9 January 2002; accepted 17 January 2002.