

# Higher algebra of $A_\infty$ -algebras in Morse theory

Thibaut Mazuir

IMJ-PRG - Sorbonne Université

Symplectic Seminar, Heidelberg University, 12/01/2022

The results presented in this talk are taken from my two recent papers : *Higher algebra of  $A_\infty$  and  $\Omega BAs$ -algebras in Morse theory I* (arXiv:2102.06654) and *Higher algebra of  $A_\infty$  and  $\Omega BAs$ -algebras in Morse theory II* (arxiv:2102.08996).

- 1 The  $A_\infty$ -algebra structure on the Morse cochains
- 2  $A_\infty$ -morphisms between the Morse cochains
- 3 Higher morphisms between  $A_\infty$ -algebras ...
- 4 ... and their realization in Morse theory
- 5 Further directions

- 1 The  $A_\infty$ -algebra structure on the Morse cochains
  - $A_\infty$ -algebras
  - The associahedra
  - $A_\infty$ -algebra structure on the Morse cochains
- 2  $A_\infty$ -morphisms between the Morse cochains
- 3 Higher morphisms between  $A_\infty$ -algebras ...
- 4 ... and their realization in Morse theory
- 5 Further directions

## Definition

Let  $A$  be a cochain complex with differential  $m_1$ . An  $A_\infty$ -algebra structure on  $A$  is the data of a collection of maps of degree  $2 - n$

$$m_n : A^{\otimes n} \longrightarrow A, \quad n \geq 1,$$


extending  $m_1$  and which satisfy

$$[m_1, m_n] = \sum_{\substack{i_1+i_2+i_3=n \\ 2 \leq i_2 \leq n-1}} \pm m_{i_1+1+i_3}(\text{id}^{\otimes i_1} \otimes m_{i_2} \otimes \text{id}^{\otimes i_3}).$$

These equations are called the  $A_\infty$ -equations.

Recall for instance that for  $n = 2$ ,

$$[m_1, m_2] := m_1 m_2 - m_2(\text{id} \otimes m_1) - m_2(m_1 \otimes \text{id}) .$$

Representing  $m_n$  as , these equations can be written as

$$[m_1, \text{tree}(1, 2, \dots, n)] = \sum_{\substack{h+k=n+1 \\ 2 \leq h \leq n-1 \\ 1 \leq i \leq k}} \pm 1 \cdot \text{tree}(1, \dots, k, d_1, d_2) .$$

In particular,

$$[m_1, m_2] = 0 ,$$

$$[m_1, m_3] = m_2(\text{id} \otimes m_2 - m_2 \otimes \text{id}) ,$$

implying that  $m_2$  descends to an associative product on  $H^*(A)$ . An  $A_\infty$ -algebra is thus simply a correct notion of a dg-algebra whose product is associative up to homotopy.

The operations  $m_n$  are the higher coherent homotopies which keep track of the fact that the product is associative up to homotopy.

## Theorem (Homotopy transfer theorem)

Let  $(A, \partial_A)$  and  $(H, \partial_H)$  be two cochain complexes. Suppose that  $H$  is a deformation retract of  $A$ , that is that they fit into a diagram

$$h \begin{array}{c} \curvearrowright \end{array} (A, \partial_A) \begin{array}{c} \xrightarrow{p} \\ \xleftarrow{i} \end{array} (H, \partial_H) ,$$

where  $\text{id}_A - ip = [\partial, h]$ . Then if  $(A, \partial_A)$  is endowed with an  $A_\infty$ -algebra structure,  $H$  can be made into an  $A_\infty$ -algebra such that  $i$  and  $p$  extend to  $A_\infty$ -morphisms.



- 1 The  $A_\infty$ -algebra structure on the Morse cochains
  - $A_\infty$ -algebras
  - The associahedra
  - $A_\infty$ -algebra structure on the Morse cochains
- 2  $A_\infty$ -morphisms between the Morse cochains
- 3 Higher morphisms between  $A_\infty$ -algebras ...
- 4 ... and their realization in Morse theory
- 5 Further directions


There exists a collection of polytopes, called the *associahedra* and denoted  $\{K_n\}$ , which encode the  $A_\infty$ -equations between  $A_\infty$ -algebras. This means that  $K_n$  has a unique cell  $[K_n]$  of dimension  $n - 2$  and that its boundary reads as

$$\partial K_n = \bigcup_{\substack{h+k=n+1 \\ 2 \leq h \leq n-1}} \bigcup_{1 \leq i \leq k} K_k \times_i K_h ,$$

where  $\times_i$  is in fact the standard  $\times$  cartesian product.

Recall that the  $A_\infty$ -equations read as

$$[m_1, \text{tree}(1, 2, \dots, n)] = \sum_{\substack{h+k=n+1 \\ 2 \leq h \leq n-1 \\ 1 \leq i \leq k}} \pm 1 \cdot \text{tree}(1, \dots, k, d_1, d_2) .$$



**Figure:** The associahedra  $K_2$ ,  $K_3$  and  $K_4$ , with cells labeled by the operations they define

- 1 The  $A_\infty$ -algebra structure on the Morse cochains
  - $A_\infty$ -algebras
  - The associahedra
  - $A_\infty$ -algebra structure on the Morse cochains
- 2  $A_\infty$ -morphisms between the Morse cochains
- 3 Higher morphisms between  $A_\infty$ -algebras ...
- 4 ... and their realization in Morse theory
- 5 Further directions

Let  $M$  be an oriented closed Riemannian manifold endowed with a Morse function  $f$  together with a Morse-Smale metric. The Morse cochains  $C^*(f)$  form a deformation retract of the singular cochains  $C_{sing}^*(M)$  as shown in [Hut08].

$$h \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} (C_{sing}^*, \partial_{sing}) \begin{array}{c} \xrightarrow{p} \\ \xleftarrow{i} \end{array} (C^*(f), \partial_{Morse}) .$$

The cup product naturally endows the singular cochains  $C_{\text{sing}}^*(M)$  with a dg-algebra structure. The homotopy transfer theorem ensures that it can be transferred to an  $A_\infty$ -algebra structure on the Morse cochains  $C^*(f)$ .

The differential on the Morse cochains is defined by a count of moduli spaces of gradient trajectories. Is it then possible to define higher multiplications  $m_n$  on  $C^*(f)$  by a count of moduli spaces such that they fit in a structure of  $A_\infty$ -algebra ?


Question solved for the first time by Abouzaid in [Abo11], drawing from earlier works by Fukaya ([Fuk97] for instance). See also [Mes18] and [AL18].




## Terminology :



A ribbon tree



A metric ribbon tree



A stable metric ribbon tree

## Definition

Define  $\mathcal{T}_n$  to be *moduli space of stable metric ribbon trees with  $n$  incoming edges*. For each stable ribbon tree type  $t$ , we define moreover  $\mathcal{T}_n(t) \subset \mathcal{T}_n$  to be the moduli space

$$\mathcal{T}_n(t) := \{\text{stable metric ribbon trees of type } t\} .$$

We then have the following cell decomposition

$$\mathcal{T}_n = \bigcup_{t \in SRT_n} \mathcal{T}_n(t) .$$

Allowing lengths of internal edges to go to  $+\infty$ , this moduli space can be compactified into a  $(n-2)$ -dimensional CW-complex  $\overline{\mathcal{T}}_n$ , where  $\mathcal{T}_n$  is seen as its unique  $(n-2)$ -dimensional stratum.

### Theorem

*The compactified moduli space  $\overline{\mathcal{T}}_n$  is isomorphic as a CW-complex to the associahedron  $K_n$ .*

This was first noticed in section 1.4. of Boardman-Vogt [BV73].




Figure: The compactified moduli space  $\overline{\mathcal{T}}_3$





Figure: The compactified moduli space  $\overline{\mathcal{T}}_4$

The goal is now to realize these moduli spaces of stable metric ribbon trees in Morse theory.



## Definition

$T := (t, \{l_e\}_{e \in E(t)})$  where  $\{l_e\}_{e \in E(t)}$  are the lengths of its internal edges of the tree  $t$ . *Choice of perturbation data* on  $T$  consists of the following data :

- (i) a vector field  $[0, l_e] \times M \xrightarrow{\mathbb{X}_e} TM$ , that vanishes on  $[1, l_e - 1]$ , for every internal edge  $e$  of  $t$  ;
- (ii) a vector field  $[0, +\infty[ \times M \xrightarrow{\mathbb{X}_{e_0}} TM$ , that vanishes away from  $[0, 1]$ , for the outgoing edge  $e_0$  of  $t$  ;
- (iii) a vector field  $] - \infty, 0] \times M \xrightarrow{\mathbb{X}_{e_i}} TM$ , that vanishes away from  $[-1, 0]$ , for every incoming edge  $e_i$  ( $1 \leq i \leq n$ ) of  $t$ .

We will write  $D_e$  for all segments  $[0, l_e]$  as well as for all semi-infinite segments  $] - \infty, 0]$  and  $[0, +\infty[$  in the rest of the talk.

### Definition ([Abo11])

A *perturbed Morse gradient tree*  $T^{Morse}$  associated to  $(T, \mathbb{X})$  is the data for each edge  $e$  of  $t$  of a smooth map  $\gamma_e : D_e \rightarrow M$  such that  $\gamma_e$  is a trajectory of the perturbed negative gradient  $-\nabla f + \mathbb{X}_e$ , i.e.

$$\dot{\gamma}_e(s) = -\nabla f(\gamma_e(s)) + \mathbb{X}_e(s, \gamma_e(s)) ,$$

and such that the endpoints of these trajectories coincide as prescribed by the edges of the tree  $T$ .



## Definition

Let  $\mathbb{X}_n$  be a smooth choice of perturbation data on  $\mathcal{T}_n$ . For critical points  $y$  and  $x_1, \dots, x_n$ , we define the moduli space

$$\mathcal{T}_n^{\mathbb{X}_n}(y; x_1, \dots, x_n) := \left\{ \begin{array}{l} \text{perturbed Morse gradient trees associated to } (T, \mathbb{X}_T) \\ \text{and connecting } x_1, \dots, x_n \text{ to } y, \text{ for } T \in \mathcal{T}_n \end{array} \right\}.$$

## Proposition

*Given a generic choice of perturbation data  $\mathbb{X}_n$ , the moduli space  $\mathcal{T}_n^{\mathbb{X}_n}(y; x_1, \dots, x_n)$  is an orientable manifold of dimension*

$$\dim(\mathcal{T}_n(y; x_1, \dots, x_n)) = n - 2 + |y| - \sum_{i=1}^n |x_i| ,$$

where  $|x| := \dim(W^S(x))$ .



Choose perturbation data  $\mathbb{X}_n$  on each moduli space  $\mathcal{T}_n$  for  $n \geq 2$ .  
 By assuming some gluing-compatibility conditions on  $(\mathbb{X}_n)_{n \geq 2}$ , the  
 1-dimensional moduli spaces  $\mathcal{T}_n(y; x_1, \dots, x_n)$  can be compactified  
 to manifolds with boundary whose boundary is given by the spaces

(i) corresponding to an internal edge breaking :

$$\mathcal{T}_{i_1+1+i_3}^{\mathbb{X}_{i_1+1+i_3}}(y; x_1, \dots, x_{i_1}, z, x_{i_1+i_2+1}, \dots, x_n) \times \mathcal{T}_{i_2}^{\mathbb{X}_{i_2}}(z; x_{i_1+1}, \dots, x_{i_1+i_2});$$

(ii) corresponding to an external edge breaking :

$$\mathcal{T}(y; z) \times \mathcal{T}_n^{\mathbb{X}_n}(z; x_1, \dots, x_n) \quad \text{and} \quad \mathcal{T}_n^{\mathbb{X}_n}(y; x_1, \dots, z, \dots, x_n) \times \mathcal{T}(z; x_i)$$



Two examples of perturbed Morse gradient trees breaking at a critical point

## Theorem ([Abo11])

For an admissible choice of perturbation data  $\mathbb{X} := (\mathbb{X}_n)_{n \geq 2}$ ,  
 defining for every  $n$  the operation  $m_n$  as

$$m_n : C^*(f) \otimes \cdots \otimes C^*(f) \longrightarrow C^*(f)$$

$$x_1 \otimes \cdots \otimes x_n \longmapsto \sum_{|y| = \sum_{i=1}^n |x_i| + 2 - n} \# \mathcal{T}_n^{\mathbb{X}}(y; x_1, \dots, x_n) \cdot y,$$

they endow the Morse cochains  $C^*(f)$  with an  $A_\infty$ -algebra structure.

Indeed, the boundary of the previous compactification is modeled on the  $A_\infty$ -equations for  $A_\infty$ -algebras :

$$[\partial_{Morse}, \text{tree}(1, 2, \dots, n)] = \sum_{\substack{h+k=n+1 \\ 2 \leq h \leq n-1 \\ 1 \leq i \leq k}} \pm \text{tree}(1, \dots, h, \text{tree}(1, \dots, k, d_1), d_2) .$$

In fact, we construct in [Maz21a] a thinner algebraic structure on  $C^*(f)$ , called an  $\Omega BAs$ -algebra structure. It corresponds to associating to each stable ribbon tree  $t$  of arity  $n$  an operation  $C^*(f)^{\otimes n} \rightarrow C^*(f)$ , and the differential for such an operation is then encoded by the codimension 1 boundary of the corresponding cell in  $\mathcal{T}_n$ . For instance,

$$\partial(\text{Y-shape}) = \pm \text{Y-shape with top-left Y} \pm \text{Y-shape with top-right Y} \pm \text{Y-shape with bottom-left Y} \pm \text{Y-shape with bottom-right Y}.$$

The  $A_\infty$ -algebra structure on the Morse cochains then stems from this  $\Omega BAs$ -algebra structure by purely algebraic arguments.

Working on the  $\Omega BAs$  and not on the  $A_\infty$  level is also more rigorous for the analysis involved in these constructions.




- 1 The  $A_\infty$ -algebra structure on the Morse cochains
- 2  $A_\infty$ -morphisms between the Morse cochains
  - $A_\infty$ -morphisms
  - The multiplihedra
  - $A_\infty$ -morphisms between the Morse cochains
- 3 Higher morphisms between  $A_\infty$ -algebras ...
- 4 ... and their realization in Morse theory
- 5 Further directions

## Definition

An  $A_\infty$ -morphism between two  $A_\infty$ -algebras  $A$  and  $B$  is a family of maps  $f_n : A^{\otimes n} \rightarrow B$  of degree  $1 - n$  satisfying

$$\begin{aligned}
 [m_1, f_n] = & \sum_{\substack{i_1+i_2+i_3=n \\ i_2 \geq 2}} \pm f_{i_1+1+i_3}(\text{id}^{\otimes i_1} \otimes m_{i_2} \otimes \text{id}^{\otimes i_3}) \\
 & + \sum_{\substack{i_1+\dots+i_s=n \\ s \geq 2}} \pm m_s(f_{i_1} \otimes \dots \otimes f_{i_s}) .
 \end{aligned}$$

Representing the operations  $f_n$  as , the operations  $m_n^B$  in red and the operations  $m_n^A$  in blue, these equations read as

$$\left[ \partial, \text{tree} \right] = \sum_{\substack{h+k=n+1 \\ 1 \leq i \leq k \\ h \geq 2}} \pm \text{tree}_1 + \sum_{\substack{i_1 + \dots + i_s = n \\ s \geq 2}} \pm \text{tree}_2 .$$

The first tree diagram in the sum has a red root with a blue child labeled  $i$  and a red child labeled  $k$ . The second tree diagram has a red root with  $s$  children, each being a tree with a red root and blue children. The children are labeled  $i_1, \dots, i_s$ .

We check that  $[\partial, f_2] = f_1 m_2^A - m_2^B(f_1 \otimes f_1)$ .

An  $A_\infty$ -morphism between  $A_\infty$ -algebras induces a morphism of associative algebras on the level of cohomology, and is a correct notion of morphism which preserves the product up to homotopy.

- 1 The  $A_\infty$ -algebra structure on the Morse cochains
- 2  $A_\infty$ -morphisms between the Morse cochains
  - $A_\infty$ -morphisms
  - **The multiplihedra**
  - $A_\infty$ -morphisms between the Morse cochains
- 3 Higher morphisms between  $A_\infty$ -algebras ...
- 4 ... and their realization in Morse theory
- 5 Further directions

There exists a collection of polytopes, called the *multiplihedra* and denoted  $\{J_n\}$ , which encode the  $A_\infty$ -equations for  $A_\infty$ -morphisms. Again,  $J_n$  has a unique  $n - 1$ -dimensional cell  $[J_n]$  and the boundary of  $J_n$  is exactly

$$\partial J_n = \bigcup_{\substack{h+k=n+1 \\ h \geq 2}} \bigcup_{1 \leq i \leq k} J_k \times_i K_h \cup \bigcup_{\substack{i_1 + \dots + i_s = n \\ s \geq 2}} K_s \times J_{i_1} \times \dots \times J_{i_s} ,$$

where  $\times_k$  is the standard cartesian product  $\times$ .

Recall that the  $A_\infty$ -equations for  $A_\infty$ -morphisms are

$$\partial(\text{diagram}) = \sum_{\substack{h+k=n+1 \\ 1 \leq i \leq k \\ h \geq 2}} \pm \text{diagram}_1 + \sum_{\substack{i_1 + \dots + i_s = n \\ s \geq 2}} \pm \text{diagram}_2 .$$

The first diagram on the right is a tree with a red root and blue children. The root has a red line labeled 1 and a blue line labeled  $k$ . The blue line has a blue line labeled  $i$  and a blue line labeled  $h$ . The second diagram is a tree with a red root and blue children. The root has a red line labeled 1 and a blue line labeled  $i_1$ . The blue line has a blue line labeled  $i_s$  and a blue line labeled  $1$ . The diagram is a tree with a red root and blue children. The root has a red line labeled 1 and a blue line labeled  $i_1$ . The blue line has a blue line labeled  $i_s$  and a blue line labeled  $1$ .




Figure: The multiplihedra  $J_1$ ,  $J_2$  and  $J_3$  with cells labeled by the operations they define in  $A_\infty$ -Morph




- 1 The  $A_\infty$ -algebra structure on the Morse cochains
- 2  **$A_\infty$ -morphisms between the Morse cochains**
  - $A_\infty$ -morphisms
  - The multiplihedra
  - **$A_\infty$ -morphisms between the Morse cochains**
- 3 Higher morphisms between  $A_\infty$ -algebras ...
- 4 ... and their realization in Morse theory
- 5 Further directions

Consider an additional Morse function  $g$  on the manifold  $M$ .

Our goal is now to construct an  $A_\infty$ -morphism from the Morse cochains  $C^*(f)$  to the Morse cochains  $C^*(g)$ , through a count of moduli spaces of perturbed Morse trees.

## Definition

A *stable two-colored metric ribbon tree* or *stable gauged metric ribbon tree* is defined to be a stable metric ribbon tree together with a length  $\lambda \in \mathbb{R}$ , which is to be thought of as a gauge drawn over the metric tree, at distance  $\lambda$  from its root, where the positive direction is pointing down.





## Definition

For  $n \geq 1$ ,  $\mathcal{CT}_n$  is the *moduli space of stable two-colored metric ribbon trees*. It has a cell decomposition by stable two-colored ribbon tree type,

$$\mathcal{CT}_n = \bigcup_{t_c \in \mathcal{SCRT}_n} \mathcal{CT}_n(t_c) .$$


Allowing again internal edges of metric trees to go to  $+\infty$ , this moduli space  $\mathcal{CT}_n$  can be compactified into a  $(n-1)$ -dimensional CW-complex  $\overline{\mathcal{CT}}_n$ .






## Theorem ([MW10])

*The compactified moduli space  $\overline{\mathcal{CT}}_n$  is isomorphic as a CW-complex to the multiplihedron  $J_n$ .*



The compactified moduli space  $\overline{\mathcal{CT}}_2$  with its cell decomposition by stable two-colored ribbon tree type



The compactified moduli space  $\overline{\mathcal{CT}}_3$  with its cell decomposition by stable two-colored ribbon tree type



## Definition

A *two-colored perturbed Morse gradient tree*  $T_g^{Morse}$  associated to a pair two-colored metric ribbon tree and perturbation data  $(T_g, \mathbb{Y})$  is the data

- (i) for each edge  $f_c$  of  $t_c$  which is above the gauge, of a smooth map

$$D_{f_c} \xrightarrow{\gamma_{f_c}} M ,$$

such that  $\gamma_{f_c}$  is a trajectory of the perturbed negative gradient  $-\nabla f + \mathbb{Y}_{f_c}$ ,


## Definition

(ii) for each edge  $f_c$  of  $t_c$  which is below the gauge, of a smooth map

$$D_{f_c} \xrightarrow{\gamma_{f_c}} M ,$$

such that  $\gamma_{f_c}$  is a trajectory of the perturbed negative gradient  $-\nabla g + \mathbb{Y}_{f_c}$ ,

and such that the endpoints of these trajectories coincide as prescribed by the edges of the two-colored tree type.



## Definition

Let  $\mathbb{Y}_n$  be a smooth choice of perturbation data on the moduli space  $\mathcal{CT}_n$ . Given  $y \in \text{Crit}(g)$  and  $x_1, \dots, x_n \in \text{Crit}(f)$ , we define the moduli spaces

$$\mathcal{CT}_n^{\mathbb{Y}_n}(y; x_1, \dots, x_n) := \left\{ \begin{array}{l} \text{two-colored perturbed Morse gradient trees associated to} \\ (T_g, \mathbb{Y}_{T_g}) \text{ and connecting } x_1, \dots, x_n \text{ to } y \text{ for } T_g \in \mathcal{CT}_n \end{array} \right\}.$$

## Proposition

*Given a generic choice of perturbation data  $\mathbb{Y}_n$ , the moduli spaces  $\mathcal{CT}_n^{\mathbb{Y}_n}(y; x_1, \dots, x_n)$  are orientable manifolds of dimension*

$$\dim(\mathcal{CT}_n(y; x_1, \dots, x_n)) = |y| - \sum_{i=1}^n |x_i| + n - 1 .$$

Given perturbation data  $\mathbb{X}^f$  and  $\mathbb{X}^g$  for the functions  $f$  and  $g$ , by assuming some gluing-compatibility conditions for a choice of perturbation data  $\mathbb{Y}_n$  for all  $n \geq 1$ , the 1-dimensional moduli spaces  $\mathcal{CT}_n^{\mathbb{Y}_n}(y; x_1, \dots, x_n)$  can be compactified into manifolds with boundary whose boundary is modeled on the  $A_\infty$ -equations for  $A_\infty$ -morphisms :

$$\left[ \partial_{Morse}, \text{Diagram} \right] = \sum_{\substack{h+k=n+1 \\ 1 \leq i \leq k \\ h \geq 2}} \pm \text{Diagram}_1 + \sum_{\substack{i_1 + \dots + i_s = n \\ s \geq 2}} \pm \text{Diagram}_2 .$$

The first diagram is a tree with a root node (red) and two children (blue). The root has a red line going down to the left and a blue line going down to the right. The left child has a blue line going down to the left and a red line going down to the right. The right child has a blue line going down to the left and a red line going down to the right. The root is labeled 1, the left child is labeled h, and the right child is labeled k.

The second diagram is a tree with a root node (red) and s children (blue). The root has a red line going down to the left and a blue line going down to the right. The left child has a blue line going down to the left and a red line going down to the right. The right child has a blue line going down to the left and a red line going down to the right. The root is labeled 1, the left child is labeled i\_1, and the right child is labeled i\_s.

## Theorem ([Maz21a])

Let  $\mathbb{X}^f$ ,  $\mathbb{X}^g$  and  $(\mathbb{Y}_n)_{n \geq 1}$  be admissible choices of perturbation data. Defining for every  $n$  the operation  $\mu_n$  as

$$\begin{aligned} \mu_n^{\mathbb{Y}} : C^*(f) \otimes \cdots \otimes C^*(f) &\longrightarrow C^*(g) \\ x_1 \otimes \cdots \otimes x_n &\longmapsto \\ &\sum_{|y| = \sum_{i=1}^n |x_i| + 1 - n} \# \mathcal{CT}_n^{\mathbb{Y}}(y; x_1, \dots, x_n) \cdot y . \end{aligned}$$

they fit into an  $A_\infty$ -morphism  $\mu^{\mathbb{Y}} : (C^*(f), m_n^{\mathbb{X}^f}) \rightarrow (C^*(g), m_n^{\mathbb{X}^g})$ .

Again, we prove in [Maz21a] that this  $A_\infty$ -morphism actually stems from an  $\Omega BAs$ -morphism between the  $\Omega BAs$ -algebras  $C^*(f)$  and  $C^*(g)$ .

We can moreover prove that this  $A_\infty$ -morphism induces an isomorphism between the Morse cohomologies.



- 1 The  $A_\infty$ -algebra structure on the Morse cochains
- 2  $A_\infty$ -morphisms between the Morse cochains
- 3 Higher morphisms between  $A_\infty$ -algebras ...**
- 4 ... and their realization in Morse theory
- 5 Further directions

Considering two  $A_\infty$ -morphisms  $F, G$ , we would like first to determine a notion giving a satisfactory meaning to the sentence " $F$  and  $G$  are homotopic". Then,  $A_\infty$ -homotopies being defined, what is now a good notion of a homotopy between homotopies ? And of a homotopy between two homotopies between homotopies ? And so on.

- 1 The  $A_\infty$ -algebra structure on the Morse cochains
- 2  $A_\infty$ -morphisms between the Morse cochains
- 3 Higher morphisms between  $A_\infty$ -algebras ...
  - $A_\infty$ -homotopies
  - Higher morphisms between  $A_\infty$ -algebras
  - The  $n$ -multiplihedra
- 4 ... and their realization in Morse theory
- 5 Further directions

## Definition

An  $A_\infty$ -homotopy between two  $A_\infty$ -morphisms  $(f_n)_{n \geq 1}$  and  $(g_n)_{n \geq 1}$  is a collection of maps

$$h_n : A^{\otimes n} \longrightarrow B ,$$

of degree  $-n$ , satisfying

$$\begin{aligned} [\partial, h_n] = & g_n - f_n + \sum_{\substack{i_1+i_2+i_3=n \\ i_2 \geq 2}} \pm h_{i_1+1+i_3}(\text{id}^{\otimes i_1} \otimes m_{i_2} \otimes \text{id}^{\otimes i_3}) \\ & + \sum_{\substack{i_1+\dots+i_s+l \\ +j_1+\dots+j_t=n \\ s+1+t \geq 2}} \pm m_{s+1+t}(f_{i_1} \otimes \dots \otimes f_{i_s} \otimes h_l \otimes g_{j_1} \otimes \dots \otimes g_{j_t}) . \end{aligned}$$

In symbolic formalism,

$$\begin{aligned}
 [\partial, \text{[0 < 1]}] &= \text{[1]} - \text{[0]} + \sum \pm \text{[0 < 1]} \\
 &+ \sum \pm \text{[0]} \text{[0]} \text{[0 < 1]} \text{[1]} \text{[1]},
 \end{aligned}$$

where we denote  $\text{[0]}$ ,  $\text{[0 < 1]}$  and  $\text{[1]}$  respectively for the  $f_n$ , the  $h_n$  and the  $g_n$ .

- 1 The  $A_\infty$ -algebra structure on the Morse cochains
- 2  $A_\infty$ -morphisms between the Morse cochains
- 3 Higher morphisms between  $A_\infty$ -algebras ...
  - $A_\infty$ -homotopies
  - Higher morphisms between  $A_\infty$ -algebras
  - The  $n$ -multiplihedra
- 4 ... and their realization in Morse theory
- 5 Further directions

Notation : the top-dimensional face of the  $n$ -simplex  $\Delta^n$  will be written as  $[0 < \cdots < n]$  and its subfaces  $I \subset \Delta^n$  as  $[i_1 < \cdots < i_k]$ .

## Definition ([MS03])

Let  $I$  be a face of  $\Delta^n$ . An *overlapping partition* of  $I$  is a sequence of faces  $(I_\ell)_{1 \leq \ell \leq s}$  of  $I$  such that

- (i) the union of this sequence of faces is  $I$ , i.e.  $\bigcup_{1 \leq \ell \leq s} I_\ell = I$  ;
- (ii) for all  $1 \leq \ell < s$ ,  $\max(I_\ell) = \min(I_{\ell+1})$ .

An overlapping 6-partition for  $[0 < 1 < 2]$  is for instance

$$[0 < 1 < 2] = [0] \cup [0] \cup [0 < 1] \cup [1] \cup [1 < 2] \cup [2] .$$



## Definition ([Maz21b])

A  $n$ -morphism from  $A$  to  $B$  is defined to be a collection of maps  $f_I^{(m)} : A^{\otimes m} \longrightarrow B$  of degree  $1 - m - \dim(I)$  for  $I \subset \Delta^n$  and  $m \geq 1$ , that satisfy

$$\begin{aligned} [\partial, f_I^{(m)}] &= \sum_{j=0}^{\dim(I)} (-1)^j f_{\partial_j I}^{(m)} + \sum_{\substack{i_1 + \dots + i_s = m \\ I_1 \cup \dots \cup I_s = I \\ s \geq 2}} \pm m_s (f_{I_1}^{(i_1)} \otimes \dots \otimes f_{I_s}^{(i_s)}) \\ &\quad + (-1)^{|I|} \sum_{\substack{i_1 + i_2 + i_3 = m \\ i_2 \geq 2}} \pm f_I^{(i_1 + 1 + i_3)} (\text{id}^{\otimes i_1} \otimes m_{i_2} \otimes \text{id}^{\otimes i_3}). \end{aligned}$$

Equivalently and more visually, a collection of maps  $\text{sing}_I$  satisfying

$$[\partial, \text{sing}_I] = \sum_{j=1}^k (-1)^j \text{sing}_{\partial_j^{\text{sing}} I} + \sum_{I_1 \cup \dots \cup I_s = I} \pm \text{sing}_{I_1} \dots \text{sing}_{I_s} + \sum \pm \text{sing}_I.$$

It is straightforward that 0-morphisms then correspond to  $A_\infty$ -morphisms and 1-morphisms correspond to  $A_\infty$ -homotopies.

Overlapping partitions are the collection of faces which naturally arise in the Alexander-Whitney coproduct.

The element  $\Delta_{\Delta^n}(I)$  corresponds to the sum of all overlapping 2-partitions of  $I$ . Iterating  $s$  times  $\Delta_{\Delta^n}$  yields the sum of all overlapping  $(s + 1)$ -partitions of  $I$ .


We point out that these equations naturally stem from the shifted bar construction viewpoint. An  $A_\infty$ -algebra structure on  $A$  is equivalent to a coderivation  $D_A$  on  $\overline{T}(sA)$  such that  $D_A^2 = 0$ . A  $n$ -morphism between  $A_\infty$ -algebras can then be defined as a morphism of dg-coalgebras  $\Delta^n \otimes \overline{T}(sA) \rightarrow \overline{T}(sB)$ , where  $\Delta^n$  is a dg-coalgebra model for the  $n$ -simplex  $\Delta^n$ .

- 1 The  $A_\infty$ -algebra structure on the Morse cochains
- 2  $A_\infty$ -morphisms between the Morse cochains
- 3 Higher morphisms between  $A_\infty$ -algebras ...
  - $A_\infty$ -homotopies
  - Higher morphisms between  $A_\infty$ -algebras
  - The  $n$ -multiplihedra
- 4 ... and their realization in Morse theory
- 5 Further directions

We would like to define a family of polytopes encoding  $n$ -morphisms between  $A_\infty$ -algebras. These polytopes will then be called  *$n$ -multiplihedra*.

We have seen that  $A_\infty$ -morphisms are encoded by the multiplihedra. A natural candidate for  $n$ -morphisms would thus be  $\{\Delta^n \times J_m\}_{m \geq 1}$ .

However,  $\Delta^n \times J_m$  does not fulfill that property as it is. Faces correspond to the data of a face of  $I \subset \Delta^n$ , and of a broken two-colored tree labeling a face of  $J_m$ . This labeling is too coarse, as it does not contain the trees



that appear in the  $A_\infty$ -equations for  $n$ -morphisms.

We prove in [Maz21b] that there exists a thinner polytopal subdivision of  $\Delta^n \times J_m$  encoding the  $A_\infty$ -equations for  $n$ -morphisms between  $A_\infty$ -algebras. We define the  *$n$ -multiplihedra* to be the polytopes  $\Delta^n \times J_m$  endowed with this polytopal subdivision and denote them  $n - J_m$ .

This thinner polytopal subdivision is obtained by lifting the combinatorics of overlapping partitions to the level of the polytopes  $\Delta^n$ .






Figure: The 1-multiplihedron  $\Delta^1 \times J_2$





Figure: The 2-multiplihedron  $\Delta^2 \times J_2$



**Figure:** The 1-multiplihedron  $\Delta^1 \times J_3$

- 1 The  $A_\infty$ -algebra structure on the Morse cochains
- 2  $A_\infty$ -morphisms between the Morse cochains
- 3 Higher morphisms between  $A_\infty$ -algebras ...
- 4 ... and their realization in Morse theory
  - Motivating question
  - $n$ -morphisms between the Morse cochains
- 5 Further directions

Given two Morse functions  $f, g$ , choices of perturbation data  $\mathbb{X}^f$  and  $\mathbb{X}^g$ , and choices of perturbation data  $\mathbb{Y}$  and  $\mathbb{Y}'$ , is  $\mu^{\mathbb{Y}}$  always  $A_\infty$ -homotopic to  $\mu^{\mathbb{Y}'}$ ? I.e., when can the following diagram be filled in the  $A_\infty$  world

$$\begin{array}{ccc}
 & \mu^{\mathbb{Y}} & \\
 C^*(f) & \begin{array}{c} \downarrow \\ \downarrow \\ \downarrow \end{array} & C^*(g) \quad ? \\
 & \mu^{\mathbb{Y}'} &
 \end{array}$$

- 1 The  $A_\infty$ -algebra structure on the Morse cochains
- 2  $A_\infty$ -morphisms between the Morse cochains
- 3 Higher morphisms between  $A_\infty$ -algebras ...
- 4 ... and their realization in Morse theory
  - Motivating question
  - $n$ -morphisms between the Morse cochains
- 5 Further directions

While the spaces parametrizing the perturbation data were the  $\mathcal{T}_m$  (a model for the associahedra  $K_m$ ) and the  $\mathcal{CT}_m$  (a model for the multiplihedra  $J_m$ ), perturbation data will now be parametrized by the  $n$ -multiplihedra  $\Delta^n \times \mathcal{CT}_m$ .

The previous pattern can then be repeated. We consider a  
 $n$ -simplex of perturbation data  $\mathbb{Y}_{\Delta^n, m} = \{\mathbb{Y}_{\delta, m}\}_{\delta \in \mathring{\Delta}^n}$  on  $\mathcal{CT}_m$ .  
 Given  $y \in \text{Crit}(g)$  and  $x_1, \dots, x_m \in \text{Crit}(f)$ , we define the moduli  
 spaces

$$\mathcal{CT}_{\Delta^n, m}^{\mathbb{Y}_{\Delta^n, m}}(y; x_1, \dots, x_m) := \bigcup_{\delta \in \mathring{\Delta}^n} \mathcal{CT}_m^{\mathbb{Y}_{\delta, m}}(y; x_1, \dots, x_m) .$$



Under some generic assumptions on  $\mathbb{Y}_{\Delta^n, m}$ , the moduli space  $\mathcal{CT}_{\Delta^n, m}(y; x_1, \dots, x_m)$  is then an orientable manifold of dimension

$$\dim(\mathcal{CT}_{\Delta^n, m}(y; x_1, \dots, x_m)) = n + m - 1 + |y| - \sum_{i=1}^m |x_i| .$$

Choose perturbation data  $\mathbb{X}^f$  and  $\mathbb{X}^g$  for the functions  $f$  and  $g$  together with perturbation data  $(\mathbb{Y}_{I,m})_{I \subset \Delta^n}^{m \geq 1}$ . By assuming some gluing-compatibility conditions on  $(\mathbb{Y}_{I,m})_{I \subset \Delta^n}^{m \geq 1}$  modeling the combinatorics of overlapping partitions, the 1-dimensional moduli spaces  $\mathcal{CT}_{I,m}^{\mathbb{Y}_{I,m}}(y; x_1, \dots, x_m)$  can be compactified into manifolds with boundary whose boundary is modeled on the  $A_\infty$ -equations for  $n$ -morphisms.

## Theorem ([Maz21b])

Let  $\mathbb{X}^f$ ,  $\mathbb{X}^g$  and  $(\mathbb{Y}_{I,m})_{I \in \Delta^n}^{m \geq 1}$  be generic choices of perturbation data. Defining for every  $m$  the operation  $\mu_I^{(m)}$  as

$$C^*(f) \otimes \cdots \otimes C^*(f) \xrightarrow{\mu_I^{(m)}} C^*(g)$$

$$x_1 \otimes \cdots \otimes x_m \longmapsto \sum_{|y| = \sum_{i=1}^m |x_i| + 1 - m + |I|} \# \mathcal{CT}_{I,m}^{\mathbb{Y}}(y; x_1, \dots, x_m) \cdot y,$$

they fit into a  $n$ -morphism  $\mu_I^{\mathbb{Y}} : (C^*(f), m_n^{\mathbb{X}^f}) \rightarrow (C^*(g), m_n^{\mathbb{X}^g})$ ,  
 $I \in \Delta^n$ .



## Proposition ([Maz21b])

Let  $\mathbb{Y}$  and  $\mathbb{Y}'$  be two admissible choices of perturbation data on the moduli spaces  $\mathcal{CT}_m$ . The  $A_\infty$ -morphisms  $\mu^{\mathbb{Y}}$  and  $\mu^{\mathbb{Y}'}$  are then  $A_\infty$ -homotopic

$$C^*(f) \begin{array}{c} \xrightarrow{\mu^{\mathbb{Y}}} \\ \Downarrow \\ \xrightarrow{\mu^{\mathbb{Y}'}} \end{array} C^*(g) \quad .$$

- 1 The  $A_\infty$ -algebra structure on the Morse cochains
- 2  $A_\infty$ -morphisms between the Morse cochains
- 3 Higher morphisms between  $A_\infty$ -algebras ...
- 4 ... and their realization in Morse theory
- 5 Further directions**

1. It is quite clear that given two compact symplectic manifolds  $M$  and  $N$ , one should be able to construct  $n$ -morphisms between their Fukaya categories  $\text{Fuk}(M)$  and  $\text{Fuk}(N)$  through counts of moduli spaces of quilted disks (see [MWW18] for the  $n = 0$  case).



2. Given three Morse functions  $f_0, f_1, f_2$ , choices of perturbation data  $\mathbb{X}^i$ , and choices of perturbation data  $\mathbb{Y}^{ij}$  defining morphisms

$$\mu^{\mathbb{Y}^{01}} : (C^*(f_0), m_n^{\mathbb{X}^0}) \longrightarrow (C^*(f_1), m_n^{\mathbb{X}^1}) ,$$

$$\mu^{\mathbb{Y}^{12}} : (C^*(f_1), m_n^{\mathbb{X}^1}) \longrightarrow (C^*(f_2), m_n^{\mathbb{X}^2}) ,$$

$$\mu^{\mathbb{Y}^{02}} : (C^*(f_0), m_n^{\mathbb{X}^0}) \longrightarrow (C^*(f_2), m_n^{\mathbb{X}^2}) ,$$

can we construct an  $A_\infty$ -homotopy such that  $\mu^{\mathbb{Y}^{12}} \circ \mu^{\mathbb{Y}^{01}} \simeq \mu^{\mathbb{Y}^{02}}$  through this homotopy ?

That is, can the following diagram be filled in the  $A_\infty$  realm

$$\begin{array}{ccc}
 C^*(f_0) & \xrightarrow{\mu^{\mathbb{Y}01}} & C^*(f_1) \\
 & \searrow \mu^{\mathbb{Y}02} & \downarrow \mu^{\mathbb{Y}12} \quad ? \\
 & & C^*(f_2)
 \end{array}$$

Work in progress, see also [MWW18] for a similar question.







3. Links between the  $n$ -multiplihedra and the 2-associahedra of Bottman (see [Bot19a] and [Bot19b] for instance) ? We may inspect this matter in a near future with Nate Bottman.





Thanks for your attention !

**Acknowledgements** : Alexandru Oancea, Bruno Vallette,  
Jean-Michel Fischer, Guillaume Laplante-Anfossi, Brice Le Grignou,  
Geoffroy Horel, Florian Bertuol, Thomas Massoni, Amiel  
Peiffer-Smadja and Victor Roca Lucio.





## References I

-  Mohammed Abouzaid, *A topological model for the Fukaya categories of plumbings*, J. Differential Geom. **87** (2011), no. 1, 1–80. MR 2786590
-  Hossein Abbaspour and Francois Laudenbach, *Morse complexes and multiplicative structures*, 2018.
-  Nathaniel Bottman, *2-associahedra*, Algebr. Geom. Topol. **19** (2019), no. 2, 743–806. MR 3924177
-  ———, *Moduli spaces of witch curves topologically realize the 2-associahedra*, J. Symplectic Geom. **17** (2019), no. 6, 1649–1682. MR 4057724

## References II

-  J. M. Boardman and R. M. Vogt, *Homotopy invariant algebraic structures on topological spaces*, Lecture Notes in Mathematics, Vol. 347, Springer-Verlag, Berlin-New York, 1973. MR 0420609
-  Kenji Fukaya, *Morse homotopy and its quantization*, Geometric topology (Athens, GA, 1993), AMS/IP Stud. Adv. Math., vol. 2, Amer. Math. Soc., Providence, RI, 1997, pp. 409–440. MR 1470740
-  Michael Hutchings, *Floer homology of families. I*, Algebr. Geom. Topol. **8** (2008), no. 1, 435–492. MR 2443235
-  Thibaut Mazuir, *Higher algebra of  $a_\infty$ -algebras in morse theory i*, 2021, arXiv:2102.06654.

## References III

-  ———, *Higher algebra of  $A_\infty$  and  $\Omega$ BAs-algebras in Morse theory II*, arXiv:2102.08996, 2021.
-  Stephan Mescher, *Perturbed gradient flow trees and  $A_\infty$ -algebra structures in Morse cohomology*, Atlantis Studies in Dynamical Systems, vol. 6, Atlantis Press, [Paris]; Springer, Cham, 2018. MR 3791518
-  James E. McClure and Jeffrey H. Smith, *Multivariable cochain operations and little  $n$ -cubes*, J. Amer. Math. Soc. **16** (2003), no. 3, 681–704. MR 1969208
-  S. Ma'u and C. Woodward, *Geometric realizations of the multiplihedra*, Compos. Math. **146** (2010), no. 4, 1002–1028. MR 2660682

## References IV



S. Ma'u, K. Wehrheim, and C. Woodward,  $A_\infty$  functors for Lagrangian correspondences, *Selecta Math. (N.S.)* **24** (2018), no. 3, 1913–2002. MR 3816496