

## Site-Directed Mutagenesis of Conserved Amino Acids in the Alpha Subunit of Toluene Dioxygenase: Potential Mononuclear Non-Heme Iron Coordination Sites

HAIYAN JIANG, REBECCA E. PARALES, NANCY A. LYNCH, AND DAVID T. GIBSON\*

Department of Microbiology and Center for Biocatalysis and Bioprocessing, The University of Iowa,  
Iowa City, Iowa 52242

Received 16 January 1996/Accepted 28 March 1996

**The terminal oxygenase component of toluene dioxygenase from *Pseudomonas putida* F1 is an iron-sulfur protein (ISP<sub>TOL</sub>) that requires mononuclear iron for enzyme activity. Alignment of all available predicted amino acid sequences for the large ( $\alpha$ ) subunits of terminal oxygenases showed a conserved cluster of potential mononuclear iron-binding residues. These were between amino acids 210 and 230 in the  $\alpha$  subunit (TodC1) of ISP<sub>TOL</sub>. The conserved amino acids, Glu-214, Asp-219, Tyr-221, His-222, and His-228, were each independently replaced with an alanine residue by site-directed mutagenesis. Tyr-266 in TodC1, which has been suggested as an iron ligand, was treated in an identical manner. To assay toluene dioxygenase activity in the presence of TodC1 and its mutant forms, conditions for the reconstitution of wild-type ISP<sub>TOL</sub> activity from TodC1 and purified TodC2 ( $\beta$  subunit) were developed and optimized. A mutation at Glu-214, Asp-219, His-222, or His-228 completely abolished toluene dioxygenase activity. TodC1 with an alanine substitution at either Tyr-221 or Tyr-266 retained partial enzyme activity (42 and 12%, respectively). In experiments with [<sup>14</sup>C]toluene, the two Tyr→Ala mutations caused a reduction in the amount of *cis*-[<sup>14</sup>C]-toluene dihydrodiol formed, whereas a mutation at Glu-214, Asp-219, His-222, or His-228 eliminated *cis*-toluene dihydrodiol formation. The expression level of all of the mutated TodC1 proteins was equivalent to that of wild-type TodC1 as judged by sodium dodecyl sulfate-polyacrylamide gel electrophoresis and Western blot (immunoblot) analyses. These results, in conjunction with the predicted amino acid sequences of 22 oxygenase components, suggest that the conserved motif Glu-X<sub>3-4</sub>-Asp-X<sub>2</sub>-His-X<sub>4-5</sub>-His is critical for catalytic function and the glutamate, aspartate, and histidine residues may act as mononuclear iron ligands at the site of oxygen activation.**

Toluene dioxygenase (TDO), a multicomponent enzyme system in *Pseudomonas putida* F1, initiates the aerobic catabolism of toluene by forming (+)-*cis*-(1*S*,2*R*)-dihydroxy-3-methylcyclohexa-3,5-diene (*cis*-toluene dihydrodiol) (21, 31, 67). TDO transfers electrons from NADH through Reductase<sub>TOL</sub> (56), and ferredoxin<sub>TOL</sub> (57), to the terminal oxygenase component iron-sulfur protein<sub>TOL</sub> (ISP<sub>TOL</sub>) (55). Reduced ISP<sub>TOL</sub> adds dioxygen to the aromatic nucleus to form *cis*-toluene dihydrodiol (Fig. 1). ISP<sub>TOL</sub> has an  $\alpha_2\beta_2$  subunit conformation (55). The presence of a Rieske-type [2Fe-2S] center in each  $\alpha$  subunit (TodC1) is supported by the presence of a characteristic binding motif (11, 40) in the deduced amino acid sequence of *todC1* (70) and recent biochemical analyses of purified TodC1 (30).

Purified ISP<sub>TOL</sub> requires exogenous mononuclear iron (Fe<sup>2+</sup>) for optimum activity (55), and TodC1 is thought to contain the iron-binding site. In related dioxygenase systems, the ISP components of naphthalene and phthalate dioxygenases each contain one atom of iron per  $\alpha\beta$  and  $\alpha$  subunit, respectively (4, 15). Mononuclear nonheme iron centers are believed to be the site of molecular oxygen activation (4, 6, 9, 19, 66), but the mechanism by which iron, dioxygen, and the substrate interact to catalyze monooxygenation or *cis*-dihydroxylation reactions is not understood. Little structural information is available regarding sites that mediate the binding of mononuclear iron in this type of enzyme.

Amino acids that usually function as iron ligands in proteins are cysteine, histidine, aspartate, glutamate, and tyrosine. For

example, histidine and glutamate residues are ligands to the diiron center of methane monooxygenase hydroxylase (5, 48) and to Fe<sup>2+</sup> at the active site in 2,3-dihydroxybiphenyl 1,2-dioxygenase (24). Histidine and tyrosine residues are ligands to Fe<sup>3+</sup> in protocatechuate 3,4-dioxygenase (43).

The  $\alpha$  subunits of all available sequenced terminal oxygenase components contain conserved histidine, tyrosine, glutamate, and aspartate residues near the middle of their amino acid sequences. The two histidine and tyrosine residues seen at His-222, His-228, and Tyr-221 in this region and Tyr-266 in TodC1 have been suggested as mononuclear iron ligands (13, 38, 40), whereas Nakatsu et al. proposed that a conserved aspartate and histidines (corresponding to Asp-219, His-222, and His-228 in TodC1) may be iron ligands in  $\alpha$  subunit alignments that do not include TodC1 (39). In this study, we individually replaced each of the above conserved histidine, tyrosine, glutamate, and aspartate residues in TodC1 with alanine by site-directed mutagenesis and determined the relationship between these mutations and the catalytic function of TDO.

### MATERIALS AND METHODS

**Bacterial strains and growth conditions.** *Escherichia coli* JM109(DE3) (Promega Corp., Madison, Wis.) and BL21(DE3) (54) were used for standard cloning and gene expression. JM109(pDTG613) (70) was used for the purification of TodC2. JM109(pGP1-2), which carries *todC1* on pT7-5 (70) and is designated here JM109(pGP1-2)(pDTG612C), and *P. putida* F106 (17), a *todC2* mutant strain of strain F1, were used as sources of TodC1, ferredoxin<sub>TOL</sub> (TodB), and reductase<sub>TOL</sub> (TodA) in studies on the reconstitution of active ISP<sub>TOL</sub>. In subsequent experiments, JM109(DE3)(pDTG628), constructed as described below, was used as a source of TodB and TodA. Where appropriate for plasmid maintenance, antibiotics were used at the following concentrations: ampicillin, 150  $\mu$ g/ml; tetracycline, 12.5  $\mu$ g/ml; kanamycin, 100  $\mu$ g/ml. *E. coli* cultures were

\* Corresponding author. Phone: (319) 335-7980. Fax: (319) 335-9999.




FIG. 1. Electron flow in the TDO complex.

grown in Luria broth medium (12) or terrific broth medium (33), and strain F106 was grown at 30°C in a mineral salts medium (53) containing 10 mM sodium pyruvate and 0.01% yeast extract, with toluene provided in the vapor phase.

**Construction of plasmids.** Plasmid pALTER-1 (Promega Corp.) contains a multiple cloning site flanked by the opposing SP6 and T7 RNA polymerase promoters. A 1.6-kb *EcoRI-PstI* fragment carrying the *todC1* gene from pDTG601A (70) was cloned into the *EcoRI-PstI* sites of the pALTER-1 vector, and this plasmid, designated pDTG632, was used as a template for mutagenesis.

A 1.7-kb *DdeI-StuI* DNA fragment carrying the *todBA* genes was purified from pDTG601 (69). The fragment was treated with S1 nuclease (49) to form blunt ends and ligated to *SmaI*-digested pK19 (46), forming pDTG627. The *todBA* genes were then excised from pDTG627 by digestion with *EcoRI* and *BamHI* and inserted into pT7-7 (2), which had been digested with the same enzymes. The resulting plasmid, pDTG628, contains the *todBA* genes under the control of the T7 promoter in pT7-7.

**Molecular techniques.** Plasmid DNA was isolated as described previously (33) or by using the Qiagen Midi Kit (Qiagen, Inc., Chatsworth, Calif.). For sequencing, DNA was further purified by being passed twice through a Centricon-100 filter unit (Amicon, Inc., Beverly, Mass.). Restriction digestions were performed as described by the enzyme suppliers (New England Biolabs, Inc., Beverly, Mass., and Promega Corp.). *E. coli* transformations were performed by the method of Hanahan (25). Ligation reactions and agarose gel electrophoresis were performed by standard procedures (49).

**Site-directed mutagenesis.** Mutagenesis of *todC1* in pDTG632 was carried out with the Altered Sites II in vitro mutagenesis system in accordance with the manufacturer's (Promega Corp.) instructions. Each mutagenic oligonucleotide was designed with a silent mutation that generated a new restriction site (Table 1) to facilitate the identification of clones carrying the desired mutation. The nucleotide sequence of the entire *todC1* gene in each mutant was determined. Sequences and phosphorylated oligonucleotides used for mutagenesis were provided by the University of Iowa DNA Facility.

**Expression of cloned genes.** *E. coli* BL21(DE3)(pDTG628), JM109(DE3)(pDTG632), and JM109(DE3) containing pDTG632 mutant derivatives were cultured in Luria broth medium at 30°C. Cultures were induced with 200  $\mu$ M isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG) for 2 h when the turbidity at 600 nm reached 0.7. Cells were harvested by centrifugation and stored at -70°C. Control strain JM109(DE3)(pALTER-1) was subjected to the same procedures as the recombinant organisms.

**Preparation of cell extracts.** Frozen cells were thawed and suspended in 50 mM 2-(*N*-morpholino)ethanesulfonic acid (MES) buffer (pH 6.8) containing 1 mM dithiothreitol. Phenylmethylsulfonyl fluoride and DNase I were added to final concentrations of 1 mM and 20  $\mu$ g/ml, respectively. The cells were broken by two cycles through a chilled French pressure cell. The broken cells were centrifuged at 150,000  $\times g$  at 4°C for 1 h to remove cell debris and membranes. The supernatant was pelleted in liquid nitrogen and stored at -70°C.

**Protein determinations.** Protein concentrations were determined by the method of Bradford (7), with bovine serum albumin as the standard.

**Purification of TodC2.** The small subunit (TodC2) of ISP<sub>TOL</sub>, encoded by pDTG613, was purified by immunoaffinity chromatography. The ligand was a monoclonal antibody (301 $\beta$ ) isolated from a hybridoma expressing anti-TodC2 antibody (35). Crude cell extract (32.1 mg) from IPTG-induced cells of

JM109(pDTG613) was applied to an 8-ml column in 50 mM MES buffer, pH 6.8, containing 1 mM dithiothreitol and 0.2 mM Fe(NH<sub>4</sub>)<sub>2</sub>(SO<sub>4</sub>)<sub>2</sub> · 6H<sub>2</sub>O. The same buffer was used to wash unbound proteins from the column (80 ml at a flow rate of 2 ml/min). The column was then inverted, and the wash procedure was repeated. Proteins were eluted from the column with buffer containing 50% (vol/vol) ethylene glycol and 1.0 M ammonium sulfate. Fractions (0.5 ml) were collected and tested for the presence of TodC2 by Western blotting. Fractions were combined, dialyzed, and concentrated in 50 mM MES buffer, pH 6.8.

**Determination of conditions for reconstitution of ISP<sub>TOL</sub> activity.** In preliminary experiments, TodC1, TodA, and TodB were provided in cell extracts from toluene-induced cells of strain F106. However, TodC1 was expressed at extremely low levels in this strain and was subsequently provided in cell extracts from heat-induced JM109(pGP1-2)(pDTG612C). Reaction mixtures which contained, in 0.40 ml of 50 mM MES buffer, pH 6.8, 7.5  $\mu$ g of purified TodC2, 0.95 mg of cell extract from JM109(pGP1-2)(pDTG612C), 0.79 mg of cell extract from strain F106, and 0.25 mM Fe(NH<sub>4</sub>)<sub>2</sub>(SO<sub>4</sub>)<sub>2</sub> · 6H<sub>2</sub>O were incubated at room temperature for various periods of time. Reactions were initiated by addition of NADH (0.25 mM final concentration) and 20  $\mu$ l of [<sup>14</sup>C]toluene (1.67 mM, 1.1  $\times 10^5$  dpm/ $\mu$ l in methanol). Aliquots (10  $\mu$ l) were applied to thin-layer chromatography (TLC) squares (1.0 cm<sup>2</sup>) and air dried to remove all remaining [<sup>14</sup>C]-toluene, and the amount of nonvolatile *cis*-toluene dihydrodiol formed was determined in a scintillation counter.

**Enzyme assays.** TDO activity was determined by a modification of the procedure described by Wackett (62). Crude cell extract containing TodC1 from JM109(DE3) carrying pDTG632 or its mutant derivatives, a saturating amount of purified TodC2 (3.3  $\mu$ g), and cell extract of IPTG-induced BL21 (DE3)(pDTG628) (1.3 mg containing of TodA and TodB) were added to 50 mM MES buffer, pH 6.8, containing 0.25 mM Fe(NH<sub>4</sub>)<sub>2</sub>(SO<sub>4</sub>)<sub>2</sub> · 6H<sub>2</sub>O. The mixture was preincubated at room temperature for 30 min. Reactions were started by the addition of NADH (0.25 mM final concentration) and 20  $\mu$ l of [<sup>14</sup>C]toluene (19 mM, 10<sup>5</sup> dpm/ $\mu$ l). The total volume of the reaction mixture was 400  $\mu$ l. Aliquots (10  $\mu$ l) were removed after 0.5, 1, and 2 min and mixed with 10  $\mu$ l of quench solution (toluene-methanol ratio, 1:1 [vol/vol]) to stop the reaction, and radioactivity was determined as described above. Reactions were carried out in triplicate. One unit of TDO activity was defined as the amount of protein required to form 1  $\mu$ mol of *cis*-toluene dihydrodiol per min. Specific activity is given as units per milligram of protein in cell extracts containing wild-type TodC1 or mutant forms of TodC1.

**Isolation of anti-TodC1 monoclonal antibody.** Pure ISP<sub>TOL</sub> (35) was separated into its subunits by electrophoresis on a denaturing 15% polyacrylamide gel (32). The region of the gel containing TodC1 was removed, minced with a razor blade, and used to immunize three BALB/c AnNHsd mice (Harlan Sprague Dawley, Indianapolis, Ind.). Subsequent steps followed standard procedures (28), and hybridomas were isolated and screened for the production of anti-TodC1 antibodies as described previously for the 301 $\beta$  antibody for TodC2 (35). One hybridoma secreted a monoclonal antibody that gave a positive Western blot with TodC1. This antibody, designated 301 $\alpha$ , was purified and used to identify wild-type TodC1 and mutant forms of TodC1 in cell extracts as described below.

**Electrophoresis and Western blot analyses.** Proteins in cell extracts prepared from *E. coli* transformants expressing either mutant or wild-type *todC1* genes were separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE; 12% polyacrylamide gels) (32). Protein detection and Western blotting with antibody 301 $\alpha$  were carried out as described previously (35).

**Detection of reaction products.** TLC, followed by autoradiography, was used to detect reaction products. Aliquots (8  $\mu$ l) from the TDO assays described above were applied to the origin of a TLC plate and developed with chloroform-acetone (80:20 [vol/vol]). X-ray films were exposed for 5 days before development.

**Computerized sequence analysis.** Sequence alignments were done with the Pileup program by using a gap weight of 3.5 and a gap length of 0.1 (Wisconsin Sequence Analysis Package; Genetics Computer Group, Madison, Wis.).

## RESULTS

**Purification of TodC2.** Studies on the function of specific amino acids in TodC1 required the development of an in vitro

TABLE 1. Oligonucleotides used for site-directed mutagenesis

Mutation	Mutagenic oligonucleotide <sup>a</sup>	Restriction site
Glu-214→Ala	5'-GGA AAT TCG <u>CCG CTC</u> AGT TTT GCA GC-3'	BsrBI
Asp-219→Ala	5'-TTG <u>CAG CGC TAT</u> GTA CCA TGC CGG GAC-3'	Eco47III
Tyr-221→Ala	5'-GCA GCG ACA TGG <u>CGC ATG CCG</u> GGA CG-3'	SphI
His-222→Ala	5'-GCA GCG ACA TGT ACG <u>CCG CGG</u> GGA CG-3'	SacII
His-228→Ala	5'-GCC GGG ACG ACC TCA <u>GCG CTG</u> TCT GGC ATC CTG-3'	Eco47III
Tyr-266→Ala	5'-GGA AGT GGC TTC <u>GCG GTC</u> GGC GAC CCC A-3'	BsiEI

<sup>a</sup> The underlined bases represent the restriction sites. The base changes are in boldface.





FIG. 2. Purification of TodC2. Fractions obtained from the 301 $\beta$  immunoaffinity column described in Materials and Methods were analyzed by SDS-PAGE (15% polyacrylamide gels). Proteins were detected by staining with Coomassie brilliant blue R250. Lanes: M, molecular mass standards; 1, purified ISP<sub>TOL</sub> (35); 2, fractions 1 to 4; 3, fractions 5 to 9; 4, fractions 10 to 14; 5, fractions 15 to 20; 6, fractions 21 to 26; 7, crude cell extract from IPTG-induced cells of JM109(DE3)(pDTG613).

assay to measure TodC1 activity. This was achieved by first purifying TodC2 by immunoaffinity chromatography (Fig. 2). No attempt was made to optimize the yield of TodC2, and the purest preparation (Fig. 2, lane 6) was used in the following reconstitution experiments.

**Reconstitution of active ISP<sub>TOL</sub> from its TodC1 and TodC2 subunits.** Purified TodC2 was preincubated with cell extracts containing TodC1, TodA, TodB, and ferrous iron as described in Materials and Methods. Temperature, pH, and preincubation times were optimized. Maximum ISP<sub>TOL</sub> activity was observed after a 30-min incubation period and remained unchanged for up to 8 h. In the following experiments with wild-type TodC1 and mutant forms of TodC1, all dioxygenase components were preincubated in the presence of ferrous iron for 30 min prior to the determination of ISP<sub>TOL</sub> activity. In addition, cell extracts from IPTG-induced cells of JM109(DE3)(pDTG628) were used as a source of TodA and TodB since they gave twice the ISP<sub>TOL</sub> activity given by extracts from strain F106 (data not shown).

**Site-directed mutagenesis of conserved amino acids.** The locations of the two proposed iron-binding regions in TodC1 are shown in Fig. 3A. Amino acid sequence alignments of TodC1 and all available related oxygenases were carried out to locate conserved amino acids that might be involved in iron binding. Figure 3B shows the region containing conserved glutamate, aspartate, tyrosine, and histidine residues which are the potential sites for mononuclear iron coordination. Tyr-266, which is not highly conserved throughout the dioxygenases in

A.



B.

			214	219	221	222	228	266													
ADP1 BenA	NWKLTA	E	N	G	A	D	G	Y	H	VSAV	H	WN	Y	AATTQHRKEK	..	QAGDTIRAMSAGSWGKHG	.GGSYGF	E	HGHMLLTQWQ		
pWwO XylX	NWKVQV	E	N	G	A	D	G	Y	H	VSTV	H	WN	Y	AATQQRKLR	..	DAGDDIRAMTASSWGGDG	.GGFYSE	E	NGHQMVMVARWG		
2CBS CbdA	NWKLQI	E	N	G	A	D	G	Y	H	VGSV	H	WN	Y	VATIGRRD	...	RTSDTIRTVDDTWSKKNIGGTYTF	E	HGHMLLTWRLP			
AC1100 TftA1	NWKLQF	E	N	GL	D	F	Y	H	FGST	H	SS	Y	VDILKLRARAVAPDATQTGVKLASEPNEQCGQTFSF	P	RHGSVMWSTFQ	P	RHGSVMWSTFQ				
C18 DoxB	NWKAPA	E	N	FVFG	D	A	Y	H	VGWT	H	AS	S	LRSGESIFSSLAGNAALPPEGAGLQMTSKYSGSMGVL	W	DGYSGVHSADL	W	DGYSGVHSADL				
9816-4 NahAc	NWKAPA	E	N	FVFG	D	A	Y	H	VGWT	H	AS	S	LRSGESIFSSLAGNAALPPEGAGLQMTSKYSGSMGVL	W	DGYSGVHSADL	W	DGYSGVHSADL				
OUS82 PahAc	NWKAPA	E	N	FVFG	D	A	Y	H	VGWT	H	AS	S	LCTGESIFSSLAGNAALPPEGAGLQMTSKYSGSMGVL	W	DGYSGVHSADL	W	DGYSGVHSADL				
G7 NahAc	NWKAPA	E	N	FVFG	D	A	Y	H	VGWT	H	AS	S	LRSGESIFASLAGNAVLPPEGAGLQMTSKYSGSMGVL	W	DGYSGVHSADL	W	DGYSGVHSADL				
DNT DntAc	NWKVFA	E	N	FVFG	D	I	Y	H	IGWT	H	AS	I	LRAGQAFAPLAGNAMLPEPGLTQATTYSGSGIGVS	L	DAYSQVQSADL	L	DAYSQVQSADL				
JS42 NtdAc	NWKFAA	E	N	FVFG	D	I	Y	H	VGWT	H	AA	A	LRAGQSVFSSLAGNAKLPPEGAGLQMTSKYSGSMGLT	W	DYYSGNFSADM	W	DYYSGNFSADM				
LB400 BphA	NWKFAA	E	Q	F	C	S	D	M	Y	H	AGTT	..	S	H	LSGILAGIPPEMDLSQAQIPTKGNQFRAAWGGHGS	G	Y	VDEPGSLLAVM			
KF707 BphA1	NWKFAA	E	Q	F	C	S	D	M	Y	H	AGTM	..	S	H	LSGILAGMPPEMDLSHAQVPVKGNQFRAWGGHGS	G	F	VDEPGMLLAVM			
KKS102 BphA1	NWKFAA	E	Q	F	C	S	D	M	Y	H	AGTM	..	S	H	LSGVLSSLPPEMDLTQVQMSKNGSQFRAAWGGHGS	G	F	INDAAILMAVM			
PpF1 TodC1	NWKFAA	E	Q	F	C	S	D	M	Y	H	AGTT	..	S	H	LSGILAGLPDEMDLADLPPTVKQYRASWGGHGS	G	Y	VGDPNMLLAIM			
P51 TcbAa	NWKFAA	E	Q	F	C	S	D	M	Y	H	AGTT	..	S	H	LSGILAGLPDGVLEADLADLPVTQVKQYRAPWGGHGTG	F	Y	IGEPNLLLAIM			
BE81 BnzA	NWKFAA	E	Q	F	C	S	D	M	Y	H	AGTT	..	S	H	LSGILAGLPDEMDLADLPPTVKQYRASWGGHGS	G	Y	VGDPNMLLAIM			
RsM5 BpdC1	NWKFAA	E	Q	F	C	S	D	M	Y	H	VGTT	..	S	H	LSGLLAGLPDEIDIREVQPTTGIQYASAPWGGHGS	G	Y	IGEMGTLLAVM			
RgP6 BphA1	NWKFAA	E	Q	F	C	S	D	M	Y	H	VGTT	..	S	H	LSGLLAGLPDEIDIREVQPTTGIQYASAPWGGHGS	G	Y	IGEMGTLLAVM			
RHA1 BphA1	NWKFAA	E	Q	F	C	S	D	M	Y	H	AGTT	..	S	H	LSGILAGLPDGVDSLAPLPTTGIQYRATWGGHGS	G	Y	IGDPNLLLAIM			
NMH102-2 Pht3	NWAQIL	E	G	A	I	.	D	S	A	H	SSSL	H	SS	D	FVPARVGGAEATSKNWLRLPSTDKAPRMQVERTSYGFR	Y	AALRRP IQNAA	Y	AALRRP IQNAA		
BR60 CbaA	NWLQNL	E	G	E	I	.	D	T	S	H	FNFL	H	VG	G	LHADEVDPDHP	....	LKYTAQVAPQYLKVTALGTG	Y	AA...QVPAEE		
POB310 PobA	HWLQNL	E	N	I	M	.	D	P	Y	H	VYIL	H	S	..	.....	TFSGIQFAENFKILPRVDFEAVDGGTV	Y	HAWR...DLED	Y	HAWR...DLED	
19151 VanA	CYRLMI	D	N	L	M	.	D	L	T	H	ETVY	H	AS	S	IGQKEI	..	DEAPVSTRVEGRDLITGRFMEGILAPPFW	R	AALRNGNLADD	R	AALRNGNLADD

FIG. 3. (A) Schematic representation of TodC1 indicating the location of the Rieske [2Fe-2S]-binding site and the potential mononuclear-iron-binding site. (B) Amino acid sequence alignments of the potential mononuclear-iron-binding region. Conserved or semiconserved residues which have been mutated in TodC1 by site-directed mutagenesis are in boxes. The  $\alpha$  subunits of the following dioxygenases are aligned: ADP1 BenA, benzoate dioxygenase from *Acinetobacter calcoaceticus* (40); pWWO XylX, toluate dioxygenase from *P. putida* mt-2 carrying TOL plasmid pWWO (26); 2CBS CbdA, 2-halobenzoate 1,2-dioxygenase from *P. cepacia* 2CBS (22); AC1100 TftA1, 2,4,5-trichlorophenoxyacetic acid oxygenase from *P. cepacia* AC1100 (10); C18 DoxB, naphthalene dioxygenase from *Pseudomonas* sp. (14); 9816-4 NahAc, naphthalene dioxygenase from *Pseudomonas* sp. strain NCIB 9816-4 (44); OUS82 PahAc, polycyclic aromatic hydrocarbon dioxygenase from *P. putida* OUS82 (61); G7 NahAc, naphthalene dioxygenase from *Pseudomonas* sp. strain DNT (59); JS42 NtdAc, 2-nitrotoluene dioxygenase from *Pseudomonas* sp. strain JS42 (44); LB400 BphA, biphenyl dioxygenase from *Pseudomonas* sp. strain LB400 (16); KF707 BphA1, biphenyl dioxygenase from *Pseudomonas* sp. strain KF707 (60); KKS102 BphA1, biphenyl dioxygenase from *Pseudomonas* sp. strain KKS102 (18); PpF1 TodC1, toluene dioxygenase from *P. putida* F1 (69); P51 TcbAa, chlorobenzene dioxygenase from *Pseudomonas* sp. strain P51 (65); BE81 BnzA, benzene dioxygenase from *P. putida* BE-81 (29); RsM5 BpdC1, biphenyl dioxygenase from *Rhodococcus* sp. strain M5 (63); RgP6 BphA1, biphenyl dioxygenase from *Rhodococcus globulus* P6 (1); RHA1 BphA1, biphenyl dioxygenase from *Rhodococcus* sp. strain RHA1 (37); NMH102-2 Pht3, phthalate dioxygenase from *P. putida* carrying plasmid NMH102-2 (41); BR60 CbaA, 3-chlorobenzoate-3,4-dioxygenase from *Alcaligenes* sp. strain BR60 (39); POB310 PobA, phenoxylbenzoate dioxygenase from *P. pseudocaligenes* POB310 (13); 19151 VanA, vanillate demethylase from *Pseudomonas* sp. strain ATCC 19151 (8). All numbers refer to amino acid positions in TodC1.




FIG. 4. (A) SDS-PAGE analysis of wild-type and mutant TodC1 proteins. The gel was stained with Coomassie brilliant blue R250. Lanes: 1, purified ISP<sub>TOL</sub> (35); 2 to 9, *E. coli* cell extracts containing wild-type or mutant TodC1; 2, wild-type TodC1; 3, Glu-214→Ala; 4, Asp-219→Ala; 5, Tyr-221→Ala; 6, His-222→Ala; 7, His-228→Ala; 8, Tyr-266→Ala; 9, cell extract from JM109(DE3) carrying the pALTER-1 vector with no *todC1* gene insert. (B) Western blot analysis of an exact replica of the SDS-PAGE gel. Monoclonal antibody 301α was used to detect the TodC1 polypeptide.

this alignment, was proposed as an iron ligand on the basis of different alignments of selected sequences (13, 40). To determine whether amino acids Glu-214, Asp-219, Tyr-221, His-222, His-228, and Tyr-266 are essential for catalytic activity, we changed each of them independently to alanine by site-directed mutagenesis. The six individual mutants were isolated, and the position of the alanine substitution was verified by determination of the entire nucleotide sequence of each mutant *todC1* gene.

**Expression of mutant and wild-type *todC1* genes.** The wild-type and mutant *todC1* genes were each expressed separately in JM109(DE3) under the control of the T7 promoter in the vector pALTER-1. High-level expression of soluble TodC1 proteins from both the wild-type and mutant *todC1* genes was achieved. As shown in Fig. 4A, SDS-PAGE analysis of crude cell extracts of the wild type and the mutants revealed equivalent levels of full-length TodC1, corresponding to the calculated 50.9-kDa molecular mass. No band was observed for negative control strain JM109(DE3)(pALTER-1). The Western blot (Fig. 4B) with monoclonal antibody 301α identified TodC1 in the crude cell extracts. It also showed that the wild type and all of the mutants expressed comparable amounts of TodC1. In the lane containing purified ISP<sub>TOL</sub> (Fig. 4B), several minor bands are apparent. These may be due to the proteolysis and aggregation of TodC1 which occurred during the purification process.

**TDO activities with wild-type and mutant TodC1 proteins.** *E. coli* crude cell extracts containing wild-type and mutant TodC1 were each analyzed for TDO activity in the presence of TodA, TodB, and TodC2 as described in Materials and Methods. As shown in Table 2, the crude wild-type TodC1 protein, when reconstituted with TodC2, gave a TDO specific activity of 469 mU mg<sup>-1</sup>, which compares favorably with the activity of 2,580 mU mg<sup>-1</sup> recently reported for purified ISP<sub>TOL</sub> (35). When amino acids Glu-214, Asp-219, His-222, and His-228 were individually mutated to alanine, no activity was detected,

TABLE 2. TDO activities obtained with wild-type and mutant TodC1 proteins<sup>a</sup>

Mutation in TodC1	Mean sp act <sup>b</sup> (mU mg <sup>-1</sup> ) ± SD	Relative activity (%)
None (wild-type)	469 ± 74	100
Glu-214→Ala	— <sup>c</sup>	—
Asp-219→Ala	—	—
Tyr-221→Ala	196 ± 19	42
His-222→Ala	—	—
His-228→Ala	—	—
Tyr-266→Ala	58 ± 8	12

<sup>a</sup> Enzyme assays were carried out in triplicate, at three time points, as described in Materials and Methods. Activities reported are the means of nine determinations.

<sup>b</sup> Units per milligram of protein in cell extract from either the wild type or mutant *todC1* clones.

<sup>c</sup> —, activity was below detectable levels (<5% of wild-type activity).

while mutations at both Tyr-221 and Tyr-266 resulted in partial TDO activity.

**Detection of reaction products.** To confirm the results in Table 2, TLC, followed by autoradiography, was used to show that toluene is oxidized to *cis*-toluene dihydrodiol (Fig. 5). The minor amounts of *o*-cresol detected are due to nonenzymatic dehydration of the dihydrodiol (21). Replacement of Tyr-221 and Tyr-266 in TodC1 reduced, but did not eliminate, the formation of *cis*-toluene dihydrodiol, whereas replacement of Glu-214, Asp-219, His-222, and His-228 completely eliminated dihydrodiol formation.

## DISCUSSION

The catabolism of aromatic hydrocarbons and related compounds by aerobic bacteria is often initiated by multicompo-




FIG. 5. Autoradiogram of the products formed from [<sup>14</sup>C]toluene by TDO with wild-type and mutant TodC1 proteins in the TDO assay system described in Materials and Methods. Lanes: 1, wild-type TodC1 cell extract; 2, Glu-214→Ala; 3, Asp-219→Ala; 4, Tyr-221→Ala; 5, His-222→Ala; 6, His-228→Ala; 7, Tyr-266→Ala; 8, cell extract from JM109(DE3)(pALTER-1). Arrows: o, origin of TLC plate; sf, solvent front; A, *o*-cresol; B, *cis*-toluene dihydrodiol.

nent oxygenases. More than 23 oxygenase systems have been identified by protein purification and/or predicted amino acid sequence analysis. The dioxygenases have been divided into three classes with subgroups based on the number and nature of the electron transport components in each system (3). A comparison of the terminal oxygenase components of dioxygenase and monooxygenase systems reveals one class that consists of dissimilar subunits,  $\alpha$  and  $\beta$ , which form an  $\alpha_2\beta_2$  or  $\alpha_3\beta_3$  conformation, depending on the enzyme. Toluene (55), naphthalene (15), biphenyl (16, 23), benzene (68), benzoate (66), and toluate (26) dioxygenases are examples of enzymes that belong to this group. In a second class, the terminal oxygenase components consist of identical  $(\alpha)_n$  subunits, where  $n = 2, 3$ , or 4. This group includes phthalate 4,5-dioxygenase (4), 4-methoxybenzoate monooxygenase (6), vanillate demethylase (8), 4-chlorophenylacetate 3,4-dioxygenase (36), 3-chlorobenzoate 3,4-dioxygenase (39), and 4-sulfobenzoate 3,4-dioxygenase (34). Extensive biophysical studies on phthalate 4,5-dioxygenase (19 and references cited therein), 4-methoxybenzoate monooxygenase (64), benzene dioxygenase (20), and benzoate dioxygenase (66) have shown that each of the  $\alpha$  subunits in these enzymes contains a Rieske  $[2Fe-2S]$  cluster (47) and mononuclear iron. The latter is believed to be the site of dioxygen activation and catalysis.

A comparison of the predicted amino acid sequences for the  $\alpha$  subunits in the  $(\alpha\beta)_n$  and  $(\alpha)_n$  oxygenases reveals a great deal of amino acid conservation within each group, while there is little overall conservation between the two groups. The exceptions are at the Rieske center-binding site (11, 40), where the motif Cys-X-His-X<sub>16-17</sub>-Cys-X<sub>2</sub>-His is conserved in all of the strains listed in Fig. 3B and in the region we propose to contain the mononuclear-iron-binding site, where the common motif Glu-X<sub>3-4</sub>-Asp-X<sub>2</sub>-His-X<sub>4-5</sub>-His is seen in both classes of terminal oxygenase components (Fig. 3B). To investigate the putative role of these amino acids in TDO activity, we developed an assay that involves the reconstitution of active ISP<sub>TOL</sub> from separate sources of its TodC1 and TodC2 subunits. This assay was essential for determination of the activity of mutant forms of TodC1. Similar experiments have been reported for the reconstitution of the terminal oxygenase component of naphthalene dioxygenase (58). However, TodC2 is the first small subunit of an  $(\alpha\beta)_n$  oxygenase to be purified and in addition to its importance in the present work, it will provide the basis for future studies on the function of the small subunit in ISP<sub>TOL</sub> activity. Harayama et al. (27) have suggested that the isofunctional small subunit in toluate 1,2-dioxygenase plays an important role in the determination of substrate specificity.

Results of site-directed mutagenesis experiments indicate that replacement of the Glu-214, Asp-219, His-222, and His-228 residues of TodC1 with alanine totally abolished TDO activity. However, alanine substitution at either of the two semiconserved tyrosines (Tyr-221→Ala and Tyr-266→Ala) resulted in proteins that retain partial TDO activity. Results of *cis*-toluene dihydrodiol formation (Fig. 5) were consistent with the results of enzyme assays in that TodC1 proteins with mutations at Glu-214, Asp-219, His-222, and His-228 formed no dihydrodiol, while the protein with mutations at Tyr-221 and Tyr-266 formed reduced amounts of this product. All mutant *todC1* genes expressed high levels of the full-length TodC1 protein (Fig. 4), indicating that loss or reduction of TDO activity was not due to truncated, unstable, or reduced levels of TodC1. Taken together, these results demonstrate that Glu-214, Asp-219, His-222, and His-228 are essential for catalytic activity and may act as mononuclear iron ligands. Iron is very loosely bound to ISP<sub>TOL</sub>, and significant enzyme activity is observed only when ferrous iron is added to the assay compo-

nents (67). Thus, we cannot unequivocally state that these amino acids are involved in iron coordination. However, the role of glutamate, aspartate, and histidine residues in the coordination of mononuclear iron has been well documented for several enzymes, including the catechol ring cleavage enzymes, catechol 2,3-dioxygenase (51), 2,3-dihydroxybiphenyl 1,2-dioxygenase (24, 50), and protocatechuate 3,4-dioxygenase (43). Thus, our results are consistent with the role of these amino acids as mononuclear iron ligands in TodC1. The results are also consistent with the suggestion that the conserved motif Asp-X<sub>2</sub>-His-X<sub>4</sub>-His in the deduced amino acid sequences of three  $(\alpha)_n$  oxygenases may serve as a binding site for mononuclear iron (39). Tyrosines at Tyr-221 and Tyr-266 in TodC1 were proposed as iron ligands (13, 38, 40) on the basis of alignments with fewer sequences than those shown in Fig. 3B. Although Tyr-221 is conserved in all  $(\alpha\beta)_n$  oxygenases, it is not present in three of the  $(\alpha)_n$  oxygenases (39) (Fig. 3B). In addition, Tyr-266 is only semiconserved in the alignments shown in Fig. 3B. Replacement of either of these two tyrosines with alanine resulted in proteins that retain partial TDO activity and are thus unlikely to interact directly with mononuclear iron. It is possible that the alanine substitution at Tyr-221 may distort the iron-binding site since tyrosine is adjacent to the essential histidine at position 222 in TodC1. Tyr-266 clearly plays an important role in ISP<sub>TOL</sub> activity, since its replacement with alanine leads to a dramatic loss in the amount of toluene converted to *cis*-toluene dihydrodiol. In some sequences, Tyr-266 is replaced with phenylalanine or tryptophan, suggesting that there may be a structural requirement for an aromatic amino acid in this region.

Recent magnetic circular dichroism studies of the dioxygen-activating site of phthalate dioxygenase from *P. cepacia* (terminal oxygenase has an  $\alpha_4$  conformation) indicate that there is one six-coordinate Fe<sup>2+</sup> site. Substrate binding converts the site to a five-coordinate species, opening up a coordination position for O<sub>2</sub> binding (19, 45). However, no data are available that identify the coordinating amino acid residues. Here we present evidence that in the related TDO system, Glu-214, Asp-219, His-222, and His-228 may be at the dioxygen activation site and play a role as mononuclear iron ligands. These amino acids are also contained in the motif Glu-X<sub>3-4</sub>-Asp-X<sub>2</sub>-His-X<sub>4-5</sub>-His, which is conserved in all oxygenase sequences that have been reported to date. The only exception is the presumably allowable substitution of aspartate for glutamate in vanillate demethylase (Fig. 3B). It will be of interest to determine whether the remaining ligands are amino acid residues or water molecules. The latter have been shown to coordinate metals in certain proteins (42, 43, 48). Although the amino acid residues that act as mononuclear iron ligands in dioxygenase  $\alpha$  subunits have been proposed for several years on the basis of sequence alignments (13, 38–40), this is the first study that has provided experimental evidence of the importance of these individual residues by replacing each with an alanine. The development of a reconstitution assay for ISP<sub>TOL</sub> with purified TodC2 paves the way for further studies of the role of Glu-214, Asp-219, His-222, and His-228 in TDO activity. To establish unequivocally that these amino acids are involved in the coordination of mononuclear iron, the individual TodC1 mutant proteins are being purified and the role of iron in catalysis will be determined by appropriate spectroscopic procedures (19, 45, 51). In addition, changing the above amino acids to residues other than alanine will provide more insight into the oxygen fixation reactions catalyzed by these unique enzymes.

## ACKNOWLEDGMENTS

This work was supported by U.S. Public Health Service grant GM29909 from the National Institute of General Medical Sciences.

We thank Kyoung Lee, John D. Haddock, and Mark Urbanowski for helpful discussions and Julie Neilson for assistance in preparing the manuscript.

## REFERENCES

- Asturias, J. A., E. Díaz, and K. N. Timmis. 1995. The evolutionary relationship of biphenyl dioxygenase from Gram-positive *Rhodococcus globerulus* P6 to multicomponent dioxygenase from Gram-negative bacteria. *Gene* **156**:11–18.
- Ausubel, F. M., R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A. Smith, and K. Struhl. 1993. Current protocols in molecular biology. John Wiley & Sons, Inc., New York.
- Batie, C. J., D. P. Ballou, and C. C. Corell. 1991. Phthalate dioxygenase reductase and related flavin-iron-sulfur containing electron transferases, p. 543–556. In F. Müller (ed.), *Chemistry and biochemistry of flavoenzymes*. CRC Press, Inc., Boca Raton, Fla.
- Batie, C. J., E. LaHaie, and D. P. Ballou. 1987. Purification and characterization of phthalate oxygenase and phthalate oxygenase reductase from *Pseudomonas cepacia*. *J. Biol. Chem.* **262**:1510–1518.
- Bender, C. J., A. C. Rosenzweig, S. J. Lippard, and J. Peisach. 1994. Nuclear hyperfine coupling of nitrogen in the coordination sphere of the diiron center of methane monooxygenase hydroxylase. *J. Biol. Chem.* **269**:15993–15998.
- Bill, E., F.-H. Bernhardt, and A. X. Trautwein. 1981. Mössbauer studies on the active Fe...[2Fe-2S] site of putidamonooxigenase, its electron transport and dioxygen activation mechanism. *Eur. J. Biochem.* **121**:39–46.
- Bradford, M. M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**:248–254.
- Brunel, F., and J. Davison. 1988. Cloning and sequencing of *Pseudomonas* genes encoding vanillate demethylase. *J. Bacteriol.* **170**:4924–4930.
- Crutcher, S. E., and P. J. Geary. 1979. Properties of the iron-sulphur proteins of the benzene dioxygenase system from *Pseudomonas putida*. *Biochem. J.* **177**:393–400.
- Danganan, C. E., R. W. Ye, D. L. Daubaras, L. Xun, and A. M. Chakrabarty. 1995. Nucleotide sequence and functional analysis of the genes encoding 2,4,5-trichlorophenoxyacetic acid oxygenase in *Pseudomonas cepacia* AC1100. *Appl. Environ. Microbiol.* **60**:4100–4106.
- Davidson, M., T. Ohnishi, E. Atta-Asofo-Adjei, and F. Daldal. 1992. Potential ligands to the [2Fe-2S] Rieske cluster of the cytochrome *bcl* complex of *Rhodobacter capsulatus* probed by site-directed mutagenesis. *Biochemistry* **31**:3342–3351.
- Davis, R. W., D. Botstein, and J. R. Roth. 1980. Advanced bacterial genetics. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- Dehmel, U., K.-H. Engesser, K. N. Timmis, and D. F. Dwyer. 1995. Cloning, nucleotide sequence, and expression of the gene encoding a novel dioxygenase involved in metabolism of carboxydiphenyl ethers in *Pseudomonas pseudoalcaligenes* POB310. *Arch. Microbiol.* **163**:35–41.
- Denome, S. A., D. C. Stanley, E. S. Olsen, and K. D. Young. 1993. Metabolism of dibenzothiophene and naphthalene in *Pseudomonas* strains: complete DNA sequence of an upper naphthalene catabolic pathway. *J. Bacteriol.* **175**:6890–6901.
- Ensley, B. D., and D. T. Gibson. 1983. Naphthalene dioxygenase: purification and properties of a terminal oxygenase component. *J. Bacteriol.* **155**:505–511.
- Erickson, B. D., and F. J. Mondello. 1992. Nucleotide sequencing and transcriptional mapping of the genes encoding biphenyl dioxygenase, a multicomponent polychlorinated biphenyl-degrading enzyme in *Pseudomonas* strain LB400. *J. Bacteriol.* **174**:2903–2912.
- Finette, B. A., V. Subramanian, and D. T. Gibson. 1984. Isolation and characterization of *Pseudomonas putida* PpF1 mutants defective in the toluene dioxygenase enzyme system. *J. Bacteriol.* **160**:1003–1009.
- Fukuda, M., Y. Yasukochi, Y. Kikuchi, Y. Nagata, K. Kimbara, H. Horiuchi, M. Takagi, and K. Yano. 1994. Identification of the *bphA* and *bphB* genes of *Pseudomonas* sp. strain KKS102 involved in degradation of biphenyl and polychlorinated biphenyls. *Biochem. Biophys. Res. Commun.* **202**:850–856.
- Gassner, G. T., D. P. Ballou, G. A. Landrum, and J. W. Whittaker. 1993. Magnetic circular dichroism studies on the mononuclear ferrous active site of phthalate dioxygenase from *Pseudomonas cepacia* show a change of ligation state on substrate binding. *Biochemistry* **32**:4820–4825.
- Geary, P. J., F. Saboowalla, D. Patil, and R. Cammack. 1984. An investigation of the iron-sulphur proteins of benzene dioxygenase from *Pseudomonas putida* by electron-spin-resonance spectroscopy. *Biochem. J.* **217**:667–673.
- Gibson, D. T., M. Hensley, H. Yoshioka, and T. J. Mabry. 1970. Formation of (+)-cis-2,3-dihydroxy-1-methylcyclohexa-4,6-diene from toluene by *Pseudomonas putida*. *Biochemistry* **9**:1626–1630.
- Haak, B., S. Fetzner, and F. Lingens. 1995. Cloning, nucleotide sequence, and expression of the plasmid-encoded genes for the two-component 2-halo-benzoate 1,2-dioxygenase from *Pseudomonas cepacia* 2CBS. *J. Bacteriol.* **177**:667–675.
- Haddock, J. D., and D. T. Gibson. 1995. Purification and characterization of the oxygenase component of biphenyl 2,3-dioxygenase from *Pseudomonas* sp. strain LB400. *J. Bacteriol.* **177**:5834–5839.
- Han, S., L. D. Eltis, K. N. Timmis, S. W. Muchmore, and J. T. Bolin. 1995. Crystal structure of biphenyl-cleaving extradiol dioxygenase from a PCB-degrading pseudomonad. *Science* **270**:976–980.
- Hanahan, D. 1983. Studies on transformation of *Escherichia coli* with plasmids. *J. Mol. Biol.* **166**:557–580.
- Harayama, S., M. Rekik, A. Bairock, E. L. Neidle, and L. N. Ornston. 1991. Potential DNA slippage structure acquired during evolutionary divergence of *Acinetobacter calcoaceticus* chromosomal *benABC* and *Pseudomonas putida* TOL pWWO plasmid *xytXYZ*, genes encoding benzoate dioxygenases. *J. Bacteriol.* **173**:7540–7548.
- Harayama, S., M. Rekik, and K. N. Timmis. 1986. Genetic analysis of a relaxed substrate specificity aromatic ring dioxygenase, toluate 1,2-dioxygenase, encoded by TOL plasmid pWWO of *Pseudomonas putida*. *Mol. Gen. Genet.* **202**:226–234.
- Harlow, E., and D. Lane. 1988. Antibodies: a laboratory manual. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- Irie, S., S. Doi, T. Yorifuji, M. Takagi, and K. Yano. 1987. Nucleotide sequencing and characterization of the genes encoding benzene oxidation enzymes of *Pseudomonas putida*. *J. Bacteriol.* **169**:5174–5179.
- Jiang, H., and D. T. Gibson. Unpublished data.
- Kobal, V. M., D. T. Gibson, R. E. Davis, and A. Garza. 1973. X-ray determination of the absolute stereochemistry of the initial oxidation product formed from toluene by *Pseudomonas putida* 39/D. *J. Am. Chem. Soc.* **95**:4420–4421.
- Laemmli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature (London)* **227**:680–685.
- Lee, S.-Y., and S. Rasheed. 1990. A simple procedure for maximum yield of high-quality plasmid DNA. *Bio/Techniques* **9**:676–679.
- Locher, H. H., T. Leisinger, and A. M. Cook. 1991. 4-Sulphobenzoate 3,4-dioxygenase: purification and properties of a desulphonative two-component enzyme system from *Comamonas testosteroni* T-2. *Biochem. J.* **274**:833–842.
- Lynch, N. A., H. Jiang, and D. T. Gibson. Rapid purification of the oxygenase component of toluene dioxygenase from a polyol-responsive monoclonal antibody. *Appl. Environ. Microbiol.*, in press.
- Markus, A., D. Krekel, and F. Lingens. 1986. Purification and some properties of component A of the 4-chlorophenylacetate 3,4-dioxygenase from *Pseudomonas* species strain CBS. *J. Biol. Chem.* **261**:12883–12888.
- Masai, E., A. Yamada, J. Healy, T. Hatta, K. Kimbara, M. Fukuda, and K. Yano. 1995. Characterization of biphenyl catabolic genes of gram-positive polychlorinated biphenyl degrader *Rhodococcus* sp. strain RHA1. *Appl. Environ. Microbiol.* **61**:2079–2085.
- Mason, J. R., and R. Cammack. 1992. The electron-transport proteins of hydroxylating bacterial dioxygenases. *Annu. Rev. Microbiol.* **46**:277–305.
- Nakatsu, C. H., N. A. Straus, and R. C. Wyndham. 1995. The nucleotide sequence of the Tn5271 3-chlorobenzoate 3,4-dioxygenase genes (*cbaAB*) unites the class IA oxygenases in a single lineage. *Microbiology* **141**:485–495.
- Neidle, E. L., C. Hartnett, L. N. Ornston, A. Bairock, M. Rekik, and S. Harayama. 1991. Nucleotide sequences of the *Acinetobacter calcoaceticus* *benABC* genes for benzoate 1,2-dioxygenase reveal evolutionary relationships among multicomponent oxygenases. *J. Bacteriol.* **173**:5385–5395.
- Nomura, Y., M. Nakagawa, N. Ogawa, S. Harashima, and Y. Oshima. 1992. Genes in PHT plasmid encoding the initial degradation pathway of phthalate in *Pseudomonas putida*. *J. Ferment. Bioeng.* **6**:333–344.
- Nordlund, P., B.-M. Sjöberg, and H. Eklund. 1990. Three-dimensional structure of the free radical protein of ribonucleotide reductase. *Nature (London)* **345**:593–598.
- Ohlendorf, D. H., A. M. Orville, and J. D. Lipscomb. 1994. Structure of protocatechuate 3,4-dioxygenase from *Pseudomonas aeruginosa* at 2.15 Å resolution. *J. Mol. Biol.* **244**:586–608.
- Parales, J. V., A. Kumar, R. E. Parales, and D. T. Gibson. Cloning and sequencing of the genes encoding 2-nitrotoluene dioxygenase from *Pseudomonas* sp. JS42. Submitted for publication.
- Pavel, E. G., L. J. Martins, W. R. J. Ellis, and E. I. Solomon. 1994. Magnetic circular dichroism studies of exogenous ligand and substrate binding to the non-heme ferrous active site in phthalate dioxygenase. *Chem. Biol.* **1**:173–183.
- Pridmore, R. D. 1987. New and versatile cloning vectors with kanamycin-resistance marker. *Gene* **56**:309–312.
- Rieske, J. S., D. H. MacLennan, and R. Coleman. 1964. Isolation and properties of an iron-protein from the (reduced coenzyme Q)-cytochrome C reductase complex of the respiratory chain. *Biochem. Biophys. Res. Commun.* **15**:338–344.
- Rosenzweig, A. C., C. A. Frederick, S. J. Lippard, and P. Nordlund. 1993. Crystal structure of a bacterial non-haem iron hydroxylase that catalyzes the biological oxidation of methane. *Nature (London)* **366**:537–543.

49. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
50. Senda, T., K. Sugiyama, H. Narita, T. Yamamoto, K. Kimbara, M. Fukuda, M. Sato, K. Yano, and Y. Mitsui. 1996. Three-dimensional structures of free form and two substrate complexes of an extradiol ring-cleavage type dioxygenase, the BphC enzyme from *Pseudomonas* sp. strain KKS102. *J. Mol. Biol.* **255**:735–752.
51. Shu, L., Y.-M. Chiou, A. M. Orville, M. A. Miller, J. D. Lipscomb, and L. Que, Jr. 1995. X-ray absorption spectroscopic studies of the Fe(II) active site of catechol 2,3-dioxygenase. Implications for the extradiol cleavage mechanism. *Biochemistry* **34**:6649–6659.
52. Simon, M. J., T. D. Osslund, R. Saunders, B. D. Ensley, S. Suggs, A. Harcourt, W.-C. Suen, D. L. Cruden, D. T. Gibson, and G. J. Zylstra. 1993. Sequences of genes encoding naphthalene dioxygenase in *Pseudomonas putida* strains G7 and NCIB 9816-4. *Gene* **127**:31–37.
53. Stanier, R. Y., N. J. Palleroni, and M. Doudoroff. 1966. The aerobic pseudomonads: a taxonomic study. *J. Gen. Microbiol.* **43**:159–271.
54. Studier, F. W., A. H. Rosenberg, J. J. Dunn, and J. W. Dubendorff. 1990. Use of T7 RNA polymerase to direct expression of cloned genes. *Methods Enzymol.* **185**:60–89.
55. Subramanian, V., T.-N. Liu, W.-K. Yeh, and D. T. Gibson. 1979. Toluene dioxygenase: purification of an iron-sulfur protein by affinity chromatography. *Biochem. Biophys. Res. Commun.* **91**:1131–1139.
56. Subramanian, V., T.-N. Liu, W.-K. Yeh, M. Narro, and D. T. Gibson. 1981. Purification and properties of NADH-ferredoxin<sub>TOL</sub> reductase: a component of toluene dioxygenase from *Pseudomonas putida*. *J. Biol. Chem.* **256**:2723–2730.
57. Subramanian, V., T.-N. Liu, W.-K. Yeh, C. M. Serdar, L. P. Wackett, and D. T. Gibson. 1985. Purification and properties of ferredoxin<sub>TOL</sub>: a component of toluene dioxygenase from *Pseudomonas putida* F1. *J. Biol. Chem.* **260**:2355–2363.
58. Suen, W.-C., and D. T. Gibson. 1994. Recombinant *Escherichia coli* strains synthesize active forms of naphthalene dioxygenase and its individual  $\alpha$  and  $\beta$  subunits. *Gene* **143**:67–71.
59. Suen, W.-C., and J. C. Spain. Unpublished data.
60. Taira, K., J. Hirose, S. Hayashida, and K. Furukawa. 1992. Analysis of *bph* operon from the polychlorinated biphenyl-degrading strain of *Pseudomonas pseudoalcaligenes* KF707. *J. Biol. Chem.* **267**:4844–4853.
61. Takizawa, N., N. Kaida, S. Torigoe, T. Moritani, T. Sawada, S. Satoh, and H. Kiyohara. 1994. Identification and characterization of genes encoding polycyclic aromatic hydrocarbon dioxygenase and polycyclic aromatic hydrocarbon dihydrodiol dehydrogenase in *Pseudomonas putida* OUS82. *J. Bacteriol.* **176**:2444–2449.
62. Wackett, L. P. 1990. Toluene dioxygenase from *Pseudomonas putida* F1. *Methods Enzymol.* **188**:39–45.
63. Wang, Y., J. Garnon, D. Labbe, H. Bergeron, and P. C. K. Lau. 1995. Sequence and expression of the *bpdC1C2BADE* genes involved in the initial steps of biphenyl chlorobiphenyl degradation by *Rhodococcus* sp. M5. *Gene* **164**:117–122.
64. Wende, P., F.-H. Bernhardt, and K. Pflieger. 1989. Substrate-modulated reactions of putidamonooxin: the nature of the active oxygen species formed and its reaction mechanism. *Eur. J. Biochem.* **181**:189–197.
65. Werlen, C., H.-P. Kohler, and J. R. van der Meer. 1996. The broad substrate chlorobenzene dioxygenase and *cis*-chlorobenzene dihydrodiol dehydrogenase of *Pseudomonas* sp. strain P51 are linked evolutionarily to the enzymes for benzene and toluene degradation. *J. Biol. Chem.* **271**:4009–4016.
66. Yamaguchi, M., and H. Fujisawa. 1982. Subunit structure of oxygenase component in benzoate-1,2-dioxygenase system from *Pseudomonas arvilla* C-1. *J. Biol. Chem.* **257**:12497–12502.
67. Yeh, W.-K., D. T. Gibson, and T.-N. Liu. 1977. Toluene dioxygenase: a multicomponent enzyme system. *Biochem. Biophys. Res. Commun.* **78**:401–410.
68. Zamanian, M., and J. R. Mason. 1987. Benzene dioxygenase in *Pseudomonas putida*. *Biochem. J.* **244**:611–616.
69. Zylstra, G. J., and D. T. Gibson. 1989. Toluene degradation by *Pseudomonas putida* F1: nucleotide sequence of the *todC1C2BADE* genes and their expression in *E. coli*. *J. Biol. Chem.* **264**:14940–14946.
70. Zylstra, G. J., and D. T. Gibson. 1991. Aromatic hydrocarbon degradation: a molecular approach, p. 183–203. *In* J. K. Setlow (ed.), *Genetic engineering: principles and methods*. Plenum Press, New York.